

Oxidative β -Cleavage of Fused Cyclobutanols Leading to Hydrofuran-Fused Polycyclic Aromatic Compounds


Hayate Kinouchi, Kazuma Sugimoto, Yousuke Yamaoka, Hiroshi Takikawa and Kiyosei Takasu*

Graduate School of Pharmaceutical Sciences, Kyoto University, Yoshida, Sakyo-ku, Kyoto 606-8501,
Japan.

E-mail: kay-t@pharm.kyoto-u.ac.jp

Fax: +81 75 753 4604; Tel: +81 75 753 4553

Graphic Abstract



Abstract


Treatment of aryl-fused bicyclo[4.2.0]octanols with an oxidant such as phenyliodine diacetate (PIDA) or hypochlorous acid gave dihydrofuran-containing polycyclic aromatic compounds by selective β -cleavage of the cyclobutanol moiety. Mechanistic studies suggest that the oxygen atom of the hydrofuran ring is incorporated from the hydroxy group of the substrate via intramolecular addition. The oxidative transformation should serve as a new method to prepare functionalized polycyclic aromatic compounds.

Introduction

Polycyclic aromatic hydrocarbons (PAHs) exhibit unique optical and electronic characteristics. PAHs have been applied to electronic devices such as organic light-emitting diodes (OLEDs), organic photovoltaics (OPVs), and organic field-effect transistors (OFETs).¹ PAHs are toxic compounds that show carcinogenic and mutagenic effects.² On the other hand, they have potential as medicinal compounds displaying antitumor activity.³ The size and geometry of the aromatic skeletons and incorporation of heteroatoms into the π -system affect the properties of PAHs. Because introducing functional groups or substituents at peripheral positions can control the properties, a variety of synthetic methods have been reported to prepare functionalized/substituted PAHs.⁴

We previously reported an intramolecular (2 + 2) cycloaddition of 2-acyl-2'-alkenyl-1,1'-biaryls **1** to furnish diareno-fused bicyclo[4.2.0]octanols **2**. This reaction is promoted by a potassium base such as KO^tBu and KHMDS (Scheme 1a).⁵ We also demonstrated that an acid-treatment of **2** provides substituted polyaromatic compounds **3** by a domino ring contraction–opening reaction (Scheme 1b).^{5,6} During our extensive studies, we found that an acid-promoted retro (2 + 2) cycloaddition of azapropellanes **4** furnishes π -extended carbazoles **5** (Scheme 1c).⁷ The promotion of these reactions should be driven by alleviating the strain of the cyclobutane rings.⁸ Oxidative carbon-carbon bond cleaving reactions are valuable synthetic tools, which yield structurally diverse organic molecules. In particular, cyclobutanols are representative substrates for bond cleaving reactions at the β -bond of the hydroxy group. The selectivity of the breaking

β -bond is actively researched.^{9,10} However, most of studies have been done on the oxidative β -cleavage of monocyclic cyclobutanols. The examples for fused cyclobutanols are limited.^{10b,d,f,h} We envisage that an oxidative β -cleavage reaction of diareno-fused bicyclo[4.2.0]octanols **2** gives substituted polyaromatic compounds **A** or cyclic biaryl compounds **B**, depending on the breaking C–C bonds (Figure 1). Herein, we describe the synthesis of dihydrofuran-fused PAHs by oxidation of **2** with an unusual regioselective bond cleavage and their further functionalization.



Scheme 1. Previous works: (a) Potassium base promoted intramolecular (2 + 2) cycloaddition, (b) domino reaction of cyclobutanols **2** giving substituted PAHs **3**, (c) retro (2 + 2) cycloaddition of azapropellanes **4** giving tribenzocarbazoles **5**.





Figure 1. This work: Oxidative β -cleavage of diareno-bicyclo[4.2.0]octanols **2**. **A**; A compound resulted in the C(1)–C(2) bond cleavage, **B**; A compound resulted in the C(1)–C(4) bond cleavage.

Results and Discussion

We initially examined the oxidation of **2a** with phenyliodine diacetate (PIDA), which has been used by Fujioka and Kita's group for β -cleavage of monocyclic cyclobutanols to give γ -hydroxyketones.^{10d} It was proposed that the reaction proceeds through a cationic pathway. When using the standard reaction in the literature in 1,1,1,3,3,3-hexafluoropropane-2-ol (HFIP) and H₂O (9 : 1 v/v), **2a** was fully consumed within 1 h to afford **6a** in 15% yield (Table 1, entry 1). ¹H and ¹³C {¹H} NMR spectra confirmed that **6a** had a phenanthrol[9,10-*b*]furan skeleton. We attributed the low yield of **6a** to the overoxidation of the electron-rich phenanthrene ring. Adding PIDA portionwise into the solution of **2a** over 15 min increased the yield of **6a** to 40% (entry 2). With 1.1 equivalents of PIDA, the yield of **2a** was increased into 66% (entry 3).

To suppress the reactivity of PIDA, the solvent was screened. Substituting acidic HFIP as a co-solvent with CH₂Cl₂ or CH₃CN of H₂O dramatically improved the yield of **6a** to 82% and 97%, respectively (entries 4 and 5). A high yield of **6a** was also achieved with 1.1 equivalents of PIDA in CH₃CN–H₂O (entry 6). The transformation of **2a** into **6a** resulted in the cleavage of the C(1)–C(2) bond (type A in Figure 1) followed by C–O bond formation. However, 8-membered biphenyl product (type B), which would be produced by C(1)–C(4) bond cleavage, was not observed. Oxidative β -cleavage reaction of fused cyclobutanols usually gave medium-sized ring products by breaking the transannular bond (type B).^{10b,d,f,h} To the best of our knowledge, this is the first case of type A regioselectivity in the oxidative β -cleavage of fused cyclobutanols.

Table 1. Optimization of the reaction conditions of PIDA-oxidation of **2a**^{a,b}














Entry	X	Solvent	Time (h)	%yield of 6a
1	2.0	HFIP	1	15
2 ^c	2.0	HFIP	0.2	40
3	1.1	HFIP	1	66
4	2.0	CH_2Cl_2	1	82
5	2.0	CH_3CN	1	97
6	1.1	CH_3CN	1	83

^aAll reactions were carried out using 0.10 mmol scale of **2a** (0.10 M). ^bIsolated yield.

^cPIDA was added portionwise over 15 min.

With the optimized conditions (entry 5 in Table 1) in hand, we then examined the substrate scope of the diareno-fused bicyclo[4.2.0]octanols (Table 2). Compounds **2b–c** and **2d**, which have electron-withdrawing and -donating groups on the aromatic ring, respectively, afforded desired phenanthrene-fused dihydrofurans **6b–d** in moderate to high yields (entries 1–3). Substrates bearing heteroaromatic or π -extended rings were applicable to this transformation to furnish dihydrofurans **6e** and **6f** in high yields (entries 4 and 5). Spirocyclic product **6g** was also obtained in high yield from **2g** (entry 6). In contrast, monomethyl substrate **2h** gave a complex mixture of unidentified products (entry 7).

Table 2. Substrate scope of the reaction of **2** with PIDA^{a,b}


Entry	Substrates	Products	%yield of 6
1	 2b (R ¹ = Cl, R ² = H)	 6b	62
2	 2c (R ¹ = H, R ² = Cl)	 6c	81
3	 2d (R ¹ = MeO, R ² = H)	 6d	91
4	 2e	 6e	99
5	 2f	 6f	59
6	 2g	 6g	97
7	 2h	 6h	complex mixture

^aAll reactions were carried out using 0.10 mmol of **2** (0.10 M). PIDA (1.1 equiv.), CH₃CN/H₂O (9:1 v/v), rt. ^bIsolated yield.

Next, we investigated oxidative β -cleavage using hypochlorous acid (HClO), which Singleton's group employed in the radical mediated oxidative cleavage of monocyclic cyclobutanols.^{10g} Treating **2a** with 4 equivalents of NaOCl·5H₂O¹¹ and AcOH, which produced HClO *in situ* in CH₂Cl₂ at 0 °C, gave chlorinated compound **7a** in 41% yield (Table 3, entry 1). The structure of **7a** was determined by X-ray crystallographic analysis. Increasing the reaction temperature of the HClO-oxidation of **2a** improved the yield of **7a** up to 81% yield (entries 2 and 3). The reaction with a stoichiometric amount of the oxidant resulted in a poor yield of **7a** (entry 4). Solvent screening revealed that toluene and CH₃CN gave lower yields of **7a** (entries

5 and 6). **7a** should be formed by the β -cleavage of the cyclobutane ring of **2a** followed by further chlorinative dearomatization (oxidation)¹² of *in situ* generated **6a**. In fact, when oxidation of **6a** was conducted under the same conditions as entry 3, **7a** was obtained in 66% yield.

Table 3. Optimization of the reaction conditions to hemiacetal **7a**^{a,b}










Entry	Solvent	Temp (°C)	% yield of 7a
1	CH ₂ Cl ₂	0	41
2	CH ₂ Cl ₂	rt	66
3	CH ₂ Cl ₂	40	81
4 ^c	CH ₂ Cl ₂	40	25
5	toluene	40	17
6	CH ₃ CN	40	43

^aAll reactions were carried out using 0.20 mmol of **2a** (0.10 M). ^bIsolated yield. ^cNaOCl·5H₂O (1.0 equiv.) and AcOH (1.0 equiv.) were used.

Under the optimal conditions (Table 3, entry 3), the HClO-oxidation of various *tert*-cyclobutanols **2** was conducted (Table 4). Compounds **2b**, **2c** and **2f** respectively afforded hemiacetals **7b**, **7c**, and **7f** in good yields (entries 1, 2, and 5). In contrast, the reactions of compounds **2d** and **2e**, which have an electron-rich aromatic ring, gave a complex mixture of unidentified products (entries 3 and 4). We speculated that the different reaction scope using HClO from using PIDA might be caused by the reaction mechanism difference in the bond-cleavage. Spirocyclic product **7g** was also obtained in high yield from **2g** (entry 6). The reaction of monomethyl substrate **2h** with HClO, followed by acetylation, afforded keto-acetate **8h** in

67 % yield as a 1 : 0.17 diastereomer mixture (entry 7). It was difficult to isolate **7h** itself, because **7h** exists as a mixture of the acetal form and a ring-opened keto-alcohol tautomer in a solution.

Table 4. Oxidation of various cyclobutanols **2** with NaOCl·5H₂O/AcOH ^{a,b}

Entry	Substrates	Products	%yield of 7
1	2b (R ¹ = Cl, R ² = H)		7b 63
2	2c (R ¹ = H, R ² = Cl)		7c 74
3	2d (R ¹ = MeO, R ² = H)		7d 0
4	2e		7e 0
5	2f		7f 64
6 ^c	2g		7g 53
7 ^d	2h		8h 67 (dr = 1 : 0.17)


^aAll reactions were carried out on a 0.10 mmol scale (0.050 M). NaOCl·5H₂O (4.0 equiv.), AcOH (4.0 equiv.), CH₂Cl₂, 40 °C. ^bIsolated yield. ^cNaOCl·5H₂O (10 equiv.), AcOH (10 equiv.). ^dAfter the oxidation of **2h** with NaOCl·5H₂O/AcOH (8.0 equiv.), acetylation of the crude mixture of **7h** with Ac₂O (3.0 equiv.) and DMAP (10 mol%) was carried out at ambient temperature for 16 h.

Previously, it was reported that oxidative β-cleavage of cyclobutanols usually gives acyclic γ-functionalized ketones, which are formed by the addition of an external nucleophile to the ring-opening intermediate.⁹ In contrast, oxidation of **2** afforded **6** or **7** with a hydrofuran ring. We speculated that the


furan ring was formed by intramolecular addition of the oxygen atom derived from the hydroxy group of **2** prior to an intermolecular trap by an external reagent. This hypothesis was also supported by the following control experiments. In the presence of MeOH (20 equiv.) as an external nucleophile, neither compound **8a** nor **9a**, which could be formed by the electrophilic addition of MeOH, were detected in the oxidation of **2a** with PIDA or NaOCl·5H₂O/AcOH, respectively (Scheme 2).

Scheme 3 summarizes a plausible mechanism. First, the reaction of cyclobutanol **2** with PIDA or HClO, which is generated from NaOCl·5H₂O/AcOH, gives intermediate **10**. Then cleavage of the C(1)–C(2) bond selectively occurs to provide tertiary-cationic intermediate **11**. This cleavage is driven by the release of the cyclobutane ring strain. The formation of benzyl-cationic intermediate **12** by the cleavage of C(1)–C(4) bond is unfavorable due to the strain of the sp²-rich 8-membered ring. Keto-type intermediate **11** readily tautomerizes into phenol intermediate **13** followed by an intramolecular attack on the tertiary cation to give dihydrofuran **6**. The higher nucleophilicity of the phenolic hydroxy group of **13** than that of the carbonyl oxygen of **11** is the rationale for the formation of the furan ring. In the reaction using HClO, phenanthrofuran **6** was further chlorinated to give **7**.

Scheme 2. Oxidative reactions of **2a** in the presence of MeOH.



Scheme 3. Plausible reaction mechanism.



Conclusions

We demonstrate that oxidative β -cleavage of dibenzo-fused bicyclo[4.2.0]octanols by PIDA gives dihydrophenanthro[9,10-*b*]furans in an unusual regioselective manner. We also show that oxidation of the alcohols by NaClO/AcOH affords higher oxidized products. This method provides various furan-fused polycyclic aromatic compounds in good yields. These oxidative transformations should serve as a new way to prepare functionalized polycyclic aromatic compounds.

Experimental Section

General Information. For reactions that require heating, oil bath was used as a heat source. Column chromatography was performed on Kanto Kagaku Silica Gel 60 N (spherical, neutral) 100-210 μm . Reactions and chromatography fractions were analyzed by thin-layer chromatography (TLC) carried out on Wako Silicagel 70 F₂₅₄ TLC Plate-Wako with visualization by ultraviolet (UV) irradiation at 254 nm, phosphomolybdic acid, anisaldehyde, ninhydrin, and/or potassium permanganate staining. NMR was

recorded on a JEOL JNM-LA (500 MHz for ^1H and 126 MHz for ^{13}C { ^1H }) or a JEOL JNM ECZ600R (600 MHz for ^1H and 151 MHz for ^{13}C { ^1H }) and measured in CDCl_3 unless otherwise mentioned. Chemical shifts and coupling constants are presented in ppm δ relative to tetramethylsilane (0.0 ppm) or $\text{C}_6\text{D}_5\text{H}$ (7.16 ppm) and Hz, respectively. Chloroform- d_1 (δ 77.0 ppm) was used as an internal standard for ^{13}C { ^1H } NMR. Abbreviations are as follows: s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; br, broad. IR spectra were recorded on a Shimadzu IRAffinity-1 and the wave numbers of maximum absorption peaks of them are presented in cm^{-1} . High-resolution mass spectra (HRMS) were recorded on a JEOL MS700 spectrometer (FAB) or a SHIMADZU LCMS-IT-TOF fitted with an ESI. Melting points were determined on YANACO micro melting point apparatus. X-ray single crystal diffraction analyses were performed on a Rigaku XtaLAB P200 apparatus. All reagents were purchased from chemical companies and used as received. Dehydrated solvents were purchased for the reactions and used without further desiccation unless otherwise mentioned.

Preparation of Substrates 2. Synthesis of substrates **2** was carried out in accordance with the reported procedure.⁵ Compound **2h** was known.⁵

(2aR*,10bR*)-2,2-Dimethyl-1,10b-dihydrocyclobuta[l]phenanthren-2a(2H)-ol (2a). White solids, Mp. 122–124 °C; ^1H NMR (500 MHz, CDCl_3) δ 8.02 (dd, J = 8.0, 1.0 Hz, 1H), 7.95 (d, J = 8.0 Hz, 1H), 7.46 (dd, J = 7.4, 1.7 Hz, 1H), 7.40 (ddd, J = 7.4, 7.4, 1.7 Hz, 1H), 7.36 (ddd, J = 7.4, 7.4, 1.5 Hz, 1H), 7.29 (ddd, J = 7.6, 7.6, 1.5 Hz, 1H), 7.24 (ddd, J = 7.4, 7.4, 1.1 Hz, 1H), 7.15 (dd, J = 7.2, 1.1 Hz, 1H), 3.72 (dd, J = 9.7, 9.7 Hz, 1H), 2.14 (brs, 1H), 2.05 (dd, J = 10.3, 9.7 Hz, 1H), 1.47 (s, 3H), 1.19 (dd, J = 10.3, 9.7 Hz, 1H), 0.84 (s, 3H); ^{13}C { ^1H } NMR (126 MHz, CDCl_3) δ 136.5, 135.8, 132.3, 130.6, 130.1, 128.3, 128.2, 127.9, 127.6, 126.8, 123.4, 122.6, 74.1, 45.7, 43.8, 38.6, 26.8, 24.2; IR (neat) 3406, 2963, 752, cm^{-1} ; HRMS (ESI) m/z : $[\text{M} + \text{Na}]^+$ Calcd for $\text{C}_{18}\text{H}_{18}\text{ONa}$ 273.1250; Found 273.1247.

(2aR*,10bR*)-4-Chloro-2,2-dimethyl-1,10b-dihydrocyclobuta[l]phenanthren-2a(2H)-ol (2b). Colorless oil, ^1H NMR (500 MHz, CDCl_3) δ 7.90 (d, J = 8.6 Hz, 1H), 7.84 (d, J = 7.2 Hz, 1H), 7.41 (d, J = 2.3 Hz, 1H), 7.32 (dd, J = 8.6, 2.3 Hz, 1H), 7.27 (td, J = 7.2, 1.4 Hz, 1H), 7.23 (td, J = 7.2, 1.4 Hz, 1H),

7.11 (dd, $J = 7.2, 1.1$ Hz, 1H), 3.65 (dd, $J = 9.6, 9.5$ Hz, 1H), 2.25 (brs, 1H), 2.05 (dd, $J = 10.7, 9.6$ Hz, 1H), 1.42 (s, 3H), 1.17 (dd, $J = 10.7, 9.6$ Hz, 1H), 0.83 (s, 3H); ^{13}C $\{^1\text{H}\}$ NMR (126 MHz, CDCl_3) δ 137.5, 136.3, 133.1, 130.9, 129.8, 129.7, 128.33, 128.28, 128.2, 126.9, 124.1, 123.3, 74.0, 45.6, 43.7, 38.7, 26.6, 24.1; IR (neat) 3390, 2967, 1447, 764 cm^{-1} ; HRMS (FAB) m/z : $[\text{M} + \text{Na}]^+$ Calcd for $\text{C}_{18}\text{H}_{17}\text{ClONa}$ 307.0866; Found 307.0871.

(2aR*,10bR*)-9-Chloro-2,2-dimethyl-1,10b-dihydrocyclobuta[l]phenanthren-2a(2H)-ol (2c).

Colorless oil, ^1H NMR (500 MHz, CDCl_3) δ 7.95 (dd, $J = 7.5, 2.0$ Hz, 1H), 7.85 (d, $J = 8.6$ Hz, 1H), 7.47–7.45 (m, 1H), 7.40 (td, $J = 7.2, 2.0$ Hz, 1H), 7.37 (td, $J = 7.2, 1.7$ Hz, 1H), 7.24 (dd, $J = 8.6, 2.3$ Hz, 1H), 7.14 (d, $J = 2.3$ Hz, 1H), 3.67 (dd, $J = 10.0, 9.5$ Hz, 1H), 2.14 (brs, 1H), 2.06 (dd, $J = 10.0, 10.0$ Hz, 1H), 1.46 (s, 3H), 1.18 (dd, $J = 10.0, 9.5$ Hz, 1H), 0.83 (s, 3H); ^{13}C $\{^1\text{H}\}$ NMR (126 MHz, CDCl_3) δ 138.3, 135.5, 133.3, 131.2, 130.1, 129.2, 128.2, 127.9, 127.8, 126.8, 124.8, 122.4, 74.0, 45.8, 43.4, 38.4, 26.6, 24.1; IR (neat) 3395, 2963, 1443, 764 cm^{-1} ; HRMS (FAB) m/z : $[\text{M} + \text{Na}]^+$ Calcd for $\text{C}_{18}\text{H}_{17}\text{ClONa}$ 307.0866; Found 307.0862.

(2aR*,10bR*)-4-Methoxy-2,2-dimethyl-1,10b-dihydrocyclobuta[l]phenanthren-2a(2H)-ol (2d). White solids, Mp. 93–95 °C; ^1H NMR (500 MHz, CDCl_3) δ 7.88 (d, $J = 9.7$ Hz, 1H), 7.80 (d, $J = 7.5$ Hz, 1H), 7.22 (td, $J = 7.1, 1.4$ Hz, 1H), 7.15 (td, $J = 7.3, 1.1$ Hz, 1H), 7.07 (dd, $J = 7.5, 1.1$ Hz, 1H), 6.92–6.89 (m, 2H), 3.80 (s, 3H), 3.62 (dd, $J = 9.7, 9.7$ Hz, 1H), 2.31 (brs, 1H), 2.01 (dd, $J = 10.7, 9.7$ Hz, 1H), 1.41 (s, 3H), 1.16 (t, $J = 10.7, 9.7$ Hz, 1H), 0.83 (s, 3H); ^{13}C $\{^1\text{H}\}$ NMR (126 MHz, CDCl_3) δ 158.9, 137.4, 135.6, 130.6, 128.1, 126.9, 126.7, 125.3, 124.0, 122.7, 114.9, 113.8, 74.3, 55.3, 45.5, 43.7, 38.5, 26.9, 24.2; IR (neat) 3410, 2959, 1485, 768 cm^{-1} ; HRMS (ESI) m/z : $[\text{M} + \text{Na}]^+$ Calcd for $\text{C}_{19}\text{H}_{20}\text{O}_2\text{Na}$ 303.1356; Found 303.1356.

(7bR*,9aR*)-9,9-Dimethyl-8,9-dihydrocyclobuta[3,4]naphtho[2,1-b]thiophen-9a(7bH)-ol (2e).

Colorless solids, Mp. 75–77 °C; ^1H NMR (500 MHz, CDCl_3) δ 7.65 (d, $J = 7.7$ Hz, 1H), 7.47 (d, $J = 5.3$ Hz, 1H), 7.42 (d, $J = 5.3$ Hz, 1H), 7.26 (td, $J = 7.5, 1.4$ Hz, 1H), 7.19 (td, $J = 7.5, 1.4$ Hz, 1H), 7.15 (dd, $J = 7.5, 1.7$ Hz, 1H), 3.79 (dd, $J = 10.0, 10.0$ Hz, 1H), 2.28 (brs, 1H), 2.03 (dd, $J = 10.6, 10.0$ Hz, 1H), 1.46 (s, 3H), 1.27 (dd, $J = 10.6, 10.0$ Hz, 1H), 1.07 (s, 3H); ^{13}C $\{^1\text{H}\}$ NMR (126 MHz, CDCl_3) δ 138.2, 136.3, 135.3,

129.3, 127.6, 127.2, 126.81, 126.78, 123.5, 122.2, 74.1, 46.2, 45.5, 38.1, 26.6, 24.0; IR (neat) 3426, 2963, 748, 718 cm^{-1} ; HRMS (ESI) m/z : $[\text{M} - \text{H}_2\text{O} + \text{H}]^+$ Calcd for $\text{C}_{16}\text{H}_{15}\text{S}$ 239.0889; Found 239.0890.

(2aR*,12bR*)-2,2-Dimethyl-1,12b-dihydrobenzo[c]cyclobuta[a]phenanthren-2a(2H)-ol (2f). White solids, Mp. 114–116 °C; ^1H NMR (500 MHz, CDCl_3) δ 8.64–8.62 (m, 1H), 7.97 (d, $J = 7.7$ Hz, 1H), 7.85–7.83 (m, 1H), 7.72 (d, $J = 8.6$ Hz, 1H), 7.51–7.47 (m, 3H), 7.25–7.16 (m, 3H), 3.71 (dd, $J = 10.2$, 10.0 Hz, 1H), 2.44 (brs, 1H), 1.88 (dd, $J = 10.5$, 10.0 Hz, 1H), 1.46 (s, 3H), 1.23 (dd, $J = 10.5$, 10.2 Hz, 1H), 0.56 (s, 3H); ^{13}C $\{^1\text{H}\}$ NMR (126 MHz, CDCl_3) δ 136.8, 134.1, 133.5, 131.8, 131.1, 129.8, 129.7, 128.8, 128.2, 127.4, 127.3, 127.11, 127.07, 126.3, 125.6, 125.4, 74.9, 46.5, 45.6, 36.3, 26.0, 24.2; IR (neat) 3387, 2967, 745 cm^{-1} ; HRMS (ESI) m/z : $[\text{M} + \text{Na}]^+$ Calcd for $\text{C}_{22}\text{H}_{20}\text{ONa}$ 323.1406; Found 323.1408.

(2aR*,10bR*)-2,2a-Dihydro-10bH-spiro[cyclobuta[l]phenanthrene-1,1'-cyclohexan]-10b-ol (2g). White solids, Mp. 120–122 °C; ^1H NMR (500 MHz, CDCl_3) δ 8.01 (dd, $J = 7.7$, 1.4, 1H), 7.93 (d, $J = 7.7$ Hz, 1H), 7.47 (dd, $J = 7.7$, 1.4 Hz, 1H), 7.38 (td, $J = 7.5$, 1.7 Hz, 1H), 7.34 (td, $J = 7.5$, 1.4 Hz, 1H), 7.29 (td, $J = 7.7$, 1.4 Hz, 1H), 7.24 (td, $J = 7.2$, 1.4 Hz, 1H), 7.14 (dd, $J = 7.5$, 1.7 Hz, 1H), 3.66 (dd, $J = 10.3$, 9.5 Hz, 1H), 2.31 (dd, $J = 10.6$, 10.3 Hz, 1H), 2.15–2.14 (m, 2H), 1.76–1.71 (m, 2H), 1.59 (m, 1H), 1.52–1.42 (m, 2H), 1.29 (td, $J = 12.9$, 3.4, 1H), 1.14 (qdd, $J = 12.7$, 3.4, 3.4 Hz, 1H), 1.14 (dd, $J = 10.6$, 9.5 Hz, 1H), 1.02 (qdd, $J = 12.9$, 2.9, 2.9 Hz, 1H), 0.93 (m, 1H); ^{13}C $\{^1\text{H}\}$ NMR (126 MHz, CDCl_3) δ 136.8, 135.4, 132.3, 130.6, 130.1, 128.3, 128.1, 127.9, 127.4, 126.8, 123.4, 122.5, 74.9, 49.9, 43.3, 36.3, 35.5, 31.6, 26.1, 22.7, 22.2; IR (neat) 3368, 2924, 1447, 748 cm^{-1} ; HRMS (ESI) m/z : $[\text{M} + \text{Na}]^+$ Calcd for $\text{C}_{21}\text{H}_{22}\text{ONa}$ 313.1563; Found 313.1565.

General Procedure for Oxidative β -Cleavage of 2a–h with PIDA. To a solution of cyclobutanol **2a–h** (0.10 mmol) in $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ (9:1, 1.0 mL) was added PIDA (0.11 mmol) at ambient temperature under Ar atmosphere, and the resulting solution was stirred for 1 h. After the completion of the reaction, the solvent was removed under reduced pressure. The residue was purified by column chromatography (hexanes/EtOAc) to give dihydrofuran **6a–h**.

2,2-Dimethyl-2,3-dihydrophenanthro[9,10-b]furan (6a). The reaction was conducted according to the

general procedure with cyclobutanol **2a** (25.0 mg, 0.10 mmol), and PIDA (64 mg, 0.20 mmol). Purification by silica gel column chromatography (hexanes/EtOAc: 10:1) afforded compound **6a** (24.5 mg, 97% yield) as white solids. Mp. 121–122 °C; ¹H NMR (500 MHz, CDCl₃) δ 8.69 (d, *J* = 7.7 Hz, 1H), 8.65 (d, *J* = 8.3 Hz, 1H), 8.09 (dd, *J* = 7.5, 1.4 Hz, 1H), 7.67–7.57 (m, 4H), 7.51–7.48 (m, 1H), 3.37 (s, 2H), 1.65 (s, 6H); ¹³C {¹H} NMR (126 MHz, CDCl₃) δ 152.1, 131.2, 130.6, 126.9, 126.6, 126.34, 126.29, 123.23, 123.16, 122.98, 122.96, 122.5, 122.2, 113.3, 87.6, 42.6, 28.8; IR (neat) 2970, 1359, 752 cm⁻¹; HRMS (ESI) *m/z*: [M + H]⁺ Calcd for C₁₈H₁₇O 249.1274; Found 249.1264.

Oxidation of 2a with PIDA in a preparative scale: To a solution of cyclobutanol **2a** (501 mg, 2.0 mmol) in CH₃CN/H₂O (9:1, 20 mL) was added PIDA (709 mg, 2.2 mmol) at ambient temperature under Ar atmosphere, and the resulting solution was stirred for 1 h. After the completion of the reaction, the solvent was removed under reduced pressure. The residue was purified by column chromatography (hexanes/EtOAc: 10 : 1) to afford dihydrofuran **6a** (426 mg, 86% yield) as white solids.

10-Chloro-2,2-dimethyl-2,3-dihydrophenanthro[9,10-b]furan (6b). The reaction was conducted according to the general procedure with cyclobutanol **2b** (28.4 mg, 0.10 mmol), and PIDA (35 mg, 0.11 mmol). Purification by silica gel column chromatography (hexanes/EtOAc: 10:1) afforded compound **6b** (17.5 mg, 62% yield) as white solids. Mp. 88–90 °C; ¹H NMR (500 MHz, CDCl₃) δ 8.58 (d, *J* = 8.9 Hz, 1H), 8.57 (d, *J* = 8.0 Hz, 1H), 8.06 (d, *J* = 2.3 Hz, 1H), 7.61–7.59 (m, 2H), 7.57 (dd, *J* = 8.9, 2.3 Hz, 1H), 7.49 (ddd, *J* = 8.5, 5.6, 2.9, 1H), 3.36 (s, 2H), 1.64 (s, 6H); ¹³C {¹H} NMR (126 MHz, CDCl₃) δ 151.1, 132.4, 130.5, 129.5, 127.2, 126.8, 126.2, 124.7, 123.6, 123.4, 123.2, 123.1, 121.6, 114.7, 88.0, 42.6, 28.8; IR (neat) 2970, 752 cm⁻¹; HRMS (FAB) *m/z*: [M + H]⁺ Calcd for C₁₈H₁₆ClO 283.0890; Found 283.0892.

5-Chloro-2,2-dimethyl-2,3-dihydrophenanthro[9,10-b]furan (6c). The reaction was conducted according to the general procedure with cyclobutanol **2c** (28.4 mg, 0.10 mmol), and PIDA (35 mg, 0.11 mmol). Purification by silica gel column chromatography (hexanes/EtOAc: 10:1) afforded compound **6c** (22.8 mg, 81% yield) as white solids. Mp. 99–101 °C; ¹H NMR (600 MHz, CDCl₃) δ 8.59 (d, *J* = 7.6 Hz, 1H), 8.53 (d, *J* = 9.0 Hz, 1H), 8.08 (d, *J* = 7.6 Hz, 1H), 7.65 (ddd, *J* = 8.3, 6.9, 1.4 Hz, 1H), 7.62 (dd, *J* = 7.6, 6.9 Hz, 1H), 7.56 (d, *J* = 2.1 Hz, 1H), 7.41 (dd, *J* = 9.0, 2.1 Hz, 1H), 3.32 (s, 2H), 1.64 (s, 6H); ¹³C

$\{^1\text{H}\}$ NMR (151 MHz, CDCl_3) δ 153.1, 132.9, 131.8, 130.8, 126.8, 126.5, 124.9, 124.8, 123.4, 122.9, 122.4, 122.3, 122.1, 112.5, 88.0, 42.4, 28.7; IR (neat) 764 cm^{-1} ; HRMS (ESI) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{18}\text{H}_{16}\text{ClO}$ 283.0884; Found 283.0882.

10-Methoxy-2,2-dimethyl-2,3-dihydrophenanthro[9,10-*b*]furan (6d). The reaction was conducted according to the general procedure with cyclobutanol **2d** (28.0 mg, 0.10 mmol), and PIDA (35 mg, 0.11 mmol). Purification by silica gel column chromatography (hexanes/EtOAc: 10:1) afforded compound **6d** (25.3 mg, 91% yield) as white solids. Mp. 121–122 °C; ^1H NMR (500 MHz, CDCl_3) δ 8.57 (d, $J = 9.2$ Hz, 1H), 8.55 (d, $J = 8.3$ Hz, 1H), 7.59 (dd, $J = 7.9, 1.0$ Hz, 1H), 7.52 (ddd, $J = 8.0, 6.9, 1.2$ Hz, 1H), 7.47 (ddd, $J = 8.3, 6.9, 1.4$ Hz, 1H), 7.42 (d, $J = 2.9$ Hz, 1H), 7.26 (dd, $J = 9.2, 2.9$ Hz, 1H), 4.00 (s, 3H), 3.37 (s, 2H), 1.65 (s, 6H); ^{13}C $\{^1\text{H}\}$ NMR (126 MHz, CDCl_3) δ 158.3, 151.7, 129.4, 126.8, 125.9, 125.5, 124.7, 123.6, 123.3, 122.9, 122.7, 117.2, 113.8, 102.0, 87.5, 55.5, 42.7, 28.8; IR (neat) $2966, 752\text{ cm}^{-1}$; HRMS (FAB) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{19}\text{H}_{19}\text{O}_2$ 279.1385; Found 279.1380.

2,2-Dimethyl-2,3-dihydrothieno[2',3':3,4]naphtho[2,1-*b*]furan (6e). The reaction was conducted according to the general procedure with cyclobutanol **2e** (25.6 mg, 0.10 mmol), and PIDA (35 mg, 0.11 mmol). Purification by silica gel column chromatography (hexanes/EtOAc: 10:1) afforded compound **6e** (25.2 mg, 99% yield) as white solids. Mp. 127–128 °C; ^1H NMR (500 MHz, CDCl_3) δ 8.32 (d, $J = 8.3$ Hz, 1H), 8.00 (d, $J = 5.2$ Hz, 1H), 7.65 (d, $J = 8.0$ Hz, 1H), 7.55 (d, $J = 5.4$ Hz, 1H), 7.52 (dd, $J = 6.9, 1.2$ Hz, 1H), 7.45 (ddd, $J = 8.3, 6.9, 1.2$ Hz, 1H), 3.39 (s, 2H), 1.66 (s, 6H); ^{13}C $\{^1\text{H}\}$ NMR (126 MHz, CDCl_3) δ 150.6, 138.1, 129.6, 125.8, 125.4, 125.2, 124.3, 123.6, 123.2, 122.9, 122.5, 113.5, 89.2, 42.3, 28.7; IR (neat) $2970, 756\text{ cm}^{-1}$; HRMS (ESI) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{16}\text{H}_{15}\text{OS}$ 255.0838; Found 255.0839.

2,2-Dimethyl-1,2-dihydrobenzo[3,4]phenanthro[1,2-*b*]furan (6f). The reaction was conducted according to the general procedure with cyclobutanol **2f** (30.0 mg, 0.10 mmol), and PIDA (35 mg, 0.11 mmol). Purification by silica gel column chromatography (hexanes/EtOAc: 10:1) afforded compound **6f** (17.6 mg, 59% yield) as a yellowish oil. ^1H NMR (500 MHz, CDCl_3) δ 9.10 (dd, $J = 8.6, 8.3$ Hz, 2H), 8.09 (d, $J = 8.6$ Hz, 1H), 8.02 (d, $J = 8.0$ Hz, 1H), 7.92 (d, $J = 8.6$ Hz, 1H), 7.72 (d, $J = 7.7$ Hz, 1H), 7.67 (t, $J = 7.7$ Hz, 1H), 7.63–7.59 (m, 2H), 7.53–7.50 (m, 1H), 3.44 (s, 2H), 1.67 (s, 6H); ^{13}C $\{^1\text{H}\}$ NMR (126 MHz,

CDCl₃) δ 152.7, 137.5, 133.4, 131.4, 130.4, 128.8, 128.7, 128.3, 128.1, 127.3, 126.5, 126.3, 126.0, 125.7, 123.0, 122.7, 120.6, 120.2, 114.9, 87.9, 42.6, 28.8; IR (neat) 2970, 760 cm⁻¹; HRMS (ESI) m/z : [M + H]⁺ Calcd for C₂₂H₁₉O 299.1430; Found 299.1429.

3'-H-Spiro[cyclohexane-1,2'-phenanthro[9,10-b]furan] (6g). The reaction was conducted according to the general procedure with cyclobutanol **2g** (29.0 mg, 0.10 mmol), and PIDA (35 mg, 0.11 mmol). Purification by silica gel column chromatography (hexanes/EtOAc: 10:1) afforded compound **6g** (28.0 mg, 97% yield) as colorless solids. Mp. 128–129 °C; ¹H NMR (500 MHz, CDCl₃) δ 8.69 (dd, J = 7.3, 2.1 Hz, 1H), 8.66 (d, J = 8.3 Hz, 1H), 8.16–8.13 (m, 1H), 7.66 (td, J = 6.9, 1.7 Hz, 1H), 7.65–7.62 (m, 2H), 7.60 (td, J = 6.9, 1.2 Hz, 1H), 7.50 (ddd, J = 9.2, 8.3, 1.6 Hz, 1H), 3.31 (s, 2H), 2.05–2.01 (m, 2H), 1.98–1.92 (m, 2H), 1.84–1.79 (m, 2H), 1.65–1.53 (m, 4H); ¹³C {¹H} NMR (126 MHz, CDCl₃) δ 152.1, 131.2, 130.7, 126.9, 126.5, 126.3, 126.2, 123.2, 123.0, 122.93, 122.86, 122.6, 122.2, 113.0, 89.5, 40.9, 37.6, 25.3, 23.1; IR (neat) 2931, 752 cm⁻¹; HRMS (FAB) m/z : [M + H]⁺ Calcd for C₂₁H₂₁O 289.1592; Found 289.1598.

General Procedure for Oxidative β -Cleavage of 2a–g with NaOCl·5H₂O/AcOH. To a solution of cyclobutanol **2a–g** (0.10 mmol) and NaOCl·5H₂O (0.40 mmol) in CH₂Cl₂ (2.0 mL) was added AcOH (0.40 mmol) at 40 °C under Ar atmosphere and the resulting solution was stirred for 1 h. After the completion of the reaction, the reaction mixture was quenched with sat. Na₂S₂O₃ aq. and extracted with EtOAc three times. The combined organic layers were washed with sat. NaHCO₃ aq. and brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The residue was purified with silica gel column chromatography (hexanes/EtOAc) to give hemiacetal **7a–g**.

(3aS*,11bS*)-3a-chloro-2,2-dimethyl-3,3a-dihydrophenanthro[9,10-b]furan-11b(2H)-ol (7a). The reaction was conducted according to the general procedure with cyclobutanol **2a** (25.0 mg, 0.10 mmol), NaOCl·5H₂O (132 mg, 0.40 mmol) and AcOH (46 μ L, 0.40 mmol). Purification by silica gel column chromatography (hexanes/EtOAc: 10:1 to 3:1) afforded compound **7a** (24.4 mg, 81% yield) as white solids. Mp. 113–115 °C; ¹H NMR (500 MHz, CDCl₃) δ 8.00 (dd, 1H, J = 7.7, 1.7 Hz), 7.92 (dd, 1H, J = 7.2, 2.0 Hz), 7.88 (dd, 1H, J = 7.7, 0.9 Hz), 7.74 (dd, 1H, J = 7.4, 1.7 Hz), 7.48 (td, 1H, J = 7.7, 1.4 Hz), 7.38–7.46 (m, 3H), 3.42 (brs, 1H), 2.99 (s, 2H), 1.57 (s, 3H), 0.76 (s, 3H); ¹H NMR (500 MHz, C₆D₆) δ 8.06–8.08 (m,

1H), 7.54–7.57 (m, 1H), 7.50–7.53 (m, 2H), 7.11–7.14 (m, 2H), 7.02–7.08 (m, 2H), 3.20 (s, 1H), 2.82 (d, $J = 12.9$ Hz, 1H), 2.64 (d, $J = 12.9$ Hz, 1H), 1.47 (s, 3H), 0.65 (s, 3H); ^{13}C $\{^1\text{H}\}$ NMR (126 MHz, CDCl_3) δ 135.7, 133.7, 131.3, 130.7, 129.8, 129.1, 128.6, 128.5, 128.3, 127.0, 124.0, 123.8, 101.9, 80.7, 74.5, 51.3, 31.2, 28.2; IR (neat) 3502, 2974, 756, 732 cm^{-1} ; HRMS (ESI) m/z : $[\text{M} + \text{Na}]^+$ Calcd for $\text{C}_{18}\text{H}_{17}\text{ClO}_2\text{Na}$ 323.0809; Found 323.0808. X-ray crystallographic analysis (CCDC 2103959): Recrystallization from pentane/MeOH gave colorless platelets suitable for X-ray crystal structural analysis: orthorhombic $P2_12_12_1$; $a = 10.3098(3)$, $b = 11.4153(3)$, $c = 12.3089.205(3)$; $V = 1448.63(6)$, $Z = 4$, $D_x = 1.379$.

Oxidation of 2a with NaOCl·5H₂O/AcOH in a preparative scale: To a solution of cyclobutanol **2a** (501 mg, 2.0 mmol) and NaOCl·5H₂O (2.64 g, 16 mmol) in CH_2Cl_2 (40 mL) was added AcOH (1.0 mL, 16 mmol) at 40 °C under Ar atmosphere and the resulting solution was stirred for 1 h. After the completion of the reaction, the reaction mixture was quenched with sat. $\text{Na}_2\text{S}_2\text{O}_3$ aq. and extracted with EtOAc three times. The combined organic layers were washed with sat. NaHCO_3 aq. and brine, dried over Na_2SO_4 , filtered, and concentrated *in vacuo*. The residue was purified with silica gel column chromatography (hexanes/EtOAc: 10 : 1 to 3 : 1) to afford hemiacetal **7a** (434 mg, 72% yield) as white solids.

(3aS*,11bS*)-3a,10-Dichloro-2,2-dimethyl-3,3a-dihydrophenanthro[9,10-b]furan-11b(2H)-ol (7b). The reaction was conducted according to the general procedure with cyclobutanol **2b** (28.4 mg, 0.10 mmol), NaOCl·5H₂O (132 mg, 0.40 mmol) and AcOH (46 μL , 0.40 mmol). Purification by silica gel column chromatography (hexanes/EtOAc: 10:1 to 3:1) afforded compound **7b** (21.1 mg, 63% yield) as white solids. Mp. 114–116 °C; ^1H NMR (600 MHz, CDCl_3) δ 8.01 (d, $J = 2.1$ Hz, 1H), 7.88–7.87 (m, 1H), 7.91 (d, $J = 9.0$ Hz, 1H), 7.73–7.72 (m, 1H), 7.46–7.42 (m, 3H), 3.43 (s, –OH), 3.03 (d, $J = 13.1$ Hz, 1H), 2.99 (d, $J = 13.1$ Hz, 1H), 1.56 (s, 3H), 0.78 (s, 3H); ^{13}C $\{^1\text{H}\}$ NMR (126 MHz, CDCl_3) δ 135.8, 135.5, 134.6, 129.90, 129.86, 129.81, 129.3, 128.8, 128.4, 127.4, 125.3, 123.9, 101.4, 81.2, 74.5, 51.2, 31.2, 28.1; IR (neat): 3541, 2974, 1477, 764, 729 cm^{-1} ; HRMS (ESI) m/z : $[\text{M} + \text{Na}]^+$ Calcd for $\text{C}_{18}\text{H}_{16}\text{Cl}_2\text{O}_2\text{Na}$ 357.0420; Found 357.0438.

(3aS*,11bS*)-3a,5-Dichloro-2,2-dimethyl-3,3a-dihydrophenanthro[9,10-b]furan-11b(2H)-ol (7c). The reaction was conducted according to the general procedure with cyclobutanol **2c** (28.4 mg, 0.10 mmol),

NaOCl·5H₂O (132 mg, 0.40 mmol) and AcOH (46 μL, 0.40 mmol). Purification by silica gel column chromatography (hexanes/EtOAc: 10:1 to 3:1) afforded compound **7c** (24.8 mg, 74% yield) as white solids. Mp. 109–111 °C; ¹H NMR (500 MHz, CDCl₃): δ 7.99 (dd, *J* = 7.4, 1.7 Hz, 1H), 7.86 (d, *J* = 8.6 Hz, 1H), 7.82 (dd, *J* = 7.7, 1.1 Hz, 1H), 7.72 (d, *J* = 2.3 Hz, 1H), 7.49 (td, *J* = 7.4, 1.7 Hz, 1H), 7.45 (td, *J* = 7.4, 1.3 Hz, 1H), 7.40 (dd, *J* = 8.6, 2.3 Hz, 1H), 3.42 (s, 1H), 2.97 (d, *J* = 13.2 Hz, 1H), 2.89 (d, *J* = 13.2 Hz, 1H), 1.57 (s, 3H), 0.84 (s, 3H); ¹³C {¹H} NMR (126 MHz, CDCl₃): δ 137.8, 134.1, 133.6, 129.9, 129.4, 129.3, 129.0, 128.4, 127.2, 125.5, 123.7, 101.8, 80.6, 74.6, 51.5, 31.1, 28.4; IR (neat): 3510, 2974, 764 cm⁻¹; HRMS (ESI) *m/z*: [M + Na]⁺ Calcd for C₁₈H₁₆Cl₂O₂Na 357.0420; Found 357.0422.

(3a*S*^{*},11b*S*^{*})-13b-Chloro-2,2-dimethyl-1,13b-dihydrobenzo[3,4]phenanthro[1,2-*b*]furan-3a(2*H*)-ol (7f). The reaction was conducted according to the general procedure with cyclobutanol **2f** (30.0 mg, 0.10 mmol), NaOCl·5H₂O (132 mg, 0.40 mmol) and AcOH (46 μL, 0.40 mmol). Purification by silica gel column chromatography (hexanes/EtOAc: 10:1 to 3:1) afforded compound **7f** (22.5 mg, 64% yield) as white solids. Mp. 114–116 °C; ¹H NMR (500 MHz, CDCl₃) δ 8.55 (d, *J* = 8.3 Hz, 1H), 8.39 (d, *J* = 8.6 Hz, 1H), 8.17 (s, 1H), 7.98–7.95 (m, 1H), 7.89–7.85 (m, 1H), 7.68–7.64 (m, 1H), 7.61–7.57 (m, 1H), 7.48 (dd, *J* = 9.5, 1.7 Hz, 1H), 7.46 (t, *J* = 6.4 Hz, 1H), 7.44 (dd, *J* = 9.5, 1.7 Hz, 1H), 3.49 (brs, 1H), 2.94 (s, 2H), 1.59 (s, 3H), 0.68 (s, 3H); ¹³C {¹H} NMR (126 MHz, CDCl₃) δ 136.0, 132.6, 132.2, 132.0, 130.5, 130.2, 129.8, 128.6, 128.2, 128.1, 128.0, 127.3, 127.0, 126.8, 125.0, 124.5, 102.3, 80.8, 75.2, 51.2, 31.1, 28.3; IR (neat) 1365, 756 cm⁻¹; HRMS (ESI) *m/z*: [M + K]⁺ Calcd for C₂₂H₁₉ClO₂K 389.0705; Found 389.0713.

(3a'*S*^{*},11b'*S*^{*})-3a'-Chloro-3',3a'-dihydro-11b'*H*-spiro[cyclohexane-1,2'-phenanthro[9,10-*b*]furan-11b'-ol (7g). The reaction was conducted according to the general procedure with cyclobutanol **2g** (29.0 mg, mmol), NaOCl·5H₂O (132 mg, 0.40 mmol) and AcOH (46 μL, 0.40 mmol). Purification by silica gel column chromatography (hexanes/EtOAc: 10:1 to 3:1) afforded compound **7g** (18.1 mg, 53% yield) as a colorless oil. ¹H NMR (600 MHz, CDCl₃) δ 8.01 (d, *J* = 7.6 Hz, 1H), 7.91 (d, *J* = 7.6 Hz, 1H), 7.87 (d, *J* = 8.3 Hz, 1H), 7.75 (dd, *J* = 7.6, 1.4 Hz, 1H), 7.48 (td, *J* = 7.6, 1.4 Hz, 1H), 7.43 (td, *J* = 8.3, 1.4 Hz, 1H), 7.41–7.38 (m, 2H), 3.48 (brs, 1H), 2.93 (d, *J* = 13.5 Hz, 1H), 2.85 (d, *J* = 13.5 Hz, 1H), 1.99–1.93 (m, 1H), 1.90–1.89 (m, 1H), 1.78–1.75 (m, 1H), 1.51–1.48 (m, 1H), 1.40–1.36 (m, 2H), 1.27–1.24 (m, 1H),

1.18–1.15 (m, 1H), 1.07–1.03 (m, 1H), 0.91–0.88 (m, 1H); ^{13}C $\{^1\text{H}\}$ NMR (151 MHz, CDCl_3) δ 136.1, 134.1, 131.2, 130.5, 129.7, 129.1, 128.6, 128.4, 128.2, 127.1, 123.9, 123.6, 101.3, 82.6, 75.7, 49.5, 40.3, 36.7, 25.1, 23.6, 23.5; IR (neat); 3507, 2932, 756, 729 cm^{-1} ; HRMS (ESI) m/z : $[\text{M} + \text{Na}]^+$ Calcd for $\text{C}_{21}\text{H}_{21}\text{ClO}_2\text{Na}$ 363.1122; Found 363.1126.

(*S*^{*})-1-[(*S*^{*} and *R*^{*})-9-chloro-10-oxo-9,10-dihydrophenanthren-9-yl]propan-2-yl acetate (8h). To a solution of cyclobutanol **2h** (49.0 mg, 0.21 mmol) and $\text{NaOCl}\cdot 5\text{H}_2\text{O}$ (273 mg, 1.66 mmol) in CH_2Cl_2 (2.0 mL) was added AcOH (95 μL , 1.66 mmol) at 40 $^\circ\text{C}$ under Ar atmosphere and the resulting solution was stirred for 30 min. After the completion of the reaction, the reaction mixture was quenched with sat. $\text{Na}_2\text{S}_2\text{O}_3$ aq. and extracted with EtOAc three times. The combined organic layers were washed with sat. NaHCO_3 aq. and brine, dried over Na_2SO_4 , filtered, and concentrated *in vacuo*. And then, the crude was dissolved in CH_2Cl_2 (2.0 mL) and acetic anhydride (59 μL , 0.62 mmol) and 4-dimethylaminopyridine (2.6 mg, 0.021 mmol) successively added to the mixture at ambient temperature and the resulting solution was stirred for 16 h. After the completion of the reaction, the reaction mixture was quenched with H_2O and brine, then extracted with EtOAc three times. The combined organic layers were washed with sat. NaHCO_3 aq. and brine, dried over Na_2SO_4 , filtered, and concentrated *in vacuo*. The residue was purified by silica gel column chromatography (hexane/AcOEt = 10:1 to 3:1) to afford product **8h** (45.6 mg, 67% over 2 steps) as colorless solids in a 1 : 0.17 diastereomeric ratio. ^1H NMR (500 MHz, CDCl_3) δ 8.28 (dd, J = 8.0, 1.4 Hz, 0.17H), 8.13 (d, J = 7.7, 1H), 8.06 (d, J = 8.6 Hz, 0.17H), 8.04–8.02 (m, 0.17H), 8.00 (d, J = 8.0 Hz, 1H), 7.98–7.96 (m, 1H), 7.87 (ddd, J = 8.6, 8.0, 7.6 Hz, 1H), 7.75–7.69 (m, 1.34H), 7.49–7.43 (m, 3.51H), 4.95 (dq, J = 8.9, 6.3, 3.2 Hz, 1H), 4.47 (dq, J = 11.2, 5.7, 2.9 Hz, 0.17H), 3.32 (dd, J = 14.0, 11.2 Hz, 0.17H), 2.85 (dd, J = 14.8, 8.9 Hz, 1H), 2.78 (dd, J = 14.8, 3.2 Hz, 1H), 2.69 (dd, J = 14.0, 2.9 Hz, 0.17H), 1.61 (s, 3H), 1.17 (s, 0.51H), 1.13 (d, J = 5.7 Hz, 0.51H), 1.12 (d, J = 6.3 Hz, 3H); ^{13}C $\{^1\text{H}\}$ NMR (126 MHz, CDCl_3 , signals from the minor diastereomer are marked with an asterisk) δ 192.4, 191.9*, 169.9, 169.5*, 137.6, 137.0*, 136.1, 136.0*, 135.2*, 135.1, 129.7*, 129.5, 129.2, 129.14, 129.10, 129.0*, 128.9*, 128.8, 128.7, 128.3, 127.7*, 123.9, 123.7*, 123.3*, 123.2, 69.5, 67.9*, 67.5, 66.1*, 48.9*, 47.3, 20.9, 20.5,

20.2*, 19.9* (three peaks are unidentified.); IR (neat) 1736, 1690, 1450, 1238, 729 cm⁻¹; HRMS (ESI) *m/z*: [M + Na]⁺ Calcd for C₁₉H₁₇ClO₃Na 351.0758; Found 351.0769.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: xxxxxx.

¹H, ¹³C {¹H} NMR spectra of new compounds and X-ray crystallography (PDF). Crystallographic data for **7a** (CCDC 2103959) (CIF).

AUTHOR INFORMATION

Corresponding Author

Kiyosei Takasu — Graduate School of Pharmaceutical Sciences, Kyoto University, Yoshida, Sakyo-ku, Kyoto 606-8501, Japan;

orcid.org/0000-0002-1798-7919; Phone: (+81)-75-753-4553; E-mail: kay-t@pharm.kyoto-u.ac.jp; Fax: (+81)-75-753-4604

Authors

Hayate Kinouchi — Graduate School of Pharmaceutical Sciences, Kyoto University, Yoshida, Sakyo-ku, Kyoto 606-8501, Japan

Kazuma Sugimoto — Graduate School of Pharmaceutical Sciences, Kyoto University, Yoshida, Sakyo-ku, Kyoto 606-8501, Japan

Yousuke Yamaoka — Graduate School of Pharmaceutical Sciences, Kyoto University, Yoshida, Sakyo-ku, Kyoto 606-8501, Japan;
orcid.org/0000-0003-3641-8823

Hiroshi Takikawa — Graduate School of Pharmaceutical Sciences, Kyoto University, Yoshida, Sakyo-ku, Kyoto 606-8501, Japan;

orcid.org/0000-0002-4414-2129

Author Contributions

All authors have given approval to the final version of the manuscript.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We thank JSPS KAKENHI Grant Number JP19H03350, JP19K22185, Platform Project for Supporting Drug Discovery and Life Science Research (BINDS) from AMED under Grant Number JP20am0101092j0004, and the Uehara Memorial Foundation.

REFERENCES

- (1) (a) Berresheim, A. J.; Müllen, K. Polyphenylene Nanostructures. *Chem. Rev.* **1999**, *99*, 1747–1786. (b) Watson, M. D.; Fechtenkötter, A.; Müllen, K. Big Is Beautiful—“Aromaticity” Revisited from the Viewpoint of Macromolecular and Supramolecular Benzene Chemistry. *Chem. Rev.* **2001**, *101*, 1267–1300. (c) Bendikov, M.; Wudl, F.; Perepichka, D. F. Tetrathiafulvalenes, Oligoacenenes, and Their Buckminsterfullerene Derivatives: The Brick and Mortar of Organic Electronics. *Chem. Rev.* **2004**, *104*, 4891–4945. (d) Anthony, J. E. The larger Acenes: Versatile Organic Semiconductors. *Angew. Chem., Int. Ed.* **2008**, *47*, 452–483. (e) Figueira-Duarte, T. M.; Müllen, K. Pyrene-Based Materials for Organic Electronics. *Chem. Rev.* **2011**, *111*, 7260–7314. (f) Sun, Z.; Ye, Q.; Chi, C.; Wu, J. Low Band Gap Polycyclic Hydrocarbons: From Closed-Shell Near Infrared Dyes and Semiconductors to Open-Shell Radicals. *Chem. Soc. Rev.* **2012**, *41*, 7857–7889. (g) Jiang, W.; Li, Y.; Wang, Z.-H. Heteroarenes as High Performance Organic Semiconductors. *Chem. Soc. Rev.* **2013**, *42*, 6113–6127.

(2) (a) Pashin, Y. V.; Bakhitova, L. M. Mutagenic and Carcinogenic Properties of Polycyclic Aromatic Hydrocarbons. *Environ. Health Perspect.* **1979**, 185–189. (b) da Silva Junior, F. C.; Felipe, M. B. M. C.; de Castro, D. E. F.; da Silva Araújo, S. C.; Sisenando, H. C. N.; de Medeiros, S. R. B. A. Look beyond the Priority: A systematic Review of the Genotoxic, Mutagenic, and Carcinogenic Endpoints of Non-priority PAHs. *Environ. Pollut.* **2021**, 278, 116838.

(3) Mohan, S. B.; Kumar, V. R.; Venkata, B.; Krishna, B. B.; Preetismita, B. Polyaromatic Hydrocarbons (PAHs): Structures, Synthesis and Biological Profile. *Curr. Org. Synth.* **2020**, 17, 625–640.

(4) (a) Niko, Y.; Hiroshige, Y.; Kawauchi, S.; Konishi, G.-i. Additional Insights into Luminescence Process of Polycyclic Aromatic Hydrocarbons with Carbonyl Groups: Photophysical Properties of Secondary *N*-Alkyl and Tertiary *N,N*-Dialkyl Carboxamides of Naphthalene, Anthracene, and Pyrene. *J. Org. Chem.* **2012**, 77, 3986–3996. (b) Achten, C.; Andersson, J. T. Overview of Polycyclic Aromatic Compounds (PAC). *Polycyclic Aromat. Compd.* **2015**, 35, 177–186. (c) Kroonblawd, M. P.; Lindsey, R. K.; Goldman, N. Synthesis of Functionalized Nitrogen-containing Polycyclic Aromatic Hydrocarbons and Other Prebiotic Compounds Impacting Glycine Solutions. *Chem. Sci.* **2019**, 10, 6091–6098. (d) Yang, X.; Hoffmann, M.; Rominger, F.; Kirchbaum, T.; Dreuw, A.; Mastalerz, M. Functionalized Contorted Polycyclic Aromatic Hydrocarbons by a One-step Cyclopentannulation and Regioselective Triflyoxylation. *Angew. Chem., Int. Ed.* **2019**, 58, 10650–10654.

(5) Nagamoto, Y.; Yamaoka, Y.; Fujimura, S.; Takemoto, Y.; Takasu, K. Synthesis of Functionalized Polycyclic Aromatic Compounds via a Formal [2 + 2]-Cycloaddition. *Org. Lett.* **2014**, 16, 1008–1011.

(6) (a) Nagamoto, Y.; Hattori, A.; Takeya, H.; Takemoto, Y.; Takasu, K. pH-Sensitive DNA Cleaving Agents: In situ Activation by Ring Contraction of Benzo-fused Cyclobutanols. *Chem. Commun.* **2013**, 49, 2622–2624. (b) Yamaoka, Y.; Taniguchi, M.; Yamada, K.; Takasu, K. Asymmetric Total Synthesis of Tylophorine via a Formal [2+2] Cycloaddition Followed by Migrative Ring Opening of a Cyclobutane. *Synthesis* **2015**, 47, 2819–2825. (c) Yamaoka, Y.; Taniguchi, M.; Yamada, K.; Takasu, K. Total Synthesis of Phenanthroquinolizidine Alkaloid Cryptopleurine and Phenanthroindolizidine Alkaloid Tylophorine. *Heterocycles* **2018**, 97, 292–305. (d) Ogawa, N.; Yamaoka, Y.; Takikawa, H.; Takasu, K. Synthesis of

Polycyclic Spirocarbocycles via Acid-Promoted Ring-Contraction/Dearomative Ring-Closure Cascade of Oxapropellanes. *Org. Lett.* **2019**, *21*, 7563–7567.

(7) Ogawa, N.; Yamaoka, Y.; Takikawa, H.; Tsubaki, K.; Takasu, K. Synthesis and Properties of Tribenzocarbazoles via an Acid-Promoted Retro (2+2)-Cycloaddition of Azapropellanes. *J. Org. Chem.* **2018**, *83*, 7994–8002.

(8) Selected examples for the synthesis PAHs from cyclobutanols: (a) Yu, J.; Yan, H.; Zhu, C. Synthesis of Multiply Substituted Polycyclic Aromatic Hydrocarbons by Iridium-Catalyzed Annulation of Ring-Fused Benzocyclobutenol with Alkyne through C–C Bond Cleavage. *Angew. Chem. Int. Ed.* **2016**, *55*, 1143–1146. (b) Mao, Y.; Zhu, C. C–C Bond (Hetero)arylation of Ring-Fused Benzocyclobutenols and Application in the Assembly of Polycyclic Aromatic Hydrocarbons. *J. Org. Chem.* **2017**, *82*, 9133–9143.

(9) Selected reviews: (a) Marek, I.; Masarwa, A.; Delaye, P.-O.; Liebeling, M. Selective Carbon-Carbon Bond Cleavage for Stereoselective Synthesis of Acyclic Systems. *Angew. Chem., Int. Ed.* **2015**, *54*, 414–429. (b) Ren, R.; Zhu, C. Radical-Mediated Ring-Opening Functionalization of Cyclobutanols: A Shortcut to γ -Substituted Ketones. *Synlett* **2016**, 1139–1144. (c) Murakami, M.; Ishida, N. β -Scission of Alkoxy Radicals in Synthetic Transformations. *Chem. Lett.* **2017**, *46*, 1692–1700. (d) Wu, X.; Zhu, C. Recent Advances in Ring-Opening Functionalization of Cycloalkanols by C–C σ -Bond Cleavage. *Chem. Rec.* **2018**, *18*, 587–598. (e) Wu, X.; Zhu, C. Recent Advances in Alkoxy Radical-Promoted C–C and C–H bond Functionalization Starting from Free Alcohols. *Chem. Commun.* **2019**, *55*, 9747–9756.

(10) Selected examples. (a) Kobayashi, K.; Itoh, M.; Suginome, H. A New Synthesis of Phthalides through β -Scission of Benzocyclobutenol Hypoiodites. *Tetrahedron Lett.* **1987**, *28*, 3369–3372. (b) Suginome, H.; Takeda, T.; Itoh, M.; Nakayama, Y.; Kobayashi, K. Photoinduced Molecular Transformations. Part 152. Ring Expansion Based on a Sensitized [2 + 2] Photoaddition of Enol Ethers of Cyclic Ketones with Olefins, Followed by a β -Scission of Alkoxy Radicals Generated from the Resulting Cyclobutanols. Two-Carbon Ring Expansion of β -Indanone, β -Tetralone and β -Suberone. *J. Chem. Soc., Perkin Trans. I* **1995**, 49–61. (c) Uemura, S.; Ohe, K.; Nishimura, T. Oxidative Transformation of *tert*-Cyclobutanols by Palladium Catalysis under Oxygen Atmosphere. *J. Org. Chem.* **2001**, *66*, 1455–1465. (d) Takasu, K.; Nagao, S.; Ihara,

M. Synthesis of Medium-sized γ -Haloketones by Radical Mediated Ring-opening Reaction of Lewis Acid Catalyzed (2+2)-Cycloadducts. *Tetrahedron Lett.* **2005**, *46*, 1005–1008. (e) Fujioka, H.; Komatsu, H.; Miyoshi, A.; Murai, K.; Kita, Y. Phenyliodine Diacetate-Mediated Oxidative Cleavage of Cyclobutanols Leading to γ -Hydroxy Ketones. *Tetrahedron Lett.* **2011**, *52*, 973–975. (f) Ren, R.; Zhao, H.; Huan, L.; Zhu, C. Manganese-Catalyzed Oxidative Azidation of Cyclobutanols: Regiospecific Synthesis of Alkyl Azides by C-C Bond Cleavage. *Angew. Chem., Int. Ed.* **2015**, *54*, 12692–12696. (g) Kurouchi, H.; Andujar-De Sanctis I. L.; Singleton D. A. Controlling Selectivity by Controlling Energy Partitioning in a Thermal Reaction in Solution. *J. Am. Chem. Soc.* **2016**, *138*, 14534–14537. (h) Huan, L.; Zhu, C. Manganese-Catalyzed Ring-Opening Chlorination of Cyclobutanols: Regiospecific Synthesis of γ -Chloroketones. *Org. Chem. Front.* **2016**, *3*, 1467–1471. (i) Lopez, M. M.; Jamey, N.; Pinet, A.; Figadere, B.; Ferrié, L. Oxidative Ring Expansion of Cyclobutanols: Access to Functionalized 1,2-Dioxanes, *Org. Lett.* **2021**, *23*, 1626–1631. (11) Kirihaara, M.; Okada, T.; Sugiyama, Y.; Akiyoshi, M.; Matsunaga, T.; Kimura, Y. Sodium Hypochlorite Pentahydrate Crystals (NaOCl·5H₂O): A Convenient and Environmentally Benign Oxidant for Organic Synthesis. *Org. Proc. Res. Dev.* **2017**, *21*, 1925–1937. (12) (a) Yin, Q.; Wang, S.-G.; Liang, X.-W.; Gao, D.-W.; Zheng, J.; You, S.-Li. Organocatalytic Asymmetric Dearomatization of Naphthols. *Chem. Sci.* **2015**, *6*, 4179–4183. (b) Wang, P.; Wang, J.; Wang, L.; Li, D.; Wang, K.; Liu, Y.; Zhu, H.; Liu, X.; Yang, D.; Wang, L. Asymmetric Dearomative Halogenation of b-Naphthols: The Axial Chirality Transfer Reaction. *Adv. Synth. Catal.* **2018**, *360*, 401–405. (c) Uyanik, M.; Sahara, N.; Ishihara, K. Regioselective Oxidative Chlorination of Arenols Using NaCl and Oxone. *Eur. J. Org. Chem.* **2019**, 27–31.