

CR manifolds in Grauert tubes

東北大学大学院理学研究科数学専攻 小泉 英介* (Eisuke Koizumi)

Mathematical Institute, Tohoku University

In this article, we introduce a result on the logarithmic term of the Szegő kernel on the boundary of two-dimensional Grauert tubes. In Section 1, we give the definition of Grauert tube and some examples. In Section 2, we introduce a result in [5]. This result plays very important roles in studying CR manifolds in Grauert tubes. Finally, in Section 3, we state the main theorem in [8] and some remarks.

1 The definition of Grauert tube and examples

Let (X, g) be an n -dimensional complete C^ω Riemannian manifold, and let $\gamma : \mathbb{R} \rightarrow X$ be a geodesic. Then we define the mapping $\psi_\gamma : \mathbb{C} \rightarrow TX$ by

$$\psi_\gamma(\sigma + i\tau) := \tau \dot{\gamma}(\sigma).$$

Definition 1.1. Let $T^r X := \{v \in TX \mid g(v, v) < r^2\}$, where $0 < r \leq \infty$. A complex structure on $T^r X$ is said to be adapted if ψ_γ is holomorphic for every geodesic γ on X .

If an adapted complex structure exists, then it is uniquely determined (see [9]).

The Grauert tube of radius r over X is the manifold $T^r X$ with the adapted complex structure. X is called the center of the Grauert tube.

Let $r_{\max}(X)$ be the maximal radius r such that the adapted complex structure is defined on $T^r X$. It is known that $r_{\max}(X) > 0$ if X is compact or X is homogeneous.

Example 1.2. Let $X := \mathbb{R}^n$. Then $T^\infty \mathbb{R}^n$ is biholomorphic to $\mathbb{C}^n$.

Example 1.3. Let $X := S^n$, the unit sphere in $\mathbb{R}^{n+1}$. Then $T^\infty S^n$ is biholomorphic to the manifold $Q^n := \{z = (z_1, \dots, z_{n+1}) \in \mathbb{C}^{n+1} \mid z_1^2 + \dots + z_{n+1}^2 = 1\}$. We call Q^n the complex quadric.

Example 1.4. Let X be the n -dimensional real hyperbolic space with constant sectional curvature -1 . Then $T^{\pi/2} X$ is biholomorphic to B^n , the unit ball in $\mathbb{C}^n$. We note that $r_{\max}(X) = \pi/2$ (see [11, Theorem 2.5]).

* e-mail: s98m13@math.tohoku.ac.jp

2 CR manifolds in Grauert tubes

Let (X, g) be an n -dimensional compact C^ω Riemannian manifold, and let $T^r X$ be the Grauert tube. We define the mapping $\rho : T^r X \rightarrow \mathbb{R}$ by $\rho(v) := 2g(v, v)$ for $v \in T^r X$.

Theorem 2.1 ([5], [9]). *ρ has the following properties:*

- (1) ρ is strictly plurisubharmonic,
- (2) $X = \rho^{-1}(0)$,
- (3) the metric $ds_{T^r X}^2$ obtained from the Kähler form $i\partial\bar{\partial}\rho/2$ is compatible with g , that is, $ds_{T^r X}^2|_X = g$, and
- (4) $(\partial\bar{\partial}\sqrt{\rho})^n = 0$ in $T^r X - X$.

Let $\Omega_\epsilon := \{\rho < \epsilon^2\} \subset T^r X$, and let $M_\epsilon := \partial\Omega_\epsilon$. Then we see that M_ϵ is a strongly pseudoconvex CR manifold.

One of the interesting problems on Grauert tube is to study relations between M_ϵ and (X, g) . Several results have been known on this problem. Stenzel [10] studied orbits of the geodesic flow and chains. Kan [7] computed the Burns-Epstein invariant, and showed that M_{ϵ_1} and M_{ϵ_2} are not CR equivalent if $\epsilon_1 \neq \epsilon_2$ when $\dim X = 2$. This Kan's result is also true for $\dim X \geq 3$ (see [12]). This implies that there exist many CR manifolds in the Grauert tube. This fact is one of the reasons why we are interested in this problem.

3 Result

Let (X, g) be a two-dimensional compact Riemannian manifold, and let $T^r X$ be the Grauert tube. We put $\Omega_\epsilon := \{\rho < \epsilon^2\} \subset T^r X$ and $M_\epsilon := \partial\Omega_\epsilon$.

Let $\theta := \iota_\epsilon^*(-i\partial\rho)$, where ι_ϵ is the embedding of M_ϵ in the Grauert tube. Then θ defines a pseudo-hermitian structure on M_ϵ . Let S_ϵ be the Szegő kernel with respect to the volume element $\theta \wedge d\theta$. Then by [2] and [1], the singularity of S_ϵ on the diagonal of M_ϵ is of the form

$$S_\epsilon(z, \bar{z}) = \varphi(z)\rho_\epsilon(z)^{-2} + \psi(z)\log\rho_\epsilon(z),$$

where $\varphi, \psi \in C^\infty(\overline{\Omega_\epsilon})$ and ρ_ϵ is a defining function of Ω_ϵ with $\rho_\epsilon > 0$ in Ω_ϵ .

Theorem 3.1 ([8]). *The boundary value of the logarithmic term coefficient $\psi_0 = \psi|_{M_\epsilon}$ has the following asymptotic expansion as $\epsilon \rightarrow +0$:*

$$(3.1) \quad \psi_0 \sim \frac{1}{24\pi^2} \sum_{l=0}^{\infty} F_l^{\psi_0} \epsilon^{2l},$$

where $F_l^{\psi_0}(\lambda^2 g) = \lambda^{-2l-4} F_l^{\psi_0}(g)$ for $\lambda > 0$.

In particular, we have

$$(3.2) \quad F_0^{\psi_0} = -\frac{1}{10}\Delta k - \frac{2}{5}(\varepsilon^2 T^2 k)|_{\varepsilon=0},$$

where k is the scalar curvature, Δ is the Laplacian and T is the unique vector field on M_ε such that $\theta(T) = 1$ and $T \lrcorner d\theta = 0$.

We now make two remarks on the term $(\varepsilon^2 T^2 k)|_{\varepsilon=0}$. One is that we can regard this term as a function on the circle bundle over X , and it is not constant on each fiber of the bundle in general (see [8, Lemma 4.5]). This means that the value to which ψ_0 tends as $\varepsilon \rightarrow +0$ varies with the way ε goes to $+0$.

The other is that

$$(3.3) \quad \int_{M_\varepsilon} (\varepsilon^2 T^2 k)|_{\varepsilon=0} \theta \wedge d\theta = c\varepsilon^2 \int_X \Delta k dV + O(\varepsilon^3),$$

where c is a constant and dV is the volume form on X (see also [7]). It follows from (3.1)–(3.3) and $\int_X \Delta k dV = 0$ that the coefficient of ε^2 in the integral

$$\int_{M_\varepsilon} \psi_0 \theta \wedge d\theta$$

is equal to 0. This is not contradict to the fact that the integral above is equal to 0.

Finally, we note that ψ_0 is a constant multiple of the Q -curvature of three-dimensional CR manifolds (see [3], [4] and [6]). In conformal geometry, there has been great progress recently in understanding the Q -curvature and its geometric meaning in low dimensions. However, roles of Q -curvature in CR geometry are not clear. We hope that this result will become an approach to studying CR Q -curvature.

References

- [1] L. Boutet de Monvel and J. Sjöstrand, *Sur la singularité des noyau de Bergman et de Szegő*, Astérisque 34–35 (1976), 123–164.
- [2] C. Fefferman, *The Bergman kernel and biholomorphic mappings of pseudoconvex domains*, Invent. Math. 26 (1974), 1–65.
- [3] C. Fefferman and K. Hirachi, *Ambient metric construction of Q -curvature in conformal and CR geometries*, preprint.
- [4] A. R. Gover and C. R. Graham, *CR invariant powers of the sub-Laplacian*, preprint.

- [5] V. Guillemin and M. Stenzel, *Grauert tubes and the homogeneous Monge-Ampère equation*, J. Differ. Geom. 34 (1993), 561–570.
- [6] K. Hirachi, *Scalar pseudo-hermitian invariantss and the Szegő kernel on three-dimensional CR manifolds*, “Complex Geometry,” Lecture Notes in Pure and Appl. Math. 143 (1993), 67–76.
- [7] S. -J. Kan, *The asymptotic expansion of a CR invariant and Grauert tubes*, Math. Ann. 304 (1996), 63–92.
- [8] E. Koizumi, *The logarithmic term of the Szegő kernel on the boundary of two-dimensional Grauert tubes*, in preparation.
- [9] L. Lempert and R. Szöke, *Global solutions of the homogeneous complex Monge-Ampère equations and complex structures on the tangent bundle of Riemannian manifolds*, Math. Ann. 290 (1991), 689–712.
- [10] M. Stenzel, *Orbits of the geodesic flow and chains on the boundary of a Grauert tube*, Math. Ann. 322 (2002), 383–399.
- [11] R. Szöke, *Complex structures on tangent bundles of Riemannian manifolds*, Math. Ann. 291 (1991), 409–428.
- [12] R. Szöke, *Adapted complex structures and Riemannian homogeneous spaces*, Ann. Polan. Math. 70 (1998), 215–220.