

A CHARACTERISTIC CAUCHY PROBLEM OF NON-LERAY TYPE IN THE COMPLEX DOMAIN

HIDESHI YAMANE¹

山根英司

§0. Introduction

We consider a Cauchy problem in the complex domain. It is assumed to be a *characteristic* problem in the sense that the characteristic points form a submanifold T (of codimension 1) of the initial hypersurface S .

Since Leray, the studies on this subject dealt with the cases where the solution is singular on a characteristic hypersurface *tangent* to S along T . See [L], [G-K-L], [H], [D], [O-Y] and [Y].

In the present paper, we consider a totally different situation: all the characteristic hypersurfaces issuing from T are *transversal* with S .

First we give two examples to show that in this kind of characteristic Cauchy problem, the solution can be singular on the above-mentioned characteristic hypersurfaces even when all the Cauchy data are regular. Next, we consider a (ramified) Cauchy problem for a certain class of operators including the examples. We perform a singular change of coordinates and reduce our problem to results of Wagschal.

§1. Examples with holomorphic data

In a neighborhood of the origin of $\mathbb{C}_t \times \mathbb{C}_x \times \mathbb{C}_z$, let us consider Cauchy problems for the operators Q_1 and Q_2 defined by

$$Q_1 = (xD_t + tD_x)D_t, \quad Q_2 = Q_1 - xt^2D_z^2.$$

1991 Mathematics Subject Classification. Primary 35A20

¹Department of Mathematical Sciences, University of Tokyo
3-8-1, Komaba, Meguro-ku, Tokyo 153, Japan

Current address: Department of Mathematics, Chiba Institute of Technology, 2-1-1 Shibazono, Narashino, Chiba 275, Japan.

We are going to solve, for $j = 1$ or 2 ,

$$\begin{cases} Q_j u(t, x, z) = 0 \\ u|_S = -\frac{\pi i}{2} x^2 \\ D_t u|_S = ix \\ S = \{t = 0\} \end{cases}$$

On the initial hypersurface S , the characteristic points form a submanifold $T = \{t = x = 0\}$. The hypersurfaces $\{x = 0\}$, $\{x = t\}$ and $\{x = -t\}$ are characteristic hypersurfaces issuing from T . They are transversal with S . Although the data are holomorphic in a neighborhood of the origin, the solution u is singular on the three characteristic hypersurfaces. In fact, we have

$$u = \frac{x^2}{2} \left\{ \frac{t}{x} \sqrt{\left(\frac{t}{x}\right)^2 - 1} - \log\left(\frac{t}{x} + \sqrt{\left(\frac{t}{x}\right)^2 - 1}\right) \right\}.$$

Since we are dealing with a multi-valued function, we have to clarify the definition of the restriction on S . Its precise meaning is that we choose a point p of S and that the initial condition is satisfied by the germ of u at p .

We will prove for a class of operators including Q_1 and Q_2 that *the singular support of the solution is contained in this kind of characteristic hypersurfaces* when the data are arbitrary holomorphic functions. As a matter of fact, we can generalize this result to the case of ramified data.

§2. Main result

In a neighborhood of the origin of $\mathbb{C}_t \times \mathbb{C}_y \times \mathbb{C}_z^n$, let us consider a second order operator $P(t, y, z; D_t, D_y, D_z)$ with holomorphic coefficients whose principal symbol $\sigma(P)$ is factorized into the form

$$\sigma(P)(t, y, z; \tau, \eta, \zeta) = \prod_{i=0,1} (\tau - \lambda_i(t, y, z; \eta, \zeta)),$$

where τ, η and $\zeta = (\zeta_1, \dots, \zeta_n)$ are the dual variables of t, y and z respectively.

We assume the following two conditions (1) and (2).

$$(1) \quad \begin{cases} \lambda_0(t, 0, z; 1, 0, \dots, 0) = 0 \\ \lambda_1(t, y, z; 1, 0, \dots, 0) = -qt^{q-1}, \\ q \text{ is an integer } \geq 2. \end{cases}$$

CHARACTERISTIC CAUCHY PROBLEM

(2) For $i = 0, 1$, the function $(\eta, \zeta) \mapsto \lambda_i(t, y, z; \eta, \zeta)$ is linear.

The most simple example is

$$\lambda_0 = 0 \text{ or } y\eta, \quad \lambda_1 = -qt^{q-1}\eta.$$

Now we consider, in a neighborhood of the origin of $\mathbb{C}_t \times \mathbb{C}_x \times \mathbb{C}_z^n$, an operator Q with holomorphic coefficients defined by

$$Q(t, x, z; D_t, D_x, D_z) = x^{2q-1}P(t, x^q, z; D_t, \frac{1}{qx^{q-1}}D_x, D_z).$$

Sometimes the exponent $2q-1$ is larger than necessary to erase negative powers of x . For example, if

$$P(t, y, z; D_t, D_y, D_z) = P(t, y; D_t, D_y) = (D_t + qt^{q-1}D_y)D_t,$$

then

$$x^{q-1}P(t, x^q; \frac{1}{qx^{q-1}}D_x) = (x^{q-1}D_t + t^{q-1}D_x)D_t.$$

When $q = 2$, this is nothing but Q_1 which we studied before.

For the purpose of formulating a Cauchy problem, put $S = \{t = 0\} \subset \mathbb{C}_t \times \mathbb{C}_x \times \mathbb{C}_z^n$, which is the initial hypersurface. It is easy to see that $T = \{t = x = 0\}$ is formed by the characteristic points of Q on S . By the condition (1), the hypersurfaces

$$K_j = \{x = \exp(j \frac{2\pi i}{q}) \cdot t\} \quad (j = 0, \dots, q-1), \quad K_q = \{x = 0\}$$

are characteristic hypersurfaces of Q issuing from T .

We then consider a ramified characteristic Cauchy problem in an open connected neighborhood Ω of the origin of $\mathbb{C}_t \times \mathbb{C}_x \times \mathbb{C}_z^n$:

$$(CP) \begin{cases} Q(t, x, z; D_t, D_x, D_z)u(t, x, z) = 0, \\ D_t^h u(t, x, z)|_S = w_h(x, z), \quad h = 0, 1. \end{cases}$$

Here we assume that there exists a point $p \in \Omega \cap (S \setminus T)$ such that for $h = 0, 1$, the function w_h is holomorphic in a neighborhood (relative to S) of the point p and can be analytically continued along all the paths from p in $\Omega \cap (S \setminus T)$ (that is, w_h is holomorphic in the universal covering space of $\Omega \cap (S \setminus T)$).

Since $p \notin T$, the usual Cauchy-Kowalevski theorem is valid there. (CP) admits a unique holomorphic solution u in a neighborhood of the point p .

We are going to prove the

Theorem 1.

There exists an open connected neighborhood Ω' of the origin of $\mathbb{C}_t \times \mathbb{C}_x \times \mathbb{C}_z^n$ such that the solution u of (CP) can be analytically continued to the universal covering space of $\Omega' \setminus \bigcup_{j=0}^q K_j$.

Of course this conclusion holds true when all the data are regular.

Proof.

Put $x = y^{1/q}$. Then $D_y = \frac{1}{qx^{q-1}} D_x$. Therefore

$$Q(t, x, z; D_t, D_x, D_z) = y^{\frac{2q-1}{q}} P(t, y, z; D_t, D_y, D_z).$$

We reduce (CP) to the following *noncharacteristic* ramified Cauchy problem, which has been solved by Wagschal in [W2].

$$(CP') \begin{cases} P(t, y, z; D_t, D_y, D_z) u(t, y^{1/q}, z) = 0, \\ D_t^h u(t, y^{1/q}, z)|_{t=0} = w_h(y^{1/q}, z), \quad h = 0, 1. \end{cases}$$

The function $w_h(y^{1/q}, z)$ is holomorphic in the universal covering space of $\{(y, z) \in \mathbb{C} \times \mathbb{C}^n; 0 < |y| \ll 1, |z| \ll 1\}$. ($a \ll 1$ means that $a \geq 0$ is sufficiently small).

Let $p' \in (\{0\} \times \mathbb{C}_y \times \mathbb{C}_z^n) \setminus \{y = 0\}$ be the point corresponding to p . Then (CP') admits a unique holomorphic solution $u(t, y^{1/q}, z)$ near p' . According to [W2], $u(t, y^{1/q}, z)$ can be analytically continued to the universal covering space of

$$\{(t, y, z) \in \mathbb{C} \times \mathbb{C} \times \mathbb{C}^n; |(t, y, z)| \ll 1\} \setminus (\{y = 0\} \cup \{y = t^q\}).$$

We finish the proof by coming back to the (t, x, z) -space. $\square$

Example.

We saw before that Q_1 was not quite the same as Q , but this does not cause any difficulty. The equation $Q_1 u = 0$ is equivalent to $x^2 Q_1 u = 0$. The operator $x^2 Q_1$ is nothing but Q .

This example suggests that the choice of the exponent of x in the definition of Q is not essential.

Remark 1.

CHARACTERISTIC CAUCHY PROBLEM

For convenience, put $y = z_0, \eta = \zeta_0$. Then, by virtue of Remarque 3.1 of [W2], (2) can be replaced by the following condition:

(3) There exists an integer $k, 0 \leq k \leq n$, such that for $i = 0, 1$, the function $\lambda_i(t, z_0, z; \zeta_0, \dots, \zeta_k, 0, \dots, 0)$ is linear in $(\zeta_0, \dots, \zeta_k)$ and does not depend on the variables $(z_{k+1}, \dots, z_n)$.

This enables us to treat Q_2 . In fact, when $n = 1, q = 2$, put

$$P(t, y, z; D_t, D_y, D_z) = D_t^2 + 2tD_tD_y - t^2D_z^2.$$

Then

$$\begin{aligned} \sigma(P) &= \tau^2 + 2t\tau\eta - t^2\zeta^2 \\ &= (\tau + t\eta)^2 - t^2(\eta^2 + \zeta^2) \\ &= \{\tau + t(\eta + \sqrt{\eta^2 + \zeta^2})\}\{\tau + t(\eta - \sqrt{\eta^2 + \zeta^2})\}, \\ Q_2 &= xP(t, x^2, z; D_t, \frac{1}{2x}D_x, D_z). \end{aligned}$$

Remark 2.

A singular change of coordinates was useful in some papers mentioned in the introduction ([L], [D], [O-Y] and [Y]). One introduces a new variable w by setting $w = (t - x^l)^{1/l}$ for some positive integer l . In the present paper, we have performed a different kind of singular change of coordinates.

§3. Inhomogeneous problem

If we choose a special class of P , we can treat an inhomogeneous problem. Assume that

$$\sigma(P)(t, y, z; \tau, \eta, \zeta) = \tau(\tau + qt^{q-1}\eta).$$

We employ the same notation as in §2. Let us consider:

$$(CP^i) \begin{cases} Q(t, x, z; D_t, D_x, D_z)u(t, x, z) = v(t, x, z), \\ D_t^h u(t, x, z)|_S = w_h(x, z), \quad h = 0, 1. \end{cases}$$

Here we assume that the function v is holomorphic in a neighborhood of p and can be analytically continued along all the paths from p in $\Omega \setminus \bigcup_{j=0}^q K_j$ (that is, v is holomorphic in the universal covering space of $\Omega \setminus \bigcup_{j=0}^q K_j$). Then we have

Theorem 2.

There exists an open connected neighborhood Ω' of the origin of $\mathbb{C}_t \times \mathbb{C}_x \times \mathbb{C}_z^n$ such that the solution u of (CP^i) can be analytically continued to the universal covering space of $\Omega' \setminus \bigcup_{j=0}^q K_j$.

Of course this conclusion holds true when all the data are regular.

Proof.

We have to solve

$$\begin{cases} P(t, y, z; D_t, D_y, D_z)u(t, y^{1/q}, z) = y^{-\frac{2q-1}{q}} v(t, y^{1/q}, z), \\ P = D_t(D_t + qt^{q-1}D_y) + \text{lower}, \\ D_t^h u(t, y^{1/q}, z)|_{t=0} = w_h(y^{1/q}, z), \quad h = 0, 1. \end{cases}$$

Since $v(t, x, z)$ is holomorphic in the universal covering space of $\Omega \setminus \bigcup_{j=0}^q K_j$, the function $y^{-\frac{2q-1}{q}} v(t, y^{1/q}, z)$ is holomorphic in the universal covering space of

$$\{(t, y, z) \in \mathbb{C} \times \mathbb{C} \times \mathbb{C}^n; |(t, y, z)| \ll 1\} \setminus (\{y = 0\} \cup \{y = t^q\}).$$

This noncharacteristic inhomogeneous problem has been solved in [W1]. $\square$

§4. Geometry

What distinguishes the present study from conventional ones is the absence of singularities on a hypersurface tangent to the initial hypersurface S . It is explained by the following

Proposition.

Under the assumption (1), there is no characteristic hypersurface of Q that is tangent to S along T .

Proof.

We have

$$\begin{aligned} \sigma(Q)(t, x, z; \tau, \xi, \zeta) &= x^{2q-1} \prod_{i=0,1} \left\{ \tau - \lambda_i(t, x^q, z; \frac{1}{qx^{q-1}}\xi, \zeta) \right\} \\ &= x \prod_{i=0,1} \left\{ x^{q-1}\tau - \lambda_i(t, x^q, z; \frac{1}{q}\xi, x^{q-1}\zeta) \right\}. \end{aligned}$$

CHARACTERISTIC CAUCHY PROBLEM

It is easy to see that S itself is not a characteristic hypersurface. A hypersurface $\neq S$ which is tangent to S along T has an expression of the form:

$$\varphi = t + x^N \psi(x, z) = 0, \quad N \geq 2$$

where ψ is a holomorphic function with $\psi(0, z) \neq 0$.

We have

$$\begin{aligned} & \sigma(Q)(t, x, z; \text{grad } \varphi) \\ &= x \prod_{i=0,1} [x^{q-1} - \lambda_i(t, x^q, z; \frac{1}{q} N x^{N-1} \psi(x, z) + \frac{1}{q} x^N D_x \psi, x^{N+q-1} D_z \psi)]. \end{aligned}$$

For a generic z we have $\psi(0, z) \neq 0$. We fix such a z . Obviously $\psi(x, z) \neq 0$ holds if $|x| \ll 1$. Then it follows that

$$\begin{aligned} & \sigma(Q)(t, x, z; \text{grad } \varphi) \\ &= x \prod_{i=0,1} [x^{q-1} - \frac{1}{q} N x^{N-1} \psi \lambda_i(t, x^q, z; 1 + \frac{x}{N\psi} D_x \psi, \frac{qx^q}{N\psi} D_z \psi)] \\ &= x \prod_{i=0,1} [x^{q-1} - \frac{1}{q} N x^{N-1} \psi (1 + \frac{x}{N\psi} D_x \psi) \lambda_i(t, x^q, z; 1, (1 + \frac{x}{N\psi} D_x \psi)^{-1} \frac{qx^q}{N\psi} D_z \psi)]. \end{aligned}$$

The assumption (1) implies that as x tends to zero

$$\begin{aligned} \lambda_0(t, x^q, z; 1, (1 + \frac{x}{N\psi} D_x \psi)^{-1} \frac{qx^q}{N\psi} D_z \psi) &= O(x^q) \\ \lambda_1(t, x^q, z; 1, (1 + \frac{x}{N\psi} D_x \psi)^{-1} \frac{qx^q}{N\psi} D_z \psi) &= -qt^{q-1} + O(x^q). \end{aligned}$$

Therefore by restricting them on the hypersurface $\{\varphi = 0\}$, we obtain

$$\begin{aligned} \lambda_0(t, x^q, z; 1, (1 + \frac{x}{N\psi} D_x \psi)^{-1} \frac{qx^q}{N\psi} D_z \psi)|_{\varphi=0} &= O(x^q) \\ \lambda_1(t, x^q, z; 1, (1 + \frac{x}{N\psi} D_x \psi)^{-1} \frac{qx^q}{N\psi} D_z \psi)|_{\varphi=0} &= -q(-x^N \psi)^{q-1} + O(x^q) = O(x^q). \end{aligned}$$

Hence $\sigma(Q)|_{\varphi=0}$ is different from zero if $0 < |x| \ll 1$. Thus $\{\varphi = 0\}$ is not a characteristic hypersurface. $\square$

HIDESHI YAMANE

REFERENCES

- [D] J.Dunau, *Un Problème de Cauchy Caractéristique*, J. Math. pures et appl. **69** (1990), 369-402.
- [G-K-L] L. Gårding, T.Kotake and J.Leray, *Problème de Cauchy, I bis et VI*, Bull. Soc. Math. de France **92** (1964), 263-361.
- [H] Y.Hamada, *Les singularités des solutions du problème de Cauchy à données holomorphes*, Recent Developments in Hyperbolic Equations (Proceedings of the conference on Hyperbolic Equations, University of Pisa, 1987), Pitman Research Notes in Mathematics Series 183 (1988), Longman, 82-95.
- [L] J.Leray, *Uniformisation de la solution du problème linéaire analytique de Cauchy près de la variété qui porte les données de Cauchy*, Bull. Soc. math. France **85** (1957), 389-429.
- [O-Y] Y.Okada, H.Yamane, *A characteristic Cauchy problem in the complex domain*, to appear in J. Math. pures et appl..
- [W1] C.Wagschal, *Problème de Cauchy ramifié pour une class d'opérateurs à caractéristique tangentes (I)*, J. Math. pures et appl. **67** (1988), 1-21.
- [W2] ———, *Problème de Cauchy ramifié: opérateurs à caractéristiques tangentes ayant un contact d'ordre constant*, Recent Developments in Hyperbolic Equations (Proceedings of the conference on Hyperbolic Equations, University of Pisa, 1987), Pitman Research Notes in Mathematics Series 183 (1988), Longman, 367-385.
- [Y] H.Yamane, *The essential singularity of the solution of a ramified characteristic Cauchy problem*, Publ. RIMS, Kyoto Univ. (to appear).