

## Nothernmost geographical distribution of *Veleva veleva* (Hydrozoa: Porpitiidae) in the Middle of the North Pacific

Shin Kubota<sup>1</sup> and Yukimitsu Imahara<sup>2</sup>

<sup>1</sup> Seto Marine Biological Laboratory, Field Science Education and Research Center,  
Kyoto University, 459 Shirahama, Nishimuro, Wakayama, 649-2211 Japan

<sup>2</sup> Wakayama Laboratory, Biological Institute on Kuroshio,  
300-11 Kire, Wakayama City, Wakayama, 640-0351, Japan

**Abstract.** The northernmost distribution of *Veleva veleva* (Hydrozoa: Porpitiidae) is recorded in the middle of the North Pacific (39° 57.87' N, 157° 40.46' W). The specimen with symbiotic algae is rather large (63 mm in length, 38 mm in width) and mature despite of the low temperature.

**Key words:** *Veleva*, occurrence, North Pacific, mature, symbiotic algae, temperature.

### Introduction

*Veleva veleva* (Linnaeus, 1758), by-the-wind-sailor, is floating on the sea surface and distributed in the temperate to tropical zone of the Pacific mainly along the warm current except tropical middle zone of the North Pacific (Bieri, 1977). It is sometimes stranded on the Japanese coasts in spring to autumn, and all the specimens are small (Bieri, 1970; Kubota, 2003, 2004). We found a large specimen in cool waters in the middle of the North Pacific as the northernmost distributional record there and reported its occurrence.

### Description

The specimen (Fig. 1) was collected by a net on board (Fig. 2: 39° 57.87' N, 157° 40.46' W) and photographed, then preserved in the formalin

seawater in situ soon after collection on June 24, 2010. The specimen, 63 mm in length and 38 mm in width, is blue in color and matured bearing many medusa buds. Many symbiotic algae were found inside the medusa buds despite the water temperature is rather cold (14.6°C) in a cool region (air temperature is 16.3°C).

### Remarks

In the western north Pacific, *Veleva* is usually up to 60 mm in length (Bieri, 1977: p. 66), therefore the present specimen is rather large, but in the eastern north Pacific it is up to 105 mm in length (Bieri, 1977: p. 65) and grows bigger than the former. On the other hand, it is often reported that coral bleaching was induced by low water temperatures, for example in Kii Peninsula, this has been confirmed three times since 1971 due to a long lasting under 14°C (20 days or more) (Nomura, 2004). It is assumed that the present large specimen may be carried to the finding site by the Kuroshio around the col-

---

\*Corresponding author :  
shkubota@medusanpolyp.mbox.media.kyoto-u.ac.jp

lected date judging from its size and possession of symbiotic algae.



Fig. 1. The large specimen of *Veleva veleva* collected in the middle of the North Pacific (in life, 63 mm in length).




Fig. 2. Geographical distribution of *Veleva veleva* (after Bieri, 1977) and the present record (★).

### Acknowledgements

We thank Mr Masahisa Muratsuka, a crew of *R/*

*V Kaiyomaru* of the Fisheries Agency of Japan, for collecting the animal on board.

### References

- Bieri, R. 1970. Droplets from the plankton net XXIV. The predominant type of *Veleva* in Japanese waters. Publ. Seto Mar. Biol. Lab., 17: 303-304.
- Bieri, R. 1977. The ecological significance of seasonal occurrence and growth rate of *Veleva* (Hydrozoa). Publ. Seto Mar. Biol. Lab., 24: 63-76.
- Kubota, S. 2003. Seasonal occurrence of strandings of large jellyfish on "Kitahama" beach at Banshozaki, Shirahama, Wakayama Prefecture, Japan. J. Jap. Drif. Soc., 1: 21-24. (In Japanese with English abstract)
- Kubota, S. 2004. Extraordinary seasonal pattern of strandings of large jellyfish on "Kitahama" beach at Banshozaki, Shirahama, Wakayama Prefecture, Japan – supplementary report with comparison to earlier records. J. Jap. Drif. Soc., 2: 25-28. (In Japanese with English abstract)
- Nomura, K. 2004. Status of coral reefs around the country. Kii Peninsula. pp. 250-257. In "The Japanese Coral Reef Society and Ministry of the Environment eds. Coral Reefs of Japan". 356 pp. Ministry of the Environment, Tokyo.

(Received May 30, 2011; Accepted June 23, 2011)