

ON THE DECOMPOSITION OF THE HOCHSCHILD COHOMOLOGY GROUP OF A MONOMIAL ALGEBRA SATISFYING A SEPARABILITY CONDITION

AYAKO ITABA

Department of Mathematics,
Tokyo University of Science

ABSTRACT. In this note, we consider the finite connected quiver $\mathcal{Q}$ having two subquivers $\mathcal{Q}^{(1)}$ and $\mathcal{Q}^{(2)}$ with $\mathcal{Q} = \mathcal{Q}^{(1)} \cup \mathcal{Q}^{(2)} = (\mathcal{Q}_0^{(1)} \cup \mathcal{Q}_0^{(2)}, \mathcal{Q}_1^{(1)} \cup \mathcal{Q}_1^{(2)})$. Suppose that $\mathcal{Q}^{(i)}$ is not a subquiver of $\mathcal{Q}^{(j)}$ where $\{i, j\} = \{1, 2\}$. For a monomial algebra $\Lambda = k\mathcal{Q}/I$ obtained by the quiver $\mathcal{Q}$, when the set $AP(n)$ ($n \geq 0$) of overlaps constructed inductively by linking generators of I satisfies a certain separability condition, we propose the method so that we construct a minimal projective resolution of Λ as a right Λ^e -module and calculate the Hochschild cohomology group of Λ .

1. INTRODUCTION

First of all, we recall the definition of Hochschild cohomology (see [S]). For a finite-dimensional algebra A over a field k , the Hochschild cohomology groups $\mathrm{HH}^n(A)$ of A is defined by

$$\mathrm{HH}^n(A) := \mathrm{Ext}_{A^e}^n(A, A) \quad (n \geq 0),$$

where $A^e := A^{\mathrm{op}} \otimes_k A$ is the enveloping algebra of A . Note that there is a natural one to one correspondence between the family of A - A -bimodules and that of right A^e -modules. Moreover, the Hochschild cohomology rings $\mathrm{HH}^*(A)$ of A is the graded algebra defined by

$$\mathrm{HH}^*(A) := \mathrm{Ext}_{A^e}^*(A, A) = \bigoplus_{i \geq 0} \mathrm{Ext}_{A^e}^i(A, A)$$

with the Yoneda product.

The low-dimensional Hochschild cohomology groups are described as follows:

- $\mathrm{HH}^0(A) = Z(A)$ is the center of A .
- $\mathrm{HH}^1(A)$ is the space of derivations modulo the inner derivations. A derivation is a k -linear map $f : A \rightarrow A$ such that $f(ab) = af(b) + f(a)b$ for all $a, b \in A$. A derivation $f : A \rightarrow A$ is an inner derivation if there is some $x \in A$ such that $f(a) = ax - xa$ for all $a \in A$.
- $\mathrm{HH}^2(A)$ measures the infinitesimal deformations of the algebra A .

One important property of Hochschild cohomology is its invariance under Morita equivalence, stable equivalence of Morita type and derived equivalence.

In general, it is not easy to calculate the Hochschild cohomology of a finite-dimensional algebra. In order to calculate the Hochschild cohomology groups of a quiver algebra, can we use calculations of the Hochschild cohomology groups of quiver algebras obtained by subquivers of the original quiver? Hence, we consider Hochschild cohomology of an algebra obtained by “linking” two algebras as the analogy of the following two studies. In [H], for

This note is a survey article of a joint work with Takahiko Furuya and Katsunori Sanada. See [IFS] for the detail.

a finite-dimensional algebra A and $M \in \text{mod } A$, Happel studied the one-point extensions $B = A[M] = \begin{pmatrix} A & M \\ 0 & k \end{pmatrix}$ of A and show that there exists the long exact sequence connecting the Hochschild cohomology of A and B . In [BO], for a finite-dimensional algebra over a field k , Bergh and Oppermann studied the Hochschild cohomology of twisted tensor products and applied this to the class of finite-dimensional algebras known as quantum complete intersections.

Let k be an algebraically closed field and $\mathcal{Q}$ a finite connected quiver. Then $k\mathcal{Q}$ denotes the path algebra of $\mathcal{Q}$ over k in this paper. Let I be an admissible ideal of $k\mathcal{Q}$. If I is generated by a finite number of paths in $\mathcal{Q}$, then I is called a monomial ideal and $\Lambda := k\mathcal{Q}/I$ a monomial algebra. For a finite-dimensional monomial algebra $\Lambda = k\mathcal{Q}/I$, using a certain set $AP(n)$ of overlaps constructed inductively by linking generators of I , Bardzell gave a minimal projective Λ^e -resolution $(P_\bullet, \phi_\bullet)$ of Λ in [B] (so called Bardzell's resolution). By using Bardzell's resolution, the Hochschild cohomology of monomial algebras are studied in the following papers [GS], [GSS], [FS], etc.


In this note, for a finite-dimensional monomial algebra Λ , we propose a method so that we easily calculate the Hochschild cohomology groups of Λ under some conditions. Let $\mathcal{Q}$ be a finite connected quiver and $\mathcal{Q}^{(i)}$ ($i = 1, 2$) a subquiver of $\mathcal{Q}$ such that $\mathcal{Q} = \mathcal{Q}^{(1)} \cup \mathcal{Q}^{(2)} = (\mathcal{Q}_0^{(1)} \cup \mathcal{Q}_0^{(2)}, \mathcal{Q}_1^{(1)} \cup \mathcal{Q}_1^{(2)})$. Let $I^{(1)} = \langle X \rangle$ (resp. $I^{(2)} = \langle Y \rangle$) be a monomial ideal of $k\mathcal{Q}^{(1)}$ (resp. $k\mathcal{Q}^{(2)}$) for X (resp. Y) a set of paths of $k\mathcal{Q}^{(1)}$ (resp. $k\mathcal{Q}^{(2)}$) and $I = \langle X, Y \rangle$ a monomial ideal of $k\mathcal{Q}$. We assume that I and $I^{(i)}$ ($i = 1, 2$) are admissible ideals. Then we define $\Lambda = k\mathcal{Q}/I$, $\Lambda_{(1)} = k\mathcal{Q}^{(1)}/I^{(1)}$ and $\Lambda_{(2)} = k\mathcal{Q}^{(2)}/I^{(2)}$. Hence Λ and $\Lambda_{(i)}$ are finite-dimensional monomial algebras for $i = 1, 2$. For the monomial algebra Λ , under a *separability condition* (i.e. $\mathcal{Q}_1^{(1)} \cap \mathcal{Q}_1^{(2)} = \emptyset$), we investigate the minimal projective Λ^e -module resolution of Λ given by Bardzell ([B]). Moreover, under an additional condition, we show that, for $n \geq 2$, the Hochschild cohomology group $\text{HH}^n(\Lambda)$ of Λ is isomorphic to the direct sum of the Hochschild cohomology groups $\text{HH}^n(\Lambda_{(1)})$ and $\text{HH}^n(\Lambda_{(2)})$.

Throughout this note, for all arrows a of $\mathcal{Q}$, we denote the origin of a by $o(a)$ and the terminus of a by $t(a)$. Also, for simplicity, we denote $\otimes_k$ by $\otimes$. For the general notation, we refer to [ASS].

2. THE SET $AP(n)$ OF OVERLAPS AND BARDZELL'S RESOLUTION

In this section, following [B] and [GS], we will summarize the definition of the set $AP(n)$ ($n \geq 0$) of overlaps.

Definition 2.1. A path $q \in k\mathcal{Q}$ overlaps a path $p \in k\mathcal{Q}$ with overlap pu if there exist u, v such that $pu = vq$ and $1 \leq l(u) \leq l(q)$, where $l(x)$ denotes the length of a path $x \in k\mathcal{Q}$. Note that we allow $l(x) = 0$ here.



A path q properly overlaps a path p with overlap pu if q overlaps p and $l(v) \geq 1$.

Let $\Lambda = k\mathcal{Q}/I$ be a finite-dimensional monomial algebra where $I = \langle \rho \rangle$ has a minimal set of generators ρ of paths of length at least 2.


Definition 2.2. For $n = 0, 1, 2$, we set

- $AP(0) := \mathcal{Q}_0$ (the set of all vertices of $\mathcal{Q}$);
- $AP(1) := \mathcal{Q}_1$ (the set of all arrows of $\mathcal{Q}$);
- $AP(2) := \rho$.

For $n \geq 3$, we define the set $AP(n)$ of all overlaps R^n formed in the following way: We say that $R^2 \in AP(2)$ maximally overlaps $R^{n-1} \in AP(n-1)$ with overlap $R^n = R^{n-1}u$ if

- (1) $R^{n-1} = R^{n-2}p$ for some path p and $R^{n-2} \in AP(n-2)$;
- (2) R^2 overlap p with overlap pu ;
- (3) there is no element of $AP(2)$ which overlaps p with overlap being a proper prefix of pu .


The construction of the paths in $AP(n)$ may be illustrated with the following picture of R^n :



Remark 2.1. ([B]) Note that for $n \geq 2$, $AP(n) = AP(n)^{\text{op}}$.

In short, overlaps are constructed by linking generators of an admissible monomial ideal I . A sequence of those generators of I is called the associated sequence of paths ([GHZ]).

Example 2.1. Let $\mathcal{Q}$ be a quiver



bound by $I = \langle a_1a_2a_3, a_2a_3a_1, a_3a_1a_2 \rangle$. We set the algebra $\Lambda = k\mathcal{Q}/I$. Then we set


- $AP(0) := \mathcal{Q}_0 = \{v_0, v_1, v_2\}$, $AP(1) := \mathcal{Q}_1 = \{a_1, a_2, a_3\}$,
- $AP(2) := \{a_1a_2a_3, a_2a_3a_1, a_3a_1a_2\}$.

For $n \geq 3$, considering all overlaps linking by generators of I inductively, we have the following:

- $AP(3) = \{a_1a_2a_3a_1, a_2a_3a_1a_2, a_3a_1a_2a_3\}$,
- $AP(4) = \{a_1a_2a_3a_1a_2, a_2a_3a_1a_2a_3, a_3a_1a_2a_3a_1\}, \dots$,
- $AP(n) = \{\underbrace{a_1a_2a_3a_1}_{n+1} \dots, \underbrace{a_2a_3a_1a_2}_{n+1} \dots, \underbrace{a_3a_1a_2a_3}_{n+1} \dots\}$.

For example, the associated sequence of paths corresponding to $a_1a_2a_3a_1, a_2a_3a_1a_2, a_3a_1a_2a_3 \in AP(4)$ are (a_1a_2, a_2a_3, a_3a_1) , (a_2a_3, a_3a_1, a_1a_2) , (a_3a_1, a_1a_2, a_2a_3) , respectively.

Example 2.2. Let $\mathcal{Q}$ be a quiver



bound by $I' = \langle a_1 a_2, a_2 a_3 \rangle$. We set the algebra $\Lambda' = k\mathcal{Q}/I'$.

- $AP(0) := \mathcal{Q}_0 = \{v_0, v_1, v_2\}$, $AP(1) := \mathcal{Q}_1 = \{a_1, a_2, a_3\}$,
- $AP(2) := \{a_1 a_2, a_2 a_3\}$.

Considering all overlaps linking by generators of I inductively,


- $AP(3) = \{a_1 a_2 a_3\}$,
- $AP(n) = \emptyset$ for all $n \geq 4$.

For a monomial algebra $\Lambda = k\mathcal{Q}/I$, by using the set $AP(n)$, Bardzell determined a minimal projective Λ^e -resolution $(P_\bullet, \phi_\bullet)$ of Λ in [B].


Definition 2.3. Let $(P_\bullet, \phi_\bullet)$ be the minimal projective Λ^e -resolution of Λ in [B]. Then, for $n \geq 0$, we set

$$P_n = \coprod_{R^n \in AP(n)} \Lambda o(R^n) \otimes t(R^n) \Lambda.$$

From [B], if $R^{2n+1} \in AP(2n+1)$, then there uniquely exist $R_j^{2n}, R_k^{2n} \in AP(2n)$ and some paths a_j, b_k such that $R^{2n+1} = R_j^{2n} a_j = b_k R_k^{2n}$.



For even degree elements $R^{2n} \in AP(2n)$, there exist $r \geq 1$, $R_l^{2n-1} \in AP(2n-1)$ and paths p_l, q_l for $l = 1, 2, \dots, r$ such that $R^{2n} = p_1 R_1^{2n-1} q_1 = \dots = p_r R_r^{2n-1} q_r$.



Remark 2.2. Note that $o(R_j^{2n}) \otimes a_j \in \Lambda o(R_j^{2n}) \otimes t(R_j^{2n}) \Lambda$ and $b_k \otimes t(R_k^{2n}) \in \Lambda o(R_k^{2n}) \otimes t(R_k^{2n}) \Lambda$. Also, note that $p_l \otimes q_l \in \Lambda o(R_l^{2n-1}) \otimes t(R_l^{2n-1}) \Lambda$.

Definition 2.4. The map $\phi_{2n+1} : P_{2n+1} \rightarrow P_{2n}$ is given as follows. If $R^{2n+1} = R_j^{2n} a_j = b_k R_k^{2n} \in AP(2n+1)$, then

$$o(R^{2n+1}) \otimes t(R^{2n+1}) \mapsto o(R_j^{2n}) \otimes a_j - b_k \otimes t(R_k^{2n}).$$

The map $\phi_{2n} : P_{2n} \rightarrow P_{2n-1}$ is given as follows. If $R^{2n} = p_1 R_1^{2n-1} q_1 = \dots = p_r R_r^{2n-1} q_r$, then

$$o(R^{2n}) \otimes t(R^{2n}) \mapsto \sum_{l=1}^r p_l \otimes q_l.$$

The following result is the main theorem in [B].

Bardzell's Theorem ([B, Theorem 4.1]) Let $\mathcal{Q}$ be a finite quiver, and suppose that $\Lambda = k\mathcal{Q}/I$ is a monomial algebra with an admissible ideal I . Then the sequence

$$\dots \rightarrow P_{n+1} \xrightarrow{\phi_{n+1}} P_n \xrightarrow{\phi_n} \dots \xrightarrow{\phi_2} P_1 \xrightarrow{\phi_1} P_0 \xrightarrow{\pi} \Lambda \rightarrow 0$$

is a minimal projective resolution of Λ as a right Λ^e -module, where π is the multiplication map.

3. THE DECOMPOSITION OF HOCHSCHILD COHOMOLOGY GROUPS

Before stating main theorem, we recall our setting.

- $\mathcal{Q} = \mathcal{Q}^{(1)} \cup \mathcal{Q}^{(2)}$,
- $I^{(1)} = \langle X \rangle$ be a monomial ideal generated by X a set of paths of $k\mathcal{Q}^{(1)}$,
- $I^{(2)} = \langle Y \rangle$ a monomial ideal generated by Y a set of paths of $k\mathcal{Q}^{(2)}$,
- $I = \langle X, Y \rangle$ a monomial ideal of $k\mathcal{Q}$,
- $\Lambda = k\mathcal{Q}/I$, $\Lambda_{(1)} = k\mathcal{Q}^{(1)}/I^{(1)}$, $\Lambda_{(2)} = k\mathcal{Q}^{(2)}/I^{(2)}$: finite-dimensional algebras,
- $AP(2) := X \cup Y$, $AP^{(1)}(2) := X$, $AP^{(2)}(2) := Y$.

Then, as in the definition of $AP(n)$ of overlaps, we define $AP^{(1)}(n)$, $AP^{(2)}(n)$. Moreover, we define projective Λ^e -modules as follows:

$$\begin{aligned} P_n^{(1)} &= \coprod_{R^n \in AP^{(1)}(n)} \Lambda o(R^n) \otimes t(R^n) \Lambda, \\ P_n^{(2)} &= \coprod_{R^n \in AP^{(2)}(n)} \Lambda o(R^n) \otimes t(R^n) \Lambda, \\ P_n &= \coprod_{R^n \in AP(n)} \Lambda o(R^n) \otimes t(R^n) \Lambda. \end{aligned}$$

To prove our main result, we need the following lemma. As mentioned in Introduction, we consider the *separability condition* $AP^{(1)}(1) \cap AP^{(2)}(1) = \emptyset$.

Lemma 3.1. ([IFS, Lemma 3.1]) *Let $i \in \{1, 2\}$. If we assume $AP^{(1)}(1) \cap AP^{(2)}(1) = \emptyset$, then we have the following:*

- (a) *For all $n \geq 1$, $AP(n) = AP^{(1)}(n) \cup AP^{(2)}(n)$.*
- (b) *For all $n \geq 1$, $AP^{(1)}(n) \cap AP^{(2)}(n) = \emptyset$.*
- (c) *Let $n \geq 1$ and $p^n \in AP(n)$. Then R^n is a path of $k\mathcal{Q}^{(i)}$ if and only if $R^n \in AP^{(i)}(n)$.*

By Bardzell's Theorem and Lemma 3.1, we have the following proposition.

Proposition 3.2. ([IFS, Proposition 3.2]) *If the condition $\mathcal{Q}_1^{(1)} \cap \mathcal{Q}_1^{(2)} = \emptyset$ holds, then, in the following minimal projective resolution of Λ :*

$$\cdots \rightarrow P_{n+1} \xrightarrow{\phi_{n+1}} P_n \xrightarrow{\phi_n} P_{n-1} \rightarrow \cdots \xrightarrow{\phi_3} P_2 \xrightarrow{\phi_2} P_1 \xrightarrow{\phi_1} P_0 \xrightarrow{\pi} \Lambda \rightarrow 0,$$

for any $n \geq 1$, P_n is isomorphic to $P_n^{(1)} \oplus P_n^{(2)}$ as right Λ^e -modules and $\phi_{n+1} = \phi_{n+1}^{(1)} \oplus \phi_{n+1}^{(2)}$, where $\phi_{n+1}^{(i)} : P_{n+1}^{(i)} \rightarrow P_n^{(i)}$ ($i = 1, 2$) is the restriction of ϕ_{n+1} .

Remark 3.1. For $i = 1, 2$, $b_k \in \Lambda_{(i)} o(R_k^{2n})$, $a_j \in t(R_j^{2n}) \Lambda_{(i)}$, $p_l \in \Lambda_{(i)} o(R_l^{2n+1})$ and $q_l \in t(R_l^{2n+1}) \Lambda_{(i)}$ actually hold. So, for $n \geq 1$, $\phi_{n+1}^{(i)}$ sends $\coprod_{R^{n+1} \in AP^{(i)}(n+1)} \Lambda_{(i)} o(R^{n+1}) \otimes t(R^{n+1}) \Lambda_{(i)}$ to $\coprod_{R^n \in AP^{(i)}(n)} \Lambda_{(i)} o(R^n) \otimes t(R^n) \Lambda_{(i)}$ (not just to $\coprod_{R^n \in AP(n)} \Lambda o(R^n) \otimes t(R^n) \Lambda$). Therefore, $(\coprod_{R^n \in AP^{(i)}(n)} \Lambda_{(i)} o(R^n) \otimes t(R^n) \Lambda_{(i)}; \phi_{n+1}^{(i)})_{n \geq 1}$ is exactly a part of degree $n \geq 1$ for the minimal projective resolution of $\Lambda_{(i)}$ ($i = 1, 2$).

The following theorem is our main result.

Theorem 3.3. ([IFS, Theorem 3.3]) *If the condition $\mathcal{Q}_1^{(1)} \cap \mathcal{Q}_1^{(2)} = \emptyset$ holds and, for each $i = 1, 2$, $o(R^n)\Lambda t(R^n) = o(R^n)\Lambda_{(i)}t(R^n)$ holds for any $n \geq 1$ and any $R^n \in AP^{(i)}(n)$, then we have the direct sum decomposition of Hochschild cohomology groups*

$$HH^n(\Lambda) \cong HH^n(\Lambda_{(1)}) \oplus HH^n(\Lambda_{(2)})$$

for any $n \geq 2$.

Proof. By Proposition 3.2, we obtain the following right Λ^e -projective resolution of Λ :

$$\cdots \rightarrow P_{n+1} \xrightarrow{\phi_{n+1}} P_n \xrightarrow{\phi_n} P_{n-1} \rightarrow \cdots \xrightarrow{\phi_3} P_2 \xrightarrow{\phi_2} P_1 \xrightarrow{\phi_1} P_0 \xrightarrow{\pi} \Lambda \rightarrow 0,$$

where for any $n \geq 1$, $P_n = P_n^{(1)} \oplus P_n^{(2)}$ and $\phi_{n+1} = \phi_{n+1}^{(1)} \oplus \phi_{n+1}^{(2)}$.

Applying $\text{Hom}_{\Lambda^e}(-, \Lambda)$ to this resolution, we have the following sequence:

$$0 \rightarrow \widehat{P_0} \xrightarrow{\widehat{\phi_1}} \widehat{P_1} \xrightarrow{\widehat{\phi_2}} \cdots \xrightarrow{\widehat{\phi_n}} \widehat{P_n} \xrightarrow{\widehat{\phi_{n+1}}} \widehat{P_{n+1}} \rightarrow \cdots,$$

where $\widehat{P_n} = \text{Hom}_{\Lambda^e}(P_n, \Lambda)$, $\widehat{\phi_n} = \text{Hom}_{\Lambda^e}(\phi_n, \Lambda)$. By the assumption, if $p^n \in AP^{(i)}(n)$, then p^n is a path of $kQ^{(i)}$ for each i ($i = 1, 2$). So we have, for any $n \geq 1$,

$$\begin{aligned} \widehat{P_n} &= \text{Hom}_{\Lambda^e}(P_n, \Lambda) \\ &= \text{Hom}_{\Lambda^e}(P_n^{(1)} \oplus P_n^{(2)}, \Lambda) \\ &= \text{Hom}_{\Lambda^e}((\coprod_{p^n \in AP^{(1)}(n)} \Lambda o(p^n) \otimes t(p^n)\Lambda) \oplus (\coprod_{p^n \in AP^{(2)}(n)} \Lambda o(p^n) \otimes t(p^n)\Lambda), \Lambda) \\ &= \text{Hom}_{\Lambda^e}((\coprod_{p^n \in AP^{(1)}(n)} \Lambda o(p^n) \otimes t(p^n)\Lambda), \Lambda) \\ &\quad \oplus \text{Hom}_{\Lambda^e}((\coprod_{p^n \in AP^{(2)}(n)} \Lambda o(p^n) \otimes t(p^n)\Lambda), \Lambda) \\ &= (\coprod_{p^n \in AP^{(1)}(n)} o(p^n)\Lambda t(p^n)) \oplus (\coprod_{p^n \in AP^{(2)}(n)} o(p^n)\Lambda t(p^n)) \\ &= (\coprod_{p^n \in AP^{(1)}(n)} o(p^n)\Lambda_{(1)}t(p^n)) \oplus (\coprod_{p^n \in AP^{(2)}(n)} o(p^n)\Lambda_{(2)}t(p^n)) \\ &= \text{Hom}_{\Lambda_{(1)}^e}((\coprod_{p^n \in AP^{(1)}(n)} \Lambda_{(1)}o(p^n) \otimes t(p^n)\Lambda_{(1)}), \Lambda_{(1)}) \\ &\quad \oplus \text{Hom}_{\Lambda_{(2)}^e}((\coprod_{p^n \in AP^{(2)}(n)} \Lambda_{(2)}o(p^n) \otimes t(p^n)\Lambda_{(2)}), \Lambda_{(2)}). \end{aligned}$$

Also, by Remark 3.1, we have, for any $n \geq 1$,

$$\begin{aligned} \widehat{\phi_{n+1}} &= \text{Hom}_{\Lambda^e}(\phi_{n+1}, \Lambda) = \text{Hom}_{\Lambda^e}(\phi_{n+1}^{(1)} \oplus \phi_{n+1}^{(2)}, \Lambda) \\ &= \text{Hom}_{\Lambda^e}(\phi_{n+1}^{(1)}, \Lambda) \oplus \text{Hom}_{\Lambda^e}(\phi_{n+1}^{(2)}, \Lambda) \\ &= \text{Hom}_{\Lambda_{(1)}^e}(\phi_{n+1}^{(1)}, \Lambda_{(1)}) \oplus \text{Hom}_{\Lambda_{(2)}^e}(\phi_{n+1}^{(2)}, \Lambda_{(2)}) = \widehat{\phi_{n+1}^{(1)}} \oplus \widehat{\phi_{n+1}^{(2)}}. \end{aligned}$$

Hence the complex giving the Hochschild cohomology groups $\mathrm{HH}^n(\Lambda)$ ($n \geq 2$)

$$\widehat{P_1} \xrightarrow{\widehat{\phi_2}} \dots \xrightarrow{\widehat{\phi_n}} \widehat{P_n} \xrightarrow{\widehat{\phi_{n+1}}} \widehat{P_{n+1}} \rightarrow \dots$$

is decomposed into the following direct sum of complexes:

$$\widehat{P_1^{(1)}} \oplus \widehat{P_1^{(2)}} \xrightarrow{\widehat{\phi_2^{(1)} \oplus \phi_2^{(2)}}} \dots \xrightarrow{\widehat{\phi_n^{(1)} \oplus \phi_n^{(2)}}} \widehat{P_n^{(1)}} \oplus \widehat{P_n^{(2)}} \xrightarrow{\widehat{\phi_{n+1}^{(1)} \oplus \phi_{n+1}^{(2)}}} \widehat{P_{n+1}^{(1)}} \oplus \widehat{P_{n+1}^{(2)}} \rightarrow \dots$$

Therefore, we have $\mathrm{HH}^n(\Lambda) \cong \mathrm{HH}^n(\Lambda_{(1)}) \oplus \mathrm{HH}^n(\Lambda_{(2)})$ for any $n \geq 2$. $\square$

Remark 3.2. For $n = 0, 1$, the above equation fails in general (see Example 4.3 for the case $n = 1$).

If $\mathcal{Q}_0^{(1)} \cap \mathcal{Q}_0^{(2)} = \{v_0\}$ and $v_0 \Lambda v_0 = k v_0$, then we have $\mathcal{Q}_1^{(1)} \cap \mathcal{Q}_1^{(2)} = \emptyset$. Also, by Lemma 3.1 and Theorem 3.3, we have the following corollary.

Corollary 3.4. ([IFS, Corollary 3.4]) *In the case $\mathcal{Q}_0^{(1)} \cap \mathcal{Q}_0^{(2)} = \{v_0\}$ and $v_0 \Lambda v_0 = k v_0$, we have the direct sum decomposition of the Hochschild cohomology groups*

$$\mathrm{HH}^n(\Lambda) \cong \mathrm{HH}^n(\Lambda_{(1)}) \oplus \mathrm{HH}^n(\Lambda_{(2)})$$


for any $n \geq 2$.

Remark 3.3. Hence, for a finite dimensional monomial algebra obtained by linking some quivers bound by monomial relations successively, we can also decompose the Hochschild cohomology groups as in Corollary 3.4.

4. EXAMPLES


In this section, we give examples of monomial algebras satisfying the condition $AP^{(1)}(1) \cap AP^{(2)}(1) = \emptyset$.

Example 4.1. Let $\mathcal{Q}$ be a quiver



bound by $I = \langle a_1 a_2, a_2 a_3, a_3 a_1, b_1 b_2, b_2 b_3, b_3 b_1 \rangle$. We set the algebra $\Lambda = k\mathcal{Q}/I$. Let $\mathcal{Q}^{(1)}$ be the subquiver of $\mathcal{Q}$ bound by $I^{(1)} = \langle a_1 a_2, a_2 a_3, a_3 a_1 \rangle$ and $\mathcal{Q}^{(2)}$ the subquiver of $\mathcal{Q}$ bound by $I^{(2)} = \langle b_1 b_2, b_2 b_3, b_3 b_1 \rangle$. We set $\Lambda_{(i)} = k\mathcal{Q}^{(i)}/I^{(i)}$ for $i = 1, 2$. Then $\mathcal{Q}_{(1)}^{(1)} \cap \mathcal{Q}_{(1)}^{(2)} = \emptyset$ holds and for each $i = 1, 2$, $o(p^n) \Lambda t(p^n) = o(p^n) \Lambda_{(i)} t(p^n)$ holds for any $n \geq 1$ and $p^n \in AP^{(i)}(n)$. Applying Corollary 3.4, we obtain the direct sum decomposition of the Hochschild cohomology groups $\mathrm{HH}^n(\Lambda) \cong \mathrm{HH}^n(\Lambda_{(1)}) \oplus \mathrm{HH}^n(\Lambda_{(2)})$ for any $n \geq 2$. Also, since $\Lambda_{(i)}$ ($i = 1, 2$) is a self-injective Nakayama algebra, we know the dimension of $\mathrm{HH}^n(\Lambda_{(i)})$ from [EH, Propositions 4.4, 5.3] for $i = 1, 2$, and so we have the dimension of $\mathrm{HH}^n(\Lambda)$ by the decomposition above.


Example 4.2. Let $\mathcal{Q}$ be a quiver



bound by


$$I = \langle a_1 a_2 \cdots a_m, a_2 a_3 \cdots a_{m+1}, \dots, a_n a_1 \cdots a_{-n+m+1}, \\ b_1 b_2 \cdots b_{m'}, b_2 b_3 \cdots b_{m'+1}, \dots, b_{n'} b_1 \cdots b_{-n'+m'+1} \rangle$$

for any integers $m, m' \geq 2$ with $m \leq n$ and $m' \leq n'$. We set the algebra $\Lambda = kQ/I$. Let $Q^{(1)}$ be the subquiver of Q bound by $I^{(1)} = \langle a_1 a_2 \cdots a_m, a_2 a_3 \cdots a_{m+1}, \dots, a_n a_1 \cdots a_{-n+m+1} \rangle$ and $Q^{(2)}$ be the subquiver of Q bound by $I^{(2)} = \langle b_1 b_2 \cdots b_{m'}, b_2 b_3 \cdots b_{m'+1}, \dots, b_{n'} b_1 \cdots b_{-n'+m'+1} \rangle$, where $Q_0^{(1)} \cap Q_0^{(1)} = \{v_0\}$ and $Q_1^{(1)} \cap Q_1^{(2)} = \emptyset$. We set $\Lambda_{(i)} = kQ^{(i)}/I^{(i)}$





for $i = 1, 2$. Then the condition of Corollary 3.4 is satisfied. Applying Corollary 3.4, we obtain the direct sum decomposition of the Hochschild cohomology groups $HH^n(\Lambda) \cong HH^n(\Lambda_{(1)}) \oplus HH^n(\Lambda_{(2)})$ for any $n \geq 2$.

Example 4.3. Let Q be a quiver



bound by $I = \langle a_1 a_2, a_2 a_3, a_3 a_4, a_4 a_1, b_1 b_2, b_2 b_3, b_3 b_4, b_4 b_1 \rangle$. We set the algebra $\Lambda = kQ/I$. Let $Q^{(1)}$ be the subquiver of Q bound by $I^{(1)} = \langle a_1 a_2, a_2 a_3, a_3 a_4, a_4 a_1 \rangle$ and $Q^{(2)}$ be the subquiver of Q bound by $I^{(2)} = \langle b_1 b_2, b_2 b_3, b_3 b_4, b_4 b_1 \rangle$, where $Q_0^{(1)} \cap Q_0^{(1)} = \{v_0, v_1\}$ and $Q_1^{(1)} \cap Q_1^{(2)} = \emptyset$.

We set $\Lambda_{(i)} = kQ^{(i)}/I^{(i)}$ for $i = 1, 2$. Then $AP^{(1)}(1) \cap AP^{(2)}(1) = \emptyset$ holds and for each $i = 1, 2$, $o(R^n)\Lambda t(R^n) = o(R^n)\Lambda_{(i)}t(R^n)$ holds for any $n \geq 1$ and any $R^n \in AP^{(i)}(n)$. Applying Theorem 3.3, we obtain the direct sum decomposition of the Hochschild cohomology groups $HH^n(\Lambda) \cong HH^n(\Lambda_{(1)}) \oplus HH^n(\Lambda_{(2)})$ for any $n \geq 2$.



On the other hand, by direct computations, we have $\dim_k \mathrm{HH}^1(\Lambda) = 3$ and $\dim_k \mathrm{HH}^1(\Lambda_{(i)}) = 1$ ($i = 1, 2$). Hence the above decomposition does not hold for $n = 1$.

REFERENCES

- [ASS] I. Assem, D. Simson and A. Skowronski, *Elements of the representation theory of associative algebras*, Vol. 1. Techniques of representation theory. London Mathematical Society Student Texts, 65. Cambridge University Press, Cambridge, 2006.
- [B] M. J. Bardzell, The alternating syzygy behavior of monomial algebras, *J. Algebra* **188** (1997), no. 1, 69–89.
- [BO] P. A. Bergh and S. Oppermann, Cohomology of twisted tensor products, *J. Algebra* **320** (2008), no. 8, 3327–3338.
- [EH] K. Erdmann and T. Holm, Twisted bimodules and Hochschild cohomology for self-injective algebras of class A_n , *Forum Math.* **11** (1999), no. 2, 177–201.
- [FS] T. Furuya and N. Snashall, Support varieties for modules over stacked monomial algebras, *Comm. Algebra* **39** (2011), no. 8, 2926–2942.
- [GHZ] E. L. Green, D. Happel and D. Zacharia, Projective resolutions over Artin algebras with zero relations, *Illinois J. Math.* **29** (1985), 180–190.
- [GS] E. L. Green and N. Snashall, The Hochschild cohomology ring modulo nilpotence of a stacked monomial algebra, *Colloq. Math.* **105** (2006), no. 2, 233–258.
- [GSS] E. L. Green, N. Snashall and Ø. Solberg, The Hochschild cohomology ring modulo nilpotence of a monomial algebra, *J. Algebra Appl.* **5** (2006), no. 2, 153–192.
- [H] D. Happel, *Hochschild Cohomology of Finite-Dimensional Algebras*, pp. 108–126, Springer Lecture Notes in Mathematics, Vol. 1404, Springer-Verlag, Berlin/New York, 1989.
- [IFS] A. Itaba, T. Furuya and K. Sanada, On the decomposition of the Hochschild cohomology group of a monomial algebra satisfying a separability condition, *Comm. Algebra* **43** (2015), no. 6, 2282–2292.
- [S] N. Snashall, Support varieties and the Hochschild cohomology ring modulo nilpotence, *Proceedings of the 41st Symposium on Ring Theory and Representation Theory*, 68–82, Ed. H. Fujita, Tsukuba, 2009.

AYAKO ITABA
 DEPARTMENT OF MATHEMATICS
 TOKYO UNIVERSITY OF SCIENCE
 KAGURAZAKA 1-3, SHINJUKU, TOKYO 162-8601, JAPAN
E-mail address: j1110701@ed.tus.ac.jp

CURRENT ADDRESS:
 DEPARTMENT OF MATHEMATICS
 FACULTY OF SCIENCE
 SHIZUOKA UNIVERSITY
 OHYA 336, SHIZUOKA 422-8529, JAPAN
E-mail address: itaba.ayako@shizuoka.ac.jp