

Microlocal *a priori* estimates and the Cauchy problem

Kunihiko Kajitani (Univ. of Tsukuba 梶谷 邦彦)
Seiichiro Wakabayashi (Univ. of Tsukuba 若林 誠一郎)

1. Microlocal *a priori* estimates.

Microlocal analysis has made a great contribution to the recent development of the theory of partial differential equations, and many authors have proved that microlocal analysis is a powerful and useful tool in studying partial differential equations. In particular, it leads us to studies on standard forms of pseudodifferential operators and clarifies essential parts to be analyzed. However, for general problems one only obtains results modulo C^∞ . To remove "modulo C^∞ " one must make an effort according to the problems. In the studies of the Cauchy problem one must distinguish the time variable from other variables, and only limited application of microlocal analysis was permitted. Our purpose is to show that microlocal analysis is fairly applicable to the studies of uniqueness and well-posedness of the Cauchy problem. In [7] we showed that microlocal *a priori* estimates (Carleman type estimates) give theorems on propagation of singularities. Here we shall show that similar microlocal estimates also give theorems on uniqueness and well-posedness of the Cauchy problem. Let $P(x, \xi) = \xi_1^m + \sum_{|\alpha| \leq m, \alpha_1 < m} a_\alpha(x) \xi^\alpha$, where $a_\alpha(x) \in C^\infty(\mathbb{R}^n)$.

Microlocal *a priori* estimate at $z^0 \in S^*\mathbb{R}^n (\simeq \mathbb{R}^n \times S^{n-1})$ ($0 < A \leq \infty$)

$P(x, \xi; \gamma) \in S_{1,0}^m$ satisfies $(E; z^0, \{t_\tau(x, \xi)\}_{\tau \in \mathcal{T}}, A) \iff^{def}$
 $\exists \psi_j(x, \xi) \in C^\infty(T^*\mathbb{R}^n \setminus 0)$ ($j = 1, 2$) (pos. homo. of deg. 0), $\exists l_j \in \mathbb{R}$ ($1 \leq j \leq 3$) s.t.
 $\psi_j(x, \xi) = 1$ in a conic nbd of z^0 ($j = 1, 2$) and $\forall \tau \in \mathcal{T}, \forall b \in \mathbb{R}, 1 \leq \exists K < A, \exists \gamma_0 \geq 1, \exists C > 0$ satisfying

$$\| \langle D \rangle_\gamma^{l_1} v \| \leq C \{ \| \langle D \rangle_\gamma^{l_2} P_\Lambda(x, D; \gamma) v \| + \| \langle D \rangle_\gamma^{l_3} (1 - \psi_{1,h}(x, D)) v \| \}$$

if $v \in H^\infty(\mathbb{R}^n)$, $\gamma \geq \gamma_0$, $h = K\gamma$ and $\Lambda(x, \xi) = (t_\tau(x, \xi) - b) \log \langle \xi \rangle (1 - \Theta(4|\xi|/h)) \psi_2(x, \xi)$, where $\Theta(t) \in C_0^\infty(\mathbb{R})$, $\Theta(t) = 1$ ($|t| \leq 1$), $\text{supp } \Theta \subset (-2, 2)$, $\psi_{1,h}(x, \xi) = (1 - \Theta(|\xi|/h)) \psi_1(x, \xi)$, $P_\Lambda(x, D; \gamma) v = (e^{-\Lambda})(x, D) P(x, D; \gamma) (e^\Lambda)(x, D) v$, $\langle \xi \rangle_\gamma = (\gamma^2 + |\xi|^2)^{1/2}$ and $\| \cdot \|$ denotes the L^2 -norm.

(P-1) $p(x, \xi)$ (=the principal part of $P(x, \xi)$) is hyperbolic w.r.t. $\vartheta = (1, 0, \dots, 0) \in \mathbb{R}^n$

Put $P(x, D; \gamma) u = e^{-\gamma \zeta(x)} P(x, D) (e^{\gamma \zeta(x)} u)$ for $\zeta(x) \in \mathcal{B}^\infty(\mathbb{R}^n)$.

(P-2) $\forall z^0 = (x^0, \xi^0) \in S^*\mathbb{R}^n$ with $x_1^0 > 0$ and $dp(z^0) = 0$, $\forall \vartheta^0 \in \Gamma(p(x^0, \cdot), \vartheta)$, $\forall k > 0$, $\exists a_0 > 0$ s.t. ' $P(x, \xi; \gamma)$ for some $\zeta(x)$ which is equal to $(x - x^0) \cdot \vartheta^0 + k \times |x - x^0|^2$ near x^0 (resp. ' $P(x, \xi + i\gamma\vartheta)$) satisfies $(E; z^0, \{at_+(x, \xi; z^0)\}_{a \geq a_0}, \infty)$ (resp. $(E; z^0, \{at_-(x, \xi; z^0)\}_{a \geq a_0}, \infty)$,' where $t_\pm(x, \xi; z^0) = \pm(x_1 - x_1^0) + |x - x^0|^2$

$+ |\xi|/|\xi| - \xi^0|^2$ in a conic nbd of z^0 , $\Gamma(p(x^0, \cdot), \vartheta)$ denotes the component of $\{\xi \in \mathbb{R}^n; p(x^0, \xi) \neq 0\}$ containing ϑ , and ${}^tP(x, D)$ denotes the transposed of $P(x, D)$.

Assume that for each $z^{0'} = (x^{0'}, \xi^{0'}) \in S^*\mathbb{R}^{n-1}$ $P(x, D)$ can be written as $P(x, D) = P^1(x, D; z^{0'}) \cdots P^\nu(x, D; z^{0'}) + R(x, D; z^{0'})$ in $\mathcal{C}'(z^{0'})$, where $\mathcal{C}'(z^{0'})$ is a conic nbd of $(0, z^{0'})$ in $\mathbb{R} \times (T^*\mathbb{R}^{n-1} \setminus 0)$, $x' = (x_2, \dots, x_n)$, $\nu \equiv \nu(z^{0'})$, $P^k(x, \xi, z^{0'}) \in S_{1,0}^{m_k,0}$ (the coef. of $\xi_1^{m_k} = 1$), $R(x, \xi; z^{0'}) \in \bigcap_l S_{1,0}^{m,l}$, $a(x, \xi) \in S_{1,0}^{m,l} \iff^{def} a(x, \xi) = \sum_{j=0}^{[m]} a_j(x, \xi') \xi_1^j$, $a_j(x, \xi') \in S_{1,0}^{m+l-j}$ (classical). Put $t'_\pm(x, \xi'; z^{0'}) = \pm x_1 + x_1^2 + |x' - x^{0'}|^2 + |\xi'/|\xi'| \pm \xi^{0'}|^2$ in a conic nbd of $(0, z^{0'})$.

(P-3) $\forall z^{0'} = (x^{0'}, \xi^{0'}) \in S^*\mathbb{R}^{n-1}$, $1 \leq \forall k \leq \nu(z^{0'})$, $\exists \mathcal{P}^k(x, \xi; z^{0'}) \in S_{1,0}^{m_k,0}$ (hyp. w.r.t. ϑ , the coef. of $\xi_1^{m_k} = 1$) s.t. (1) $\mathcal{P}^k(x, \xi; z^{0'}) = P^k(x, \xi; z^{0'})$ ($(x, \xi') \in \mathcal{C}'(z^{0'})$, $|\xi'| \geq 1$). (2) $\mathcal{P}^k(x, \xi; z^{0'}) = \mathcal{P}^k(\xi; z^{0'})$ ($x \notin \Omega(z^{0'})$ ($\subset \subset \mathbb{R}^n$): a nbd of $(0, x^{0'})$), where $\mathcal{P}^k(\xi; z^{0'})$ is pos. homo. for $|\xi'| \geq 1$. (3) $\forall z^1 = (x^1, \xi^1) \in S^*\mathbb{R}^n$ with $x^1 \in \overline{\Omega(z^{0'})}$ and $d\mathcal{P}_0^k(z^1; z^{0'}) = 0$, $\exists a_0, a'_0 > 0$ s.t. ' $\mathcal{P}^k(x, \xi - i\gamma\vartheta; z^{0'})$ ($(+)\text{sign}$) ($resp.$ ${}^t\mathcal{P}^k(x, \xi + i\gamma\vartheta; z^{0'})$ ($(-)\text{sign}$)) satisfies $(E; z^1, \{at_\pm(x, \xi; z^1) + a't'_\pm(x, \pm\xi'; z^{0'}) \log\langle \xi' \rangle \times 1/\log\langle \xi \rangle\}_{a \geq a_0, a' \geq a'_0, \infty})$, where $\mathcal{P}_0^k(x, \xi; z^{0'})$ denotes the principal symbol of $\mathcal{P}^k(x, \xi; z^{0'})$.

For $x^0 \in \mathbb{R}^n$ define $K_{x^0}^\pm = \{x(t); \pm t \geq 0\}$, and $\{x(t)\}$ is a Lipschitz cont. curve in $\mathbb{R}^n$ satisfying $(d/dt)x(t) \in \Gamma(p(x(t), \cdot), \vartheta)^*$ (a.e. t) and $x(0) = x^0$, where $\Gamma^* = \{y \in \mathbb{R}^n; y \cdot \eta \geq 0 \text{ for } \forall \eta \in \Gamma\}$.

(P-4) $K_{x^0}^- \cap \{x_1 \geq 0\}$ is bdd for every $x^0 \in \mathbb{R}^n$.

Theorem 1. Assume that (P-1)–(P-4) are satisfied. Then $\forall f \in \mathcal{D}'$ (resp. C^∞) with $\text{supp } f \subset \{x_1 \geq 0\}$, $\exists! u \in \mathcal{D}'$ (resp. C^∞) s.t. $P(x, D)u = f$, $\text{supp } u \subset \{x_1 \geq 0\}$. Moreover, $\text{supp } u \subset \{x \in \mathbb{R}^n; x \in K_y^+ \text{ for some } y \in \text{supp } f\}$.

To prove Theorem 1 we first derive local *a priori* estimates (Carleman type estimates) from microlocal *a priori* estimates. This proves uniqueness of the Cauchy problem. This argument can be also applicable to operators which are not necessarily hyperbolic. Next we derive hyperbolic estimates for ${}^tP(x, D)$ from (P-3), which proves existence of solutions to the Cauchy problem. In deriving hyperbolic estimates the condition (2) of (P-3) plays an important role. For the detail we refer to [?].

2. The Cauchy problem.

We shall assume that $P(x, D)$ satisfies at least one of the conditions (A) $_{x^0}$, (B) $_{x^0}$ and (C) $_{x^0}$ for $z^0 \in S^*\mathbb{R}^n$ with $dp(z^0) = 0$, which will be defined below. Let $z^0 = (x^0, \xi^0) \in S^*\mathbb{R}^n$ satisfy $dp(z^0) = 0$. We say that $P(x, D)$ satisfies the condition (A) $_{x^0}$ if (A-1) $_{x^0}$ and (A-2) $_{x^0}$ below are fulfilled.

(A-1) $_{x^0}$ $\exists$ conic nbd $\mathcal{C}$ of z^0 , $\exists$ conic nbd $\tilde{\mathcal{C}}$ of (y^0, η^0) , $\exists$ homo. canonical transf. $\chi: \tilde{\mathcal{C}} \xrightarrow{\sim} \mathcal{C}$, $\exists$ symbols $q(\eta)$, $e(y, \eta)$ s.t. $z^0 = \chi(y^0, \eta^0)$, $q(\eta)$: pos. homo. of deg. $m' \in \mathbb{N}$ and $p(\chi(y, \eta)) = e(y, \eta)q(\eta)$, $e(y, \eta) \neq 0$ for $(y, \eta) \in \tilde{\mathcal{C}}$.

Let F_1 and F_2 be classical Fourier integral operators corresponding to χ and χ^{-1} which are elliptic at (y^0, η^0) and z^0 , respectively. Under (A-1) $_{x^0}$ we can write

$$\sigma(F_2 P(x, D) F_1)(y, \eta) = \tilde{e}(y, \eta)(q(\eta) + s(y, \eta))$$

in a conic nbd $\tilde{C}_0$ of (y^0, η^0) if $|\eta| \geq 1$, where $\tilde{e}(y, \eta)$ is a classical symbol, which is elliptic in $\tilde{C}_0$, and $s(y, \eta) \in S_{1,0}^{m'-1}$ is a classical symbol.

(A-2) $_{z^0}$ $\exists C_0 > 0, \exists C_\alpha > 0$ for $\forall \alpha$ s.t.

$$|s_{(\alpha)}(y, \eta)| \leq C_\alpha |q(\eta; \tilde{\vartheta})| \quad \text{for } (y, \eta) \in \tilde{C}_0 \text{ with } |\eta| \geq C_0,$$

where $\tilde{\vartheta} = d\eta_{z^0}(0, \vartheta)$, $\chi^{-1}(x, \xi) = (y(x, \xi), \eta(x, \xi))$ and $q(\eta; \zeta) = \sum_{|\alpha| \leq m} (-i\zeta)^\alpha q^{(\alpha)}(\eta) \times \alpha!^{-1}$ for $\zeta \in \mathbb{R}^n$.

We note that $q(\eta)$ is microhyperbolic with respect to $\tilde{\vartheta}$ at (y^0, η^0) (see [16]). By the definition of microhyperbolicity we may assume that $|q(\eta; s\tilde{\vartheta})| \geq cs^m$ for $(y, \eta) \in \tilde{C}_0$ and $0 < s \leq s_0$, where $c > 0$ and $s_0 > 0$. This implies that $\tilde{\vartheta} \neq 0$ (see, also, Lemma 3.2 in [15]). We say that $P(x, D)$ satisfies the condition (B) $_{z^0}$ if (B-1) $_{z^0}$ –(B-3) $_{z^0}$ below are fulfilled.

(B-1) $_{z^0}$ $\exists$ conic nbd $\mathcal{C}$ of z^0 , $\exists$ conic nbd $\tilde{\mathcal{C}}$ of (y^0, η^0) , $\exists$ homo. canonical transf. $\chi : \tilde{\mathcal{C}} \xrightarrow{\sim} \mathcal{C}$, $\exists$ real-valued funs $t_j(y) \in B^\infty(\mathbb{R}^n)$ ($1 \leq j \leq N$), $\exists$ real-valued symbols $\lambda(y, \eta')$, $\alpha(y, \eta')$, $e(y, \eta)$ and $\exists C > 0$ s.t. $z^0 = \chi(y^0, \eta^0)$, $\eta^0 = (0, \dots, 0, 1)$, $d\chi_{(y^0, \eta^0)}(0, \vartheta), d\chi_{(y^0, \eta^0)}(0, \nabla t_j(y^0)) \in \Gamma(p_{z^0}, (0, \vartheta))$ ($1 \leq j \leq N$), $\lambda(y, \eta')$, $\alpha(y, \eta')$: pos. homo. of deg. 1, 2 (resp.), $p(\chi(y, \eta)) = e(y, \eta)(\eta_1(\eta_1 - \lambda(y, \eta')) - \alpha(y, \eta'))$ and $e(y, \eta) \neq 0$ for $(y, \eta) \in \tilde{\mathcal{C}}$, $\lambda(y^0, \eta^{0'}) = \alpha(y^0, \eta^{0'}) = 0$, and $\alpha(y, \eta') \geq 0$ and $T(y)\partial\alpha/\partial y_1(y, \eta') \leq C\alpha(y, \eta')$ for $(y, \eta') \in \tilde{\mathcal{C}}'$, where $p_{z^0}(\delta z)$ ($\neq 0$ in $\delta z \in \mathbb{R}^{2n}$) is defined as $p(z^0 + s\delta z) = s^\mu(p_{z^0}(\delta z) + o(1))$ when $s \rightarrow 0$, $\Gamma(p_{z^0}, (0, \vartheta))$ is the connected component of $\{\delta z \in \mathbb{R}^{2n}; p_{z^0}(\delta z) \neq 0\}$ containing $(0, \vartheta)$ ($\in \mathbb{R}^{2n}$), $T(y) = \min_{1 \leq j \leq N} |t_j(y)|$ and $\tilde{\mathcal{C}}' = \{(y, \eta'); (y, \eta) \in \tilde{\mathcal{C}} \text{ for some } \eta_1\}$.

Let F_1 and F_2 be classical Fourier integral operators corresponding to χ and χ^{-1} which are elliptic at (y^0, η^0) and z^0 , respectively. Under (B-1) $_{z^0}$ we can write

$$\sigma(F_2 P(x, D) F_1)(y, \eta) = \tilde{e}(y, \eta) \{ \eta_1(\eta_1 - \lambda(y, \eta')) - \alpha(y, \eta') + \beta(y, \eta) \}$$

in a conic nbd $\tilde{C}_0$ of (y^0, η^0) if $|\eta| \geq 1$, where $\tilde{e}(y, \eta)$ is a classical symbol, which is elliptic in $\tilde{C}_0$, and $\beta(y, \eta) \in S_{1,0}^1$ is a classical symbol.

(B-2) $_{z^0}$ $\exists$ classical symbol $A(y, \eta') \in S_{1,0}^0$, $\exists C > 0$ s.t.

$$T(y) |\Im \beta(y, 0, \eta') - \sum_{j=2}^n (\partial^2 \alpha / \partial y_j \partial \eta_j)(y, \eta') / 2 - A(y, \eta') \lambda(y, \eta')| \leq C(\alpha(y, \eta')^{1/2} + 1) \quad \text{for } (y, \eta') \in \tilde{C}_0'.$$

We say that the condition (B-3) $_{z^0}$ is satisfied if at least one of the following conditions (B-3-1) $_{z^0}$ and (B-3-2) $_{z^0}$ is satisfied:

(B-3-1) $_{z^0}$ $\exists$ classical symbol $B(y, \eta') \in S_{1,0}^0$, $\exists C > 0$ s.t.

$$T(y) |\Re \beta(y, 0, \eta') - B(y, \eta') \lambda(y, \eta')| \leq C(\alpha(y, \eta')^{1/2} + 1)$$

for $(y, \eta') \in \tilde{\mathcal{C}}'_0$.

(B-3-2) $_{z^0}$ $\Re P_{m-1}(z^0) < \text{Tr}^+ F_p$,
where F_p denotes the Hamilton map (fundamental matrix) corresponding to the Hessian of $p(z)/2$ at $z = z^0$ and $\text{Tr}^+ F_p = \sum \lambda_j$, the $i\lambda_j$ are the eigenvalues of F_p with pos. imaginary parts.

We say that $P(x, D)$ satisfies the condition (C) $_{z^0}$ if the following conditions (C-1) $_{z^0}$ and (C-2) $_{z^0}$ are fulfilled:

(C-1) $_{z^0}$ $\exists$ conic nbd $\mathcal{C}$ of z^0 , $\exists$ conic nbd $\tilde{\mathcal{C}}$ of (y^0, η^0) , $\exists$ homo. canonical transf. $\chi: \tilde{\mathcal{C}} \xrightarrow{\sim} \mathcal{C}$, $\exists$ real-valued funs $t_j(y) \in \mathcal{B}^\infty(\mathbb{R}^n)$ ($1 \leq j \leq N$), $\exists c > 0$ s.t. $z^0 = \chi(y^0, \eta^0)$, $d\chi_{(y^0, \eta^0)}(0, \nabla t_j(y^0)) \in \Gamma(p_{z^0}, (0, \vartheta))$ ($1 \leq j \leq N$),

$$h_{m-1}(x, \xi) \geq ch_{m-2}(x, \xi)T(x, \xi)^2$$

for $(x, \xi) \in \mathcal{C}$ with $|\xi| = 1$, where $h_k(x, \xi)$ ($0 \leq k \leq m$) are defined as $|p(x, \xi - is\vartheta)|^2 = \sum_{j=0}^m s^{2j} h_{m-j}(x, \xi)$ for $(x, \xi) \in T^*\mathbb{R}^n$ and $s \in \mathbb{R}$, $T(x, \xi) = \min_{1 \leq j \leq N} |t_j(y(x, \xi))|$ and $\chi^{-1}(x, \xi) = (y(x, \xi), \eta(x, \xi))$.

(C-2) $_{z^0}$ $\exists C > 0$ s.t.

$$|P_j(x, \xi)| \leq Ch_{2j-m}(x, \xi)^{1/2}$$

for $(x, \xi) \in \mathcal{C}$ with $|\xi| = 1$ and $[m/2] + 1 \leq j \leq m-1$, where $P_j(x, \xi) = \sum_{|\alpha|=j} a_\alpha(x) \xi^\alpha$ ($0 \leq j \leq m-1$).

Theorem 2. Assume that (P-1) and (P-4) are satisfied and that at least one of the conditions (A) $_{z^0}$, (B) $_{z^0}$ and (C) $_{z^0}$ is satisfied if $z^0 = (x^0, \xi^0) \in S^*\mathbb{R}^n$, $x_1^0 \geq 0$ and $dp(z^0) = 0$. Then $\forall f \in \mathcal{D}'$ (resp. C^∞) with $\text{supp } f \subset \{x_1 \geq 0\}$, $\exists! u \in \mathcal{D}'$ (resp. C^∞) s.t. $P(x, D)u = f$, $\text{supp } u \subset \{x_1 \geq 0\}$. Moreover, $\text{supp } u \subset \{x \in \mathbb{R}^n; x \in K_y^+ \text{ for some } y \in \text{supp } f\}$.

Theorem 2 can be proved, applying Theorem 1 (or an improved version) (see Theorem 5.2 in [8]). If one finds other conditions to give microlocal *a priori* estimates at $z^0 \in S^*\mathbb{R}^n$, one can add these conditions to (A) $_{z^0}$, (B) $_{z^0}$ and (C) $_{z^0}$ and improve Theorem 2. The condition (A) $_{z^0}$ is satisfied if the characteristic roots of $p(x, \xi) = 0$ are involutive and the lower order terms of $P(x, D)$ satisfy the so-called Levi conditions, which was studied by Zeman [17] (see, also, [1], [10]). The condition (B) $_{z^0}$ implies that $p(x, \xi)$ has double characteristics at z^0 . If $P(x, D)$ is effectively hyperbolic, then $P(x, D)$ satisfies both the conditions (B) $_{z^0}$ and (C) $_{z^0}$. C^∞ well-posedness of the Cauchy problem for effectively hyperbolic operators was proved by Iwasaki [6] (see, also, [3], [5], [11], [12], [14]). The operators, which was treated by Ivrii [4], satisfy the condition (B) $_{z^0}$ with $T(y) \equiv 1$ (see, also, [2]). We can treat both effectively hyperbolic cases and non effectively hyperbolic cases at the same time by introducing $T(y)$ in (B) $_{z^0}$. The condition (C) $_{z^0}$ is closely related to the conditions given in [13]. Some results on propagation of singularities were essentially obtained in [15] if the operators are microhyperbolic at z^0 and satisfy (A) $_{z^0}$. For microhyperbolic operators satisfying (B) $_{z^0}$ we obtained theorems on propagation of singularities in [13]. Applying the methods

developed in [7], we can also obtain theorems on propagation of singularities for microhyperbolic operators which satisfy at least one of the conditions $(A)_{x^0}$, $(B)_{x^0}$ and $(C)_{x^0}$. For the proof of Theorem 2 and the detail we refer to [9].

References

1. J. Chazarain, Opérateurs hyperboliques à caractéristiques de multiplicité constant, Ann. Inst. Fourier, 24 (1974), 173–202.
2. L. Hörmander, The Cauchy problem for differential equations with double characteristics, J. Analyse Math., 32 (1977), 118–196.
3. V. Ja. Ivrii, Sufficient conditions for regular and completely regular hyperbolicity, Trudy Moskov. Mat. Obšč., 33 (1976), 3–66.
4. V. Ja. Ivrii, The well-posedness of the Cauchy problem for nonstrictly hyperbolic operators. III. The energy integral, Trudy Moskov. Mat. Obšč., 34 (1977), 151–170.
5. N. Iwasaki, The Cauchy problem for effectively hyperbolic equations (a standard type), Publ. RIMS, Kyoto Univ., 20 (1984), 551–592.
6. N. Iwasaki, The Cauchy problem for effectively hyperbolic equations (general case), J. Math. Kyoto Univ., 25 (1985), 727–743.
7. K. Kajitani and S. Wakabayashi, Propagation of singularities for several classes of pseudodifferential operators, to appear.
8. K. Kajitani and S. Wakabayashi, Microlocal *a priori* estimates and the Cauchy problem I, preprint.
9. K. Kajitani and S. Wakabayashi, Microlocal *a priori* estimates and the Cauchy problem II, preprint.
10. S. Mizohata and Y. Ohya, Sur la condition de E. E. Levi concernant des équations hyperboliques, Publ. RIMS, Kyoto Univ., 4 (1968), 511–526.
11. T. Nishitani, Local energy integrals for effectively hyperbolic operators, I, J. Math. Kyoto Univ., 24 (1984), 623–658.
12. T. Nishitani, Local energy integrals for effectively hyperbolic operators, II, J. Math. Kyoto Univ., 24 (1984), 659–666.
13. T. Nishitani, Hyperbolic operators with symplectic multiple characteristics, to appear.
14. O. A. Oleinik, On the Cauchy problem for weakly hyperbolic equations, Comm. Pure Appl. Math., 23 (1970), 569–586.
15. S. Wakabayashi, Singularities of solutions of the Cauchy problems for operators with nearly constant coefficient hyperbolic principal part, Comm. in Partial Differential Equations, 8 (1983), 347–406.
16. S. Wakabayashi, Generalized Hamilton flows and singularities of solutions of the hyperbolic Cauchy problem, Taniguchi Symp., HERT, Kyoto 1984, Kinokuniya, Tokyo, pp415–423.
17. M. Zeman, The well-posedness of the Cauchy problem for partial differential equations with multiple characteristics, Comm. in Partial Differential Equations, 2 (1977), 223–249.