

Self-organization of droplets and buckling of droplet chains in free-standing liquid crystalline films

Otto-von-Guericke Universität Magdeburg, Institut für Experimentelle
Physik, Magdeburg, Germany Camilla Völtz ¹

While nematic liquid crystals are good models for anisotropic bulk emulsions [1], free-standing films represent a convenient model for two-dimensional systems. Here free-standing smectic films are investigated at the transition from the smectic C phase to the isotropic phase. In the vicinity of the bulk transition temperature isotropic droplets of micrometer size appear in the film. A characteristic feature of the droplets is their mutual interaction by elastic distortions of the c -director [2, 3]. The director deformations created by isotropic droplets of different sizes are discussed. Depending upon droplet size and anchoring conditions, topological defects can be induced in the c -director field.

The isotropic droplets can interact and self-organize in different patterns, like chains, lattices and rings. If the c -director field in a free-standing film is prepared as a target pattern with a continuous radial deformation, the droplets align tangentially in regular chains in the structure of the c -director field. Incorporation of additional droplets lengthens the chains at given ring diameter until they form complete rings. Further chain growth introduces a reversible buckling-instability (see Fig. 1) with a characteristic wavelength [4]. The phenomenon is similar in many respects to growth processes in biological systems or Euler buckling in polymer foils.

References

- [1] P. Poulin, H. Stark, T.C. Lubensky, and D.A. Weitz, *Science* **275** (1997).

¹ Current address: Liquid Crystal Nano-System Project, ERATO/SORST, Japan Science and Technology Agency (JST), 5-9-9 Tokodai, Tsukuba, Ibaraki 300-2635, Japan




Figure 1: Buckling of isotropic droplet chains in a target pattern of the c -director field. The width of the image is $655.2\ \mu\text{m}$. Individual droplets in the chains are only resolved in the central part of the image; otherwise the chains are recognized by dark lines. The innermost rings are replete and start to develop a modulation pattern with characteristic wave length.

- [2] P. Cluzeau, V. Bonnand, G. Joly, V. Dolganov, and H. T. Nguyen, Eur. Phys. J **10** (2003), 231.
- [3] C. Völtz and R. Stannarius, Phys. Rev. E **70** (2004), 061702.
- [4] C. Völtz and R. Stannarius, Phys. Rev. E (2005), in press.