

On the isomorphisms between the centers of the principal p -block algebras induced by the Glauberman-Dade correspondence

Fuminori Tasaka (田阪文規)

Division of Mathematical Science and Physics, Chiba Univ. (千葉大学自然科学研究科)

1

For a prime p , let $(\mathcal{K}, \mathcal{O}, k)$ be a p -modular system where $\mathcal{O}$ is a complete discrete valuation ring having the residue field k of characteristic p which is algebraically closed and having the quotient field $\mathcal{K}$ of characteristic zero which will be assumed to be large enough for any finite group we consider in this article. Below, by a character, we mean a $\mathcal{K}$ -character.

Glauberman showed in [5] that there is a bijective correspondence between the set $\text{Irr}(G)^S$ of all S -invariant irreducible characters of G and the set $\text{Irr}(C_G(S))$ of all irreducible characters of $C_G(S)$, called the *Glauberman correspondence* of characters, where G is a finite group and S is a finite solvable group such that S acts on G via automorphism and $(|G|, |S|) = 1$, where $|G|$ and $|S|$ denote the orders of G and S , respectively. When S is cyclic, a basic relation between $\chi \in \text{Irr}(G)^S$ and the Glauberman corresponding character $\beta_\chi \in \text{Irr}(C_G(S))$ is

$$\hat{\chi}(cs) = \epsilon_\chi \beta_\chi(c), \quad (\#)$$

where $\hat{\chi}$ is a unique extension of χ to the semi-direct product $G \rtimes S$ satisfying $S \subseteq \text{Ker}(\det(\hat{\chi}))$, called the *canonical extension*, c is any element of $C_G(S)$, s is any generator of S and ϵ_χ is a uniquely determined sign, see [5, Theorem 3].

Dade gives in [4] a new approach to the Glauberman correspondence without considering the canonical extension and the relation $(\#)$ above, and partly generalizes it: he gives in [4, Theorem 6.8] a bijective correspondence between $\text{Irr}(G)^E$ and $\text{Irr}(G')^{E'}$, where E is a finite group, G is a normal subgroup of E such that E/G is cyclic, E' is a subgroup of E such that $E = GE'$, and $G' = G \cap E'$ (hence, G' is normal in E' and E'/G' is isomorphic to the cyclic group E/G) with the condition that E'_0 , the subset of E' consisting of every element of E' whose canonical image in E'/G' is a generator of E'/G' , is a trivial intersection subset of E with E' as its normalizer, that is, $E'_0 \cap E_0'^t$ is the empty set for any $t \in E - E'$. In fact, the correspondence of characters is given for twisted group algebras of G and G' over $\mathcal{K}$. It is shown that, with the notations above, when S is cyclic, the above conditions are satisfied by taking $G \rtimes S$, G , $C_G(S) \times S$ and $C_G(S)$ as E , G , E' and G' , respectively ([4, Lemma 7.5]), and,

in this case, Dade's correspondence coincides with the Glauberman correspondence ([4, Proposition 7.8]). We call the above correspondence of characters the *Glauberman-Dade correspondence*.

Watanabe began in [11] a p -block-theoretical study of the Glauberman correspondence and she showed that, when b is an S -invariant p -block of G with an S -centralized defect group, the Glauberman correspondence induces a perfect isometry (for this notion, see [2]) between $\mathbb{Z}\text{Irr}(b)$ and $\mathbb{Z}\text{Irr}(w(b))$ for some uniquely determined p -block $w(b)$ of G^S .

In general, by the theorem of Broué ([2]), perfectly isometric p -blocks have isomorphic centers, and so $Z(\mathcal{O}Gb) \simeq Z(\mathcal{O}G^S w(b))$ as $\mathcal{O}$ -algebras.

On the other hand, in [8], Okuyama gives an explicit isomorphism between the centers of $\mathcal{O}Gb$ and $\mathcal{O}G^S w(b)$ using the relation (#).

In this article, we note that, when $|E/G|$ is a prime, under the condition analogous to the one of Watanabe (see, Condition 3.1), the Glauberman-Dade correspondence induces a bijective correspondence between the characters belonging to the principal p -blocks of G and G' (Proposition 3.3), and give an explicit isomorphism between the centers of the principal p -block algebras of G and G' (Theorem 3.5) which coincides with the one given by Okuyama when the hypotheses are the same. For a more general situation, see [10], and for standard facts, see [7].

2

Below, we always assume the following:

Condition 2.1. E is a finite group with a normal subgroup G such that the quotient group $F = E/G$ is cyclic of prime order q . E' is a subgroup of E such that $E = GE'$. G' is a normal subgroup of E' defined by $G' = G \cap E'$. Set $E_0 = E - G$ and $E'_0 = E' - G'$. E'_0 is a trivial intersection subset of E with E' as its normalizer, that is, $E'_0 \cap E'^\tau = \emptyset$, the empty set, for any $\tau \in E - E'$.

Take $s \in E'_0$. Then $E = \langle G, s \rangle$ and $E' = \langle G', s \rangle$.

Denote by π the canonical epimorphism from E to E/G , and set $F = \pi(G) = \pi_{E'}(E') \simeq E'/G'$, where $\pi_{E'}$ is the restriction of π to E' .

Choose once and for all an isomorphism $F \simeq \text{Hom}(F, \mathcal{K}^\times)$ of groups. Denote by $\hat{F}$ the cyclic group $\text{Hom}(F, \mathcal{K}^\times)$, and let λ be a generator of $\hat{F}$. Then $\hat{F}$ acts on $\text{Irr}(E)$ by $(\lambda\theta)(x) = \lambda(\pi(x))\theta(x)$ for $\theta \in \text{Irr}(E)$ and $x \in E$, and on $\text{Irr}(E')$ by $(\lambda\theta')(x') = \lambda(\pi_{E'}(x'))\theta'(x')$ for $\theta' \in \text{Irr}(E')$ and $x' \in E'$. Note that a primitive q -th root of unity $\lambda(\pi(s))$ is in $\mathcal{O}^\times$. (For a ring R , we denote by $R^\times$ the multiplicative group consisting of all units of R .)

Recall from [4, Proposition 1.19] the following correspondence. Similar for E' and G' . Note that for Proposition 2.2, we only need E/G being cyclic.

Proposition 2.2. (Dade) *There is a bijective correspondence between $\text{Irr}(G)^E$ and the set of regular $\hat{F}$ -orbits of irreducible characters of E . By this correspondence, $\phi \in \text{Irr}(G)^E$ corresponds to $\Psi = \text{Irr}(E \mid \phi) = \{\psi \in \text{Irr}(E) \mid [\psi \downarrow_G^E, \phi]_G \neq 0\}$ and $\phi = \psi \downarrow_G^E$ for any $\psi \in \Psi$.*

We recall the correspondence of Dade in the case of $|E/G|$ being a prime (for the statements under the condition that the order of the cyclic group E/G is divided by several primes, see [4, Theorems 6.8 and 6.9]).

Theorem 2.3. (Dade) *There is a bijection*

$$\text{Irr}(G)^E \longrightarrow \text{Irr}(G')^{E'}, \quad \phi \mapsto \phi_{(G')} \quad (*)$$

which satisfies the following:

- When q is odd, there are a unique sign $\epsilon_\phi \in \{\pm 1\}$ and a unique bijection

$$\text{Irr}(E \mid \phi) \longrightarrow \text{Irr}(E' \mid \phi_{(G')}), \quad \psi \mapsto \psi_{(E')} \quad (**)$$

such that

$$(\psi - \lambda^i \psi) \downarrow_{E'}^E = \epsilon_\phi (\psi_{(E')} - \lambda^i (\psi_{(E')})) \quad (***)$$

holds for any i as generalized characters.

- When q is 2, if we choose a sign ϵ_ϕ arbitrary, there is a unique bijection $(**)$ such that $(***)$ holds.

Moreover, $\lambda^i(\psi_{(E')}) = (\lambda^i \psi)_{(E')}$ (hence denoted by $\lambda^i \psi_{(E')}$) for any i .

We call both the correspondences $(*)$ and $(**)$ in Theorem 2.3 the *Glauberman-Dade correspondence* of characters. We denote by $\phi'_{(G)}$ the character of $\text{Irr}(G)^E$ corresponding to $\phi' \in \text{Irr}(G')^{E'}$.

In the remainder of this section, we rewrite the Glauberman-Dade correspondence in terms of the elements of the group algebras (Proposition 2.6).

Let $\mathcal{R} \in (\mathcal{K}, \mathcal{O}, k)$, H a finite group with a subgroup L , and $r_h \in \mathcal{R}$ for $h \in H$. Denote by Pr_L^H an $\mathcal{R}$ -linear map from $\mathcal{R}H$ to $\mathcal{R}L$ defined by $\text{Pr}_L^H(\sum_{h \in H} r_h h) = \sum_{l \in L} r_l l$, which induces an $\mathcal{R}$ -linear map from $Z(\mathcal{R}H)$ to $Z(\mathcal{R}L)$. Denote by Tr_L^H an $\mathcal{R}$ -linear map from $(\mathcal{R}H)^L (\supset Z(\mathcal{R}L))$ to $Z(\mathcal{R}H)$ defined by $\text{Tr}_L^H(\tau) = \sum_{x \in [L \setminus H]} \tau^x$ for $\tau \in (\mathcal{R}H)^L$, where $(\mathcal{R}H)^L$ is the subalgebra of $\mathcal{R}H$ consisting of all the elements fixed by the conjugation action by L and $[L \setminus H]$ is a set of left coset representatives of L in H . For a conjugacy class C of H , we denote $\widehat{C} = \sum_{x \in C} x \in \mathcal{R}H$. For $\theta \in \text{Irr}(H)$, we denote by e_θ the primitive idempotent of $Z(\mathcal{K}H)$ corresponding to θ .

Denote by $C(x)$ the conjugacy class of E containing $x \in E$ and by $C(x)'$ the conjugacy class of E' containing $x' \in E'$.

Since $E_0 = \sqcup_{t \in [E' \setminus E]} (E'_0)^t$ (disjoint union), see [4, Lemma 6.5], we have:

Lemma 2.4. *For $x \in E'_0$, it holds that:*

$$(1) \frac{|C(x)|}{|E|} = \frac{|C(x)'|}{|E'|}.$$

$$(2) \text{Pr}_{E'}^E(\widehat{C(x)}) = \widehat{C(x)'}, \quad \text{Tr}_{E'}^E(\widehat{C(x)'}) = \widehat{C(x)}.$$

For $c \in \mathcal{K}$, we denote by $\bar{c}$ the complex conjugate of c (we view $\mathcal{K}$ as a subfield of the complex number field).

Lemma 2.5. *Let $\phi, \xi \in \text{Irr}(G)^E$, and let $\psi, \eta \in \text{Irr}(E)$ be extensions of ϕ and ξ , respectively. Then, for $x, y \in E'_0$, we have $\psi(x) \overline{\eta(y)} = \epsilon_\phi \epsilon_\xi \psi_{(E')}(x) \overline{\eta_{(E')}(y)}$.*

Proof. Let q be odd. Let $c, d \in \mathbb{Z}$ be such that $\pi(x) = \pi(s)^c$ and $\pi(y) = \pi(s)^d$. Let $c', d' \in \mathbb{Z}$ be such that $\underline{c'} = \underline{c}^{-1}$ and $\underline{d'} = \underline{d}^{-1}$ in $\mathbb{Z}/q\mathbb{Z}$ where the canonical image of $a \in \mathbb{Z}$ in the residue ring $\mathbb{Z}/q\mathbb{Z}$ is denoted by $\underline{a}$. By Theorem 2.3, we have

$$\begin{aligned} q \psi(x) \overline{\eta(y)} &= \sum_{i=1}^{\frac{q-1}{2}} [(\psi - \lambda^{ic'} \psi)(x) \overline{(\eta - \lambda^{id'} \eta)(y)}] \\ &= \epsilon_\phi \epsilon_\xi \sum_{i=1}^{\frac{q-1}{2}} [(\psi_{(E')} - \lambda^{ic'} \psi_{(E')})(x) \overline{(\eta_{(E')} - \lambda^{id'} \eta_{(E')})(y)}] = \epsilon_\phi \epsilon_\xi q \psi_{(E')}(x) \overline{\eta_{(E')}(y)}. \end{aligned}$$

In the case of $q = 2$, we see that $\psi(x) = \epsilon_\phi \psi_{(E')}(x)$ since $(\psi - \lambda\psi)(x) = 2\psi(x)$. Hence, the assertions follow. $\square$

With the notations in section 1, for cyclic $S = \langle s \rangle$, using (#), Okuyama showed in [8]

$$\text{Pr}_{G^S}^G(s^{-1} \widehat{C(s)} e_\chi) = e_{\beta_\chi} \quad \text{for } \chi \in \text{Irr}(G)^S. \quad (\star)$$

We can show the analogous statement which implies $(\star)$ for $|S| = q$ (without using (#) since we treat the Glauberman-Dade correspondence):

Proposition 2.6. *Let $x \in E'_0$. Then*

- (1) $\text{Pr}_{E'}^E(\widehat{C(x)} e_\phi) = \widehat{C(x)'} e_{\phi_{(G')}} \text{ for } \phi \in \text{Irr}(G)^E.$
- (2) $\text{Tr}_{E'}^E(\widehat{C(x)'} e_{\phi'}) = \widehat{C(x)} e_{\phi'_{(G')}} \text{ for } \phi' \in \text{Irr}(G')^{E'}.$

Proof. Let $\psi \in \text{Irr}(E)$ be an extension of ϕ . Then,

$$\begin{aligned} \widehat{C(x)} e_\phi &= \widehat{C(x)} \sum_{i=0}^{q-1} e_{\lambda^i \psi} \\ &= \frac{|C(x)|}{\phi(1)} \sum_{i=0}^{q-1} \lambda^i \psi(x) e_{\lambda^i \psi} \\ &= \frac{|C(x)|}{\phi(1)} \frac{\phi(1)}{|E|} \sum_{i=1}^{q-1} \sum_{y \in E} \lambda^i \psi(x) \overline{\lambda^i \psi(y)} y \\ &= \frac{|C(x)|}{|E|} \sum_{z \in E_0, \pi(z)=\pi(x)} q \psi(x) \overline{\psi(z)} z. \end{aligned}$$

From this and Lemmas 2.4 and 2.5, the assertions follow immediately. $\square$

3

In this section, we show that under Condition 3.1 below (this condition implies G' controls p -fusion in G), the Glauberman-Dade correspondence in the case $|E/G|$ being a prime induces a one-to-one correspondence between the characters belonging to the principal (p -)blocks of G and G' (Proposition 3.3) and an isomorphism of the centers of the principal (p -)block algebras of G and G' (Theorem 3.5).

Condition 3.1. There is some Sylow p -subgroup of E which is contained in G' (hence, $p \neq q$) and centralized by some element s of E'_0 .

Below, we assume Condition 3.1 and take $s \in E'_0$ as in it.

A primitive idempotent of $Z(\mathcal{O}G)$ is called a block (idempotent) of G . Let b (resp. b') be the principal block (idempotent) of G (resp. G'). That is, a primitive idempotent of $Z(\mathcal{O}G)$ such that $1_G(b) \neq 0$ where 1_G is the trivial character of G . (We don't distinguish $\text{Irr}(G)$ and $\text{Irr}(\mathcal{K}G)$.) Denote $\text{Irr}(b) = \{\phi \in \text{Irr}(G) \mid \phi(b) \neq 0\}$. $\phi \in \text{Irr}(b)$ is called a character belonging to the block b . Similar for the other groups and blocks.

By [3] (see also [1]), b has the following primitive idempotent decomposition in $Z(\mathcal{O}Eb)$: $b = \sum_{i=0}^{q-1} \hat{b}_i$. In this situation, $\text{Irr}(b)^E = \text{Irr}(b)$ and the restriction and some extension of the characters determine a one-to-one correspondence between $\text{Irr}(b)$ and $\text{Irr}(\hat{b}_i)$ for any i . Similar for b' .

We denote by $\hat{b}_0$ (resp. $\hat{b}'_0$) the principal block of E (resp. E'), and use $\hat{\cdot}$ for the extensions of characters in $\text{Irr}(b)$ (resp. $\text{Irr}(b')$) belonging to the principal block, that is, $\text{Irr}(\hat{b}_0) = \{\hat{\phi} \mid \phi \in \text{Irr}(b)\}$ (resp. $\text{Irr}(\hat{b}'_0) = \{\hat{\phi}' \mid \phi' \in \text{Irr}(b')\}$).

Then, we see immediately that, for $0 \leq i \leq q-1$, a set $\{\lambda^i \hat{\phi} \mid \phi \in \text{Irr}(b)\}$ (resp. $\{\lambda^i \hat{\phi}' \mid \phi' \in \text{Irr}(b')\}$) forms the set of characters of E (resp. E') belonging to some block of E (resp. E') and we may denote it by $\hat{b}_i$ (resp. $\hat{b}'_i$).

Lemma 3.2. (Osima [9] or see [6, Theorem 12.4.12]) *For a finite group H and $\theta \in \text{Irr}(H)$, θ belongs to the principal p -block of H if and only if $\sum_{x \in H_{p'}} \theta(x) \neq 0$, where $H_{p'}$ is the set of elements of H with the order prime to p .*

Proposition 3.3.

$$(1) \text{Irr}(b') = \{\phi_{(G')} \mid \phi \in \text{Irr}(b)\}$$

$$(2) \text{Irr}(\hat{b}'_i) = \{(\lambda^i \hat{\phi})_{(E')} \mid \lambda^i \hat{\phi} \in \text{Irr}(\hat{b}_i)\} = \{\lambda^i \widehat{\phi_{(G')}} \mid \phi \in \text{Irr}(b)\} \text{ for } 0 \leq i \leq q-1, \\ \text{if we choose signs } \epsilon_\phi \text{ appropriately in the case } q=2.$$

Proof. By Theorem 2.3 and Lemmas 2.4(1) and 3.2, for $\phi \in \text{Irr}(b)$ and $1 \leq i \leq q-1$,

$$\begin{aligned} 0 \neq \sum_{x \in E_{p'}} \hat{\phi}(x) &= \sum_{x \in E_{p'}} (\hat{\phi} - \lambda^i \hat{\phi})(x) = \sum_{x \in (E_0)_{p'}} (\hat{\phi} - \lambda^i \hat{\phi})(x) \\ &= \epsilon_\phi \frac{|E|}{|E'|} \sum_{x' \in (E'_0)_{p'}} (\hat{\phi}_{(E')} - \lambda^i \hat{\phi}_{(E')})(x') = \epsilon_\phi \frac{|E|}{|E'|} \sum_{x' \in E'_{p'}} (\hat{\phi}_{(E')} - \lambda^i \hat{\phi}_{(E')})(x'). \end{aligned}$$

Hence, $\hat{\phi}_{(E')}$ or $\lambda^i \hat{\phi}_{(E')}$ must belong to $\hat{b}'_0$, and so $\phi_{(G')} \in \text{Irr}(b')$. If $q = 2$, we can choose e_ϕ so that $\hat{\phi}_{(E')}$ belongs to the principal block $\hat{b}'_0$. If q is odd, we see that $\hat{\phi}_{(E')} \in \text{Irr}(\hat{b}'_0)$ since $i \neq 0$ is arbitrary.

Hence, in any case, the Glauberman-Dade correspondence in Theorem 2.3 induces injections from $\text{Irr}(b)$ to $\text{Irr}(b')$ and from $\text{Irr}(\hat{b}_0)$ to $\text{Irr}(\hat{b}'_0)$.

Onto part also follows from the similar consideration.

Note that $\hat{\phi}_{(E')} = \widehat{\phi_{(G')}}$. The remaining statement follows from the commutativity of the action of $\hat{F}$ and the Glauberman-Dade correspondence. $\square$

Since $b = \sum_{\phi \in \text{Irr}(b)} e_\phi$ and $b' = \sum_{\phi' \in \text{Irr}(b')} e_{\phi'}$, by Propositions 2.6 and 3.3, we have:

Proposition 3.4. $\text{Pr}_{E'}^E(\widehat{C(x)b}) = \widehat{C(x)'b'}$ and $\text{Tr}_{E'}^E(\widehat{C(x)'b'}) = \widehat{C(x)b}$

We see $\widehat{C(s)b} \in Z(\mathcal{O}Eb)^\times$ and $\widehat{C(s)'b'} \in Z(\mathcal{O}E'b')^\times$. In fact, $\widehat{C(s)b_i} \in Z(\mathcal{O}Eb_i)^\times$ for any i , since $\omega_{b_i}(\widehat{C(s)b_i}) \neq 0$ by Condition 3.1 where ω_{b_i} is a unique algebra homomorphism from the local algebra $Z(\mathcal{O}Eb_i)$ to k whose value at $\widehat{C(s)b_i}$ is the canonical image in k of $\frac{|E|\lambda^i \mathbf{1}_E(s)}{|\widehat{C_E(s)}|\lambda^i \mathbf{1}_E(1)}$.

Note that $bb' \neq 0$. Denote $\gamma = (\widehat{C(s)'b'})^{-1} \widehat{C(s)b} \in (\mathcal{O}Ebb')^{E'}$ and let $\gamma' = \gamma^{-1} = (\widehat{C(s)b})^{-1} \widehat{C(s)'b'}$ the inverse of γ in $(\mathcal{O}Ebb')^{E'}$. Here we mean $(\widehat{C(s)'b'})^{-1}$ the inverse of $\widehat{C(s)'b'}$ in $Z(\mathcal{O}E'b')$ and $(\widehat{C(s)b})^{-1}$ the inverse of $\widehat{C(s)b}$ in $Z(\mathcal{O}Eb)$. Considering the grading given by π and noting $Z(\mathcal{O}Gb)$ is a local algebra, we see that $(\widehat{C(s)b})^q \in Z(\mathcal{O}Gb)^\times$. Hence, we see $(\widehat{C(s)b})^{-1}$ (similar for $(\widehat{C(s)'b'})^{-1}$) is a linear combination of the elements $x \in E$ such that $\pi(x) = \pi(s)^{q-1}$. Hence, $\gamma, \gamma' \in \mathcal{O}G$.

The following in the situation of the Glauberman correspondence appeared in [8].

Theorem 3.5. *With the notations above, there is an $\mathcal{O}$ -algebra isomorphism from $Z(\mathcal{O}Gb)$ to $Z(\mathcal{O}G'b')$ mapping $z \in Z(\mathcal{O}Gb)$ to $\text{Pr}_{G'}^G(\gamma z)$. The inverse is given by the map sending $z' \in Z(\mathcal{O}G'b')$ to $\text{Tr}_{G'}^G(\gamma' z')$.*

Proof. Firstly, note that multiplying $(\widehat{C(s)'b'})^{-1}$ to

$$\widehat{C(s)'b'} e_{\phi_{(G')}} = \widehat{C(s)'e_{\phi_{(G')}}} = \text{Pr}_{E'}^E(\widehat{C(s)e_\phi})$$

where $\phi \in \text{Irr}(b)$ (see Proposition 2.6(1)), we have

$$\begin{aligned} e_{\phi_{(G')}} &= b' e_{\phi_{(G')}} = (\widehat{C(s)'b'})^{-1} \text{Pr}_{E'}^E(\widehat{C(s)e_\phi}) \\ &= \text{Pr}_{E'}^E[(\widehat{C(s)'b'})^{-1} \widehat{C(s)e_\phi}] \\ &= \text{Pr}_{G'}^G[(\widehat{C(s)'b'})^{-1} \widehat{C(s)e_\phi}] \\ &= \text{Pr}_{G'}^G(\gamma e_\phi). \end{aligned}$$

Similarly, multiplying $(\widehat{C(s)b})^{-1}$ to

$$\widehat{C(s)b}e_{\phi'_{(G)}} = \widehat{C(s)}e_{\phi'_{(G)}} = \mathrm{Tr}_{E'}^E(\widehat{C(s)'}e_{\phi'})$$

where $\phi' \in \mathrm{Irr}(b')$ (see Proposition 2.6(2)), we have

$$\begin{aligned} e_{\phi'_{(G)}} &= be_{\phi'_{(G)}} = (\widehat{C(s)b})^{-1}\mathrm{Tr}_{E'}^E(\widehat{C(s)'}e_{\phi'}) \\ &= \mathrm{Tr}_{E'}^E[(\widehat{C(s)b})^{-1}\widehat{C(s)'}e_{\phi'}] \\ &= \mathrm{Tr}_{G'}^G[(\widehat{C(s)b})^{-1}\widehat{C(s)'}e_{\phi'}] \\ &= \mathrm{Tr}_{G'}^G(\gamma'e_{\phi'}). \end{aligned}$$

Then, statements follow as in [8]. The $\mathcal{K}$ -linear map sending $z_{\mathcal{K}} \in Z(\mathcal{K}Gb)$ to $\mathrm{Pr}_{G'}^G(\gamma z_{\mathcal{K}})$ and the $\mathcal{K}$ -linear map sending $z'_{\mathcal{K}} \in Z(\mathcal{K}G'b_{(G')})$ to $\mathrm{Tr}_{G'}^G(\gamma' z'_{\mathcal{K}})$ are mutually inverse $\mathcal{K}$ -algebra isomorphisms between $Z(\mathcal{K}Gb)$ and $Z(\mathcal{K}G'b')$, since $\{e_{\phi} | \phi \in \mathrm{Irr}(b)\}$ and $\{e_{\phi_{(G')}} | \phi \in \mathrm{Irr}(b)\}$ are orthogonal $\mathcal{K}$ -bases of $Z(\mathcal{K}Gb)$ and $Z(\mathcal{K}G'b')$ respectively. Moreover, since these maps send elements with coefficient in $\mathcal{O}$ to elements with coefficient in $\mathcal{O}$ by definitions, these maps restrict to $\mathcal{O}$ -algebra isomorphisms between $Z(\mathcal{O}Gb)$ and $Z(\mathcal{O}G'b')$. $\square$

References

- [1] J. L. Alperin, Isomorphic blocks, *J. Algebra* **43** (1976), 694–698.
- [2] M. Broué, Isométries parfaites, types de blocs, catégories dérivées, *Astérisque* **181-182** (1990), 61–92.
- [3] E. C. Dade, Remarks on isomorphic blocks, *J. Algebra* **45** (1977), 254–258.
- [4] E. C. Dade, A new approach to Glauberman’s correspondence, *J. Algebra* **270** (2003), 583–628.
- [5] G. Glauberman, Correspondence of characters for relatively prime operator groups, *Canad. J. Math.* **20** (1968), 1465–1488.
- [6] G. Karpilovsky, *Group Representations*, Vol. 5 North-Holland Mathematics Studies **183**, North-Holland Amsterdam, 1996.
- [7] H. Nagao, Y. Tsushima, *Representations of Finite Groups*, Academic Press, Boston, 1989.
- [8] T. Okuyama, A talk at a Seminar in Ochanomizu Univ. on 5 November, 2005.
- [9] M. Osima, On some properties of group characters, *Proc. Japan Acad.* **36** (1960), 18–21.
- [10] F. Tasaka, preprint, 2006
- [11] A. Watanabe, The Glauberman character correspondence and perfect isometries for blocks of finite groups, *J. Algebra* **216** (1999), 548–565.