

Highlights

- ▶ We studied the active normal faults in the intracontinental Weihe Graben, China.
- ▶ Active faults are characterized by stepwise fault scarps with dip-angles of 40° – 71° .
- ▶ Active normal fault scarps are distributed in a wide zone of up to 500 m.
- ▶ Structural and topographic features indicate a pure dip-slip sense of the fault zone.
- ▶ Average dip-slip rate is estimated to be ~ 3.0 – 4.0 mm/yr for the normal fault zone.

**Tectonic activity and structural features of active
intracontinental normal faults in the Weihe Graben, central
China**

Gang Rao^{1, 2#}, Aiming Lin^{1*}, Bing Yan^{1, 2}, Zhikun Ren³, Dong Jia⁴, and Xiaojun Wu⁴

¹Department of Geophysics, Graduate School of Science, Kyoto University

Kyoto 606-8502, Japan

²Graduate School of Science and Technology, Shizuoka University

Shizuoka 422-8529, Japan

³State Key Laboratory of Earthquake Dynamics, Institute of Geology, China

Earthquake Administration, Beijing 100029, China

⁴Department of Earth Sciences, Nanjing University

Nanjing 210093, China

#Current address: Department of Earth Sciences, Zhejiang University

Hangzhou 310027, China

***Corresponding author:**

Department of Geophysics

Graduate School of Science, Kyoto University

Kyoto 606-8502, Japan

Tel: 81-75-753-3941

E-mail: slin@kugi.kyoto-u.ac.jp (A. Lin)

Abstract

This study examines the tectonic activity and structural features of active normal faults in the Weihe Graben, central China. The Weihe Graben is an area with a high level of historic seismicity, and it is one of the intracontinental systems that developed since Tertiary in the extensional environment around the Ordos Block. Analysis of high-resolution remote-sensing imagery data, field observations, and radiocarbon dating results reveal the following: i) active normal faults are mainly developed within a zone <500 m wide along the southern border of the eastern part of the Weihe Graben; ii) the active faults that have been identified are characterized by stepwise fault scarps dipping into the graben at angles of 40°–71°; iii) there are numerous discontinuous individual fault traces, ranging in length from a few tens of meters to 450 m (generally <200 m); iv) fault zone structures, topographic features, and fault striations on the main fault planes indicate almost pure normal-slip; and v) late Pleistocene–Holocene terrace risers, loess, and alluvial deposits have been vertically offset by up to ~80 m, with an average dip-slip rate (throw-rate) of ~3.0–4.0 mm/yr. Our results reveal that active normal faults have been developing in the Weihe Graben under an ongoing extensional environment, probably associated with the pre-existing graben and spreading of the continental crust, and this is in contrast with the Ordos Block and neighboring orogenic regions. These results provide new insights into the nature of extensional tectonic deformation in intracontinental graben systems.

Keywords: active intracontinental normal fault, fault scarp, tectonic activity, Weihe Graben, Ordos Block

1. Introduction

The Tibetan Plateau has been uplifted due to the ongoing collision between the Indian and Eurasia plates since the Early Eocene (e.g., Harrison et al., 1992; Yin and Harrison, 2000). The plateau and neighboring regions in East Asia are ideal places for studying active tectonics, because the orogeny is still active and the collision has produced a variety of geological features including large-scale thrust, normal, and strike-slip fault systems (e.g., Tapponnier and Molnar, 1977; Molnar and Deng, 1984; Armijo et al., 1986; Molnar and Lyon–Caen, 1989; Liu et al., 2004). The co-seismic rupture structures associated with thrusting and strike-slip movements, in and around the Tibetan Plateau (Fig. 1), have been well investigated on the basis of numerous large earthquakes, including the 2008 M_w 7.9 Wenchuan earthquake due to thrusting (e.g., Lin et al., 2009; Jia et al., 2010; Li et al., 2010), the 2001 M_w 7.8 Kunlun earthquake due to strike-slip movement (e.g., Lin et al., 2002), and the 2010 M_w 6.9 Yushu earthquake, also due to strike-slip movement (e.g., Lin et al., 2011; Rao et al., 2011; Guo et al., 2012). In contrast, we have relatively less understanding of the tectonic geomorphology and structural features related to active intracontinental normal faults, or the co-seismic rupturing mechanisms of such faults (e.g., Armijo et al., 1986; Sun et al., 2011). Although previous studies have reported some structural features of co-seismic surface ruptures associated with normal faulting locally such as western USA (e.g., Myers and Hamilton, 1964; Wallace, 1984; Crone et al., 1987), large earthquakes caused by normal faulting are still rare worldwide in contrast with the thrust-type and strike-slip earthquakes. Recently, the 2011 M_w 6.6 Fukushima

1 earthquake, one of the aftershocks triggered by the 2011 M_w 9.0 Tohoku earthquake in
2
3 Japan, generated a ~23-km-long surface rupture zone along the pre-existing active
4
5 normal faults developed in the intracontinental rift environment characterized by thick
6
7 half-graben sediments as that of the Weihe Graben, the target of this study, which
8
9 provides a typical example of large earthquake triggered by normal faulting (e.g., Lin
10
11 et al., 2013a; Toda and Tsutsumi, 2013).
12
13
14
15
16

17 The Weihe Graben is one of the intracontinental graben systems that has
18
19 developed around the Ordos Block in central China since the Eocene, and it provides
20
21 a unique natural laboratory for studying the long-term tectonic history of active
22
23 intracontinental normal-faults in an extensional environment (Fig. 1). In previous
24
25 studies, the active faults in this region have been traced as a simple linear structure
26
27 along the topographic boundary between the Weihe Graben and the Qinling and
28
29 Huashan Mountains (e.g., Li and Ran, 1983; Zhang et al., 1995; Hou et al., 1998;
30
31 Yuan and Feng, 2010), but the surface traces, structural features, and geometric
32
33 distribution of the active faults were not mapped in detail and hence remain unclear.
34
35 Although it has been inferred that the active faults in the Weihe Graben are dominated
36
37 by normal slip (SSB, 1988; Deng et al., 2003), a large component of sinistral
38
39 strike-slip movement has also been suggested (e.g., Li and Ran, 1983; Wang, 1987;
40
41 Zhang et al., 1998; Mercier et al., 2013), and it seems, therefore, that the sense of
42
43 tectonic movement is also an issue in dispute.
44
45
46
47
48
49
50
51
52
53
54

55 In this study, we focus on the recent activity, geometric distribution pattern, and,
56
57 structural features of the active normal faults in the Weihe Graben, and discuss their
58
59
60
61
62
63
64
65

1 implications for the extensional tectonic deformation in the intracontinental rift basin
2
3 systems around the Ordos Block, central China.
4
5
6
7

8 9 **2. Tectonic setting**

10
11 The Weihe Graben is located at the southern margin of the stable Ordos Block, which
12
13 has a pre-Mesozoic crystalline basement (Ma and Wu, 1987; SSB, 1988). The graben
14
15 is bordered to the south by the Qinling Mountains, which were formed by the collision
16
17 of the North China Craton (NCC) and the South China Block (SCB) in the Triassic
18
19 (Fig. 1; e.g., Meng and Zhang, 2000; Rastchbacher et al., 2003). The Weihe Graben is
20
21 just one of several intracontinental graben systems that are developed around the
22
23 Ordos Block (Fig. 1). As a result of extension, the Weihe Graben has received
24
25 Cenozoic deposits up to 7000 m in thickness since the Eocene, and the extension has
26
27 been accompanied by the uplift of mountainous blocks (e.g., the Huashan Mountains
28
29 and the Weinan Loess Tableland) along its southern border (SSB, 1988; Zhang et al.,
30
31 1998; Liu et al., 2013). The upthrown Huashan Mountains are mainly composed of
32
33 pre-Mesozoic metamorphic basement rocks; in contrast, the Weinan Loess Tableland
34
35 is characterized by Quaternary loess and alluvial deposits (Fig. 1; SSB, 1988). Fission
36
37 track dating data have shown that the uplift of the Huashan Mountains started at
38
39 around 68.2 Ma, and that uplift rates became faster from around 17.8 Ma, with an
40
41 average rate of ~0.19 mm/yr (Yin et al., 2001).
42
43
44
45
46
47
48
49
50
51
52
53
54
55

56 More than 10 large historical earthquakes of $M \geq 7$, including four with $M \geq 8$,
57
58 have occurred in the graben systems around the Ordos Block, three of those occurred
59
60
61
62
63
64
65

1 in the Weihe Graben (Fig. 1; Deng, 2007). The 1556 M ~8.5 Huaxian great
2
3 earthquake occurred in the Weihe Graben and caused >830,000 deaths, and it has been
4
5 suggested that an active fault zone ruptured for up to 70 km along the southern margin
6
7 of the Ordos Block between the cities of Weinan and Huayin (e.g., Kuo, 1957; Wang,
8
9 1980; SSB, 1988; Xie, 1992; CENC, 2007; Yuan and Feng, 2010). The high level of
10
11 historical seismicity indicates that the normal faults in the Weihe graben are currently
12
13 active as seismogenic faults (e.g., SSB, 1988; Zhang et al., 1998; Deng, 2007).
14
15
16
17
18
19
20
21
22

23 **3. Identification of active faults**

24
25 Since crustal deformation associated with active faults is represented by displaced
26
27 landforms on the surface of the Earth, a perspective view of topographic features
28
29 provides an important method of detecting and identifying active faults. In this study,
30
31 we identified active faults by using perspective view of high-resolution satellite
32
33 remote-sensing images (0.5 m WorldView and 1 m IKONOS), together with
34
35 fieldwork that was guided by the images. The remote-sensing images were processed
36
37 and analyzed in 3D by draping on the 30 m resolution ASTER Global Digital
38
39 Elevation Model (GDEM) data (Figs. 2–4). These multi-perspective views made it
40
41 possible to identify the active faults more easily than by using traditional methods
42
43 such as aerial photographs. On the basis of geomorphic markers such as terrace risers
44
45 and alluvial fans, we were able to identify active fault traces that are mostly
46
47 perpendicular or oblique to valleys and river channels. As shown in Fig. 2, a series of
48
49 irregular linear traces is developed near the heads of the alluvial fans along the
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 piedmont of the Huashan Mountains, and the traces are characterized by a series of
2
3 parallel or sub-parallel fault scarps that face NNW. The topographic and structural
4
5 features of the fault scarps indicate that faulting has taken place since the formation of
6
7 the alluvial fans, and in combination with the results of radiocarbon dating, it can
8
9 therefore be demonstrated that the observed faults have been active in the Holocene
10
11
12
13
14 (Fig. 2; Table 1).
15
16
17
18
19

20 **5. Structural features and activity of the recent faults**

21

22 The analysis of the remote sensing images, together with field investigations, shows
23
24 that the active faults are mainly distributed along the northern marginal zones of the
25
26 Huashan Mountains and the Weinan Loess Tableland. Numerous discontinuous fault
27
28 traces are present, and they range in length from a few tens of meters to 450 m
29
30 (generally <200 m) (Figs. 2–4). These faults are mainly characterized by numerous
31
32 stepped normal fault scarps, which are mostly restricted to a narrow zone 10–500 m
33
34 wide (generally <300 m), and the spacing of the individual fault scarps is in the range
35
36 ~20–200 m (generally <100 m) (Figs. 2–5). Along these scarps, many outcrops of the
37
38 faults can be observed in the field, and four typical examples (Locs. 1–4), where the
39
40 loess, alluvial deposits, and basement rocks are displaced, are described below (Figs.
41
42 6–11).
43
44
45
46
47
48
49
50
51
52

53 At Loc. 1, the stepped fault scarps can be seen along the piedmont of the Huashan
54
55 Mountains near Huayin city, and the height of each individual fault scarp is as much
56
57 as 19 m (Fig. 6a–b; see Figs. 1c and 4b for details of the location). Under one scarp,
58
59
60
61
62
63
64
65

1 the main shear zone of the fault crops out, and fault breccia and cataclasite sourced
2
3 from the basement rocks, together with disturbed loess and alluvial deposits, are
4
5 bounded by distinct fault planes that dip NW at angles of 40°–54° (Fig. 6c, d). The
6
7 fault breccias are mostly dragged into positions parallel or subparallel to the main
8
9 fault planes, indicating a slip-sense that is dominantly normal (Fig. 6c, d). Some
10
11 brownish-gray veins composed of fine-grained microbreccia have been injected into
12
13 both the loess and sandy gravel deposits, indicating a recent faulting event that cuts all
14
15 the deposits at this site (Fig. 6d, e). Such injection veins within active fault shear
16
17 zones have commonly been reported elsewhere, and they are generally considered to
18
19 form rapidly during large paleo-earthquakes, indicating the co-seismic rupture
20
21 structure of seismogenic fault zone (e.g., Lin et al., 2012, 2013b).
22
23
24
25
26
27
28
29
30

31 At Loc. 2, the shear zone of the fault is well exposed in four parallel quarry pits
32
33 (T1–T4) on the piedmont of the Huashan Mountains (Figs. 7–8; see Figs. 2 and 3a for
34
35 details of the location). The shear zone is exposed along a fault scarp, approximately
36
37 10 m in height that is developed in the alluvial sediments and loess deposits (Fig. 7).
38
39 Sketches of the shear zone structures in seven exposures are shown in Fig. 7c; the
40
41 structures are concentrated in a zone less than 5 m in width, and 3–5 sub-faults are
42
43 present. The loess deposits and alluvial sand gravels are downthrown to the north
44
45 along the main fault planes that dip north at angles of 40°–71°, indicating a normal
46
47 slip-sense, consistent with the topographic features of the stepped fault scarps (Fig.
48
49 7c). All the deposits within the shear zone have been disturbed, and the gravels were
50
51 mostly dragged into positions parallel or subparallel to the main fault planes, with
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 some injected into fractures or forming wedge-shaped structures (Figs. 7c and 8). The
2
3 calcareous materials collected in the loessial deposits which are interlayered by the
4
5 loess and alluvial deposit composed of sandy gravels, yield radiocarbon ages of
6
7
8 ~5500–19,450 yr BP (Fig. 7c; Table 1), indicating late Pleistocene–Holocene activity
9
10
11 of the faults at this site.
12

13
14 At Loc. 3, the shear zone of the fault was observed at the boundary between basin
15
16 and mountain, ~600 m east of Loc. 2. Here, the fault plane dips north at an angle of
17
18 56°, and it separates unconsolidated alluvial deposits in the hanging wall from
19
20 basement rocks in the footwall (Fig. 9; see Figs. 2 and 3a for details of the location).
21
22 The shear zone is characterized by a narrow fault gouge and breccia zone less than 30
23
24 cm thick where the sheared material has been mainly dragged into positions parallel
25
26 or subparallel to the fault plane. Fault steps and striations are present, indicating a
27
28 mainly normal sense of slip (Fig. 9b–e). The alluvial deposits that bound the shear
29
30 zone were disturbed and mixed with the fault breccias, and a downthrown movement
31
32 of hanging wall is indicated, which is also consistent with that indicated by the fault
33
34 striations and fault steps observed on the main fault plane.
35
36
37
38
39
40
41
42
43

44 At Loc. 4, fault scarps are developed on the terrace risers of the Chishuihe River
45
46 along the northern marginal zone of the Weinan Loess Tableland. The scarps are
47
48 clearly seen in the remote-sensing perspective views (Fig. 10a), and they can also be
49
50 seen in the field (Fig. 10b–d). The fault outcrops are exposed along both banks of the
51
52 Chishuihe River, where alluvial sand–pebble and cobble deposits are faulted against
53
54 weakly consolidated silts and sandstones. The main fault plane strikes N40°W and
55
56
57
58
59
60
61
62
63
64
65

dips NE at an angle of 55° . A fault gouge zone less than 5 cm thick is present, and fault striations indicate a mainly normal sense of slip, consistent with the sense of movement indicated by the fault scarps at this site (Figs. 10e and 11). The pebbly deposits that bound the main fault plane were mostly dragged into positions parallel or subparallel to the fault plane (Fig. 11a, b). The equal-area projection of measured striations and topographic features indicates an almost pure normal fault movement at this site (Fig. 11c). The deposits of Unit 4 are overlaid by Unit 1-3 in which the current plastic materials are involved, indicating a formation age of <100 years. The results of ^{14}C age dating show that the alluvial sediments (Unit 5) in the hanging wall were deposited at $18,270 \pm 60$ yr B.P. indicating activity of the fault since the late Pleistocene (Fig. 11b; Table 1).

6. Discussion

6.1. Geometric characteristics of active normal faults

Normal faults in the upper crust generally have high dip angles of $\sim 60^\circ$, and most large-magnitude earthquakes caused by normal faulting are triggered by faults with dips of $>30^\circ$ (e.g., Jackson, 1987; Doser and Smith, 1989). However, many normal faults have a listric geometry, and flatten out with depth (e.g., Shelton, 1984; Xiao and Suppe, 1992); in addition, some seismogenic faults are known to be low-angle normal faults (e.g., Abers et al., 1997; Axen, 1999; Abbott et al., 2001), although the mechanism by which they trigger earthquakes is still under debate (e.g., Jackson and White, 1989; Lecomte et al., 2012). As stated above, the dip angles of the main faults

1 in the Weihe graben vary from 40° to 71°, but are mostly >50°, consistent with the
2
3 subsurface fault geometry revealed by seismic reflection data (Xie et al., 2011).
4
5
6 Previous study also showed that the dip angles of the active faults in the study area
7
8 range from 45° to 80°, which were measured in the field (SSB, 1988). It seems clear,
9
10 therefore, that the active normal faulting in the study area is not on low-angle
11
12 structures at the near-surface, reflecting the middle-high dip angle of seismogenic
13
14 fault zone.
15
16
17
18
19

20 In plan view, the surface traces of active faults in the study area are characterized
21
22 by numerous discontinuous and short lineaments with variable trends, and they are
23
24 mostly distributed in a zone 10–500 m wide (Figs. 2–4). These geometric features
25
26 contrast with those reported in previous studies where the surface fault traces were
27
28 inferred to form a simple straight lineament along the topographic boundary between
29
30 the mountains and the basin; however, the geometry and distribution pattern of the
31
32 active faults were not mapped in detail (e.g., Li and Ran, 1983; Zhang et al., 1995;
33
34 Hou et al., 1998). Previous studies also show that a single fault zone and/or a
35
36 co-seismic surface rupture zone occurred along a pre-existing active fault is generally
37
38 characterized by numerous fault stands formed a zone ranging from a few meters to
39
40 hundreds meters in width at surface (e.g., Steven et al., 1996; Lin et al., 2002, 2009;
41
42 Kim et al., 2004; Berg and Skar, 2005). The parallel–subparallel active fault traces
43
44 identified in the remote-sensing images have also been confirmed in the field, as
45
46 noted above, and they are characterized by stepwise fault scarps with similar
47
48 orientations, and they form steep slopes in topographic profiles (Figs. 2–5). Such
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

geometric features are comparable with those of typical active normal fault zones in extensional tectonic regimes worldwide (e.g., Armijo et al., 1986; Yeats et al., 1997; McCalpin, 2009). Based on geological and geophysical data, the stepwise fault scarps are generally considered to be a response to the refraction of master fault-planes bounding the graben (e.g., Gibbs, 1984; Stewart and Hancock, 1991; McCalpin, 2009). In the study area, seismic reflection data also show that the stepwise faults join together in a master fault below the surface (e.g., Xie et al., 2011). Our findings show that the active normal faults developed in the Weihe Graben have irregular geometries, trends, distributions, and continuity, and at the surface they do not form a regular straight lineament as previously reported.

6.2. Slip rates

The slip rate of an active fault is an important and quantitative parameter for assessing the seismic hazard and tectonic activity of the fault, including the recurrence interval of large earthquakes (e.g., Yeats et al., 1997; McCalpin, 2009). Two factors generally affect the precision of slip-rate estimates in extensional regimes: the actual amount by which topographic markers have been offset, and the age of any surface displacement marker. An estimate of the amount of offset mainly depends on the distribution pattern and geometric characteristics of the active fault, as determined by field mapping, especially in regions, such as the present study area, that are characterized by a stepwise distribution of faults. Neighboring normal faults with similar orientations commonly slip at different rates, and they may be active over different time intervals

(Cowie and Roberts, 2001); in some cases it may be difficult to distinguish a tectonic fault from a gravitational fault such as the slide surface formed during landsliding (e.g., Moro et al., 2012; Carbonel et al., 2013). Therefore, care must be taken when choosing sites for assessing slip rates.

Previous researchers have estimated slip rates for the active normal faults in the Weihe Graben of 1–3 mm/yr (Li and Ran, 1983; SSB, 1988; Deng et al., 2003) to 10.4 mm/yr (Li and Ran, 1983). These estimates differ by one order of magnitude, and they remain open to dispute; clearly, the slip rates of the active normal faults in the Weihe Graben are poorly constrained. These slip rates were generally estimated on the presumption that the faults formed simple straight lineaments. There was no detailed mapping of the geometric patterns and fault distribution, and no accurate dating of surface markers; instead, various inferred ages of displaced landforms were used, involving large uncertainties (Li and Ran, 1983; SSB, 1988). In other words, the large differences among the previous calculations of slip rates probably resulted from a lack of knowledge of the detailed geometry of the active faults, and a lack of reliable age constraints for the surface markers that are offset and deformed. Variation of slip-rates (extension rate of the Graben) might also be affected by the inherited Mesozoic and Cenozoic structures with complex deformation history (e.g., Mercier et al., 2013), as well as the far field effects of the India-Asian collision (e.g., Zhang et al., 1998; Liu et al., 2004).

As stated above, the late Pleistocene–Holocene terrace risers, alluvial fans, and loess slopes have all been systematically displaced, and we have used them as

1 topographic surface markers for estimating slip rates. The loess is wind-borne
2
3 sediment that contains calcareous materials that can be used for radiocarbon dating
4
5
6 (Ding et al., 1994), and this allows the loess sequences to be used for assessing the
7
8 ages of surface markers such as the loess plateau surface and alluvial fans.
9
10 Radiocarbon dating results give ages that are generally younger than 20 ka for the
11
12 alluvial fans and the loess plateau surfaces on which the stepwise fault scarps are
13
14 developed (Figs. 7 and 11; Table 1).
15
16

17
18 On the basis of our observations, the active deformation zones are mainly
19
20 characterized by a series of stepped fault scarps. Therefore, the total vertical offset can
21
22 be estimated by summing the vertical offsets of individual fault scarps across the fault
23
24 zone (Fig. 5). To avoid the effects of variations in the geometric distribution of the
25
26 fault traces, we selected six typical topographic profiles across the fault zone to
27
28 measure the total vertical offset amounts; the profiles range from 15 m to 80 m in
29
30 length (generally 30–40 m) (Fig. 5; Table 2). The results of radiocarbon dating show
31
32 that the loess plateau surfaces and alluvial terrace risers formed at around 5500 to
33
34 16,000 yr. B.P. (Tables 1 and 2). Accordingly, the slip rates are calculated to be in the
35
36 range 2.1–5.7 mm/yr, with an average value of ~4 mm/yr (Table 2). Our results
37
38 contrast with those previously estimated (between 1–3 mm/yr and 10.4 mm/yr).
39
40
41
42
43
44
45
46
47
48
49
50
51
52

53 **6.3. Tectonic implications for intracontinental rift basins**

54
55 Understanding the geometric and kinematic characteristics of active normal faults
56
57 in the Weihe Graben would assist in investigating seismotectonic features and the
58
59
60
61
62
63
64
65

1 tectonic–sedimentary evolution of active extensional basins around the Ordos Block
2
3 (e.g., Peltzer et al., 1985; Zhang et al., 1985, 1998). The active normal faults
4
5 identified in this study are mostly developed along the northern margins of the
6
7
8
9 Huashan Mountains and Weinan Loess Tableland, and they are characterized by pure
10
11 dip-slip movements (Fig. 12a) that indicate a NW–SE extensional stress-direction,
12
13 consistent with the regional extensional stress-direction inferred from earthquake
14
15 mechanisms (Ma et al., 1989). Moreover, this result is in general agreement with GPS
16
17 observations that show crustal movement velocities within the entire North China
18
19
20
21 Craton decelerate gradually from WNW to ESE (e.g., Wang et al., 2001). Furthermore,
22
23 the complex geometry and distribution pattern of the surface fault trends mapped in
24
25 the study area show that the map patterns of normal faults are highly irregular, similar
26
27
28 to those of co-seismic surface zones caused by large earthquakes in the
29
30
31 intracontinental rift environments, such as the 1954 Dixie Valley earthquake (M_s 7.2)
32
33
34 and 1915 Pleasant Valley earthquake (M_s 7.6) (Yeats et al., 1997), which provide us
35
36
37 the mapped patterns of fault trends for assessing the normal faulting process and slip
38
39
40 rate of active normal faults that triggered a giant earthquake ($M \sim 8.5$) in the Weihe
41
42
43 Graben as documented above.
44
45

46
47 The Weihe Graben can be classified as a traditional rift basin (e.g., Friedmann
48
49 and Burbank, 1995), characterized by thick half-graben sediments (>7000 m), a long
50
51 history (~ 50 Ma) (e.g., SSB, 1988; Zhang et al., 1998; Liu et al., 2013), and normal
52
53 faults with high dip angles (40° – 71° ; this study). The variations in the lithospheric
54
55 structure (crustal thinning) of the Weihe rift basin relative to its neighboring regions
56
57
58
59
60
61
62
63
64
65

1 are probably the result of extension in the lower crust due to underlying
2
3 asthenospheric mantle flow (Fig. 12b), as revealed by geophysical observations (e.g.,
4
5 Huang et al., 2008; Bao et al., 2011). On the basis of numerical simulations and shear
6
7 wave splitting analysis, it has been suggested that the structures in the mantle were
8
9 affected by the collision of the North and South China blocks in the Mesozoic, with
10
11 the lateral flow of the mantle caused by driving forces associated with Indo-Asian
12
13 collision (Liu et al., 2004; Huang et al., 2008).
14
15
16
17
18
19

20 The active normal faults are currently under an ongoing extensional environment,
21
22 which is probably related to the pre-existing spreading and rifting of the continental
23
24 crust in this area (in contrast to the Ordos Block and other neighboring orogenic
25
26 regions) (Fig. 12b). Because of the absence of magma activity, the strain accumulated
27
28 within the brittle upper crust is thought to have been released mainly by repeated
29
30 movements along the active normal faults. These movements have caused major
31
32 earthquakes such as the 1556 M ~8.5 Huaxian great earthquake, the relative uplift of
33
34 adjacent mountainous blocks (e.g., the Huashan Mountains and the Weinan Loess
35
36 Tableland), and the accumulation of a thick succession of sediments in the rift basin
37
38 (Fig. 12). However, more work is required if we are to have a better understanding of
39
40 the nature of the deformation associated with the seismogenic source fault that
41
42 triggered the 1556 earthquake, and if we wish to improve our assessments of the
43
44 seismic hazards within the densely populated area of the Weihe Graben.
45
46
47
48
49
50
51
52
53
54

55 Similar lithospheric structures have been observed in association with other rift
56
57 basins around the Ordos Block, such as in the Yinchuan Graben (e.g., Zhang et al.,
58
59
60
61
62
63
64
65

1985; Ye et al., 1987; SSB, 1988; Tian et al., 2011). We suggest, therefore, that our findings concerning the tectonic activity and structural features of active normal faults in the Weihe Graben, in combination with geophysical observations of the underlying lithospheric structures, will assist in understanding the seismogenic mechanisms of other large intracontinental earthquakes caused by normal-faulting around the Ordos Block (e.g., Zhang et al., 1985; Huang et al., 2008; Bao et al., 2011), and also provide insights into the Cenozoic extensional deformation in these graben systems (e.g., SSB, 1988; Zhang et al., 1998; He et al., 2004).

7. Conclusions

On the basis of the analysis of remote-sensing images, combined with field investigations, we have reached the following conclusions.

- (i) Active normal faults are mainly developed within a zone <500 m wide along the southern border of the eastern part of the Weihe Graben.
- (ii) The observed active faults are characterized by a distinctive series of stepped fault scarps that dip into the graben at angles of 40°–71°. The fault traces themselves are discontinuous, with individual lengths ranging from a few tens of meters to 450 m (generally <200 m).
- (iii) Topography, geomorphic features, and striations developed on the main fault planes indicate an almost pure normal sense of slip.
- (iv) Late Pleistocene–Holocene terrace risers, loess, and alluvial fan deposits have been offset vertically by up to tens of meters, and the estimated

average dip-slip rate (throw-rate) is 3.0–4.0 mm/yr.

- (v) The normal faults in the Weihe Graben are currently active under an ongoing extensional regime, which is probably related to the pre-existing spreading and rifting of the continental crust in this area (in contrast to the Ordos Block and other neighboring orogenic regions).

Acknowledgements

We thank Earth Remote Sensing Data Analysis Center (ERSDAC) for making ASTER GDEM data freely available from their web site. This work was supported by the Science Project (Project no. 23253002 awarded to A. Lin) from the Ministry of Education, Culture, Sports, Science and Technology of Japan and partially by the National S & T Major Project of China (2011ZX05003-002 and 2011ZX05009-001).

References

- Abbott, R.E., Louie, J.N., Caskey, S.J. and Pullammanappallil, S., 2001. Geophysical confirmation of low-angle normal slip on the historically active Dixie Valley fault, Nevada. *Journal of Geophysical Research* 106, 4169-4181.
- Abers, G.A., Mutter, C.Z. and Fang, J., 1997. Shallow dips of normal faults during rapid extension: Earthquakes in the Woodlark-D'Entrecasteaux rift system, Papua New Guinea. *Journal of Geophysical Research* 102, 15301-15317.
- Armijo, R., Tapponnier, P., Mercier, J.L. and Tong-Lin, H., 1986. Quaternary extension in southern Tibet: field observations and tectonic implications. *Journal*

- of Geophysical Research 91, 13803-13872.
- Axen, G.J., 1999. Low-angle normal fault earthquakes and triggering. *Geophysical Research Letters* 26, 3693-3696.
- Bao, X., Xu, M., Wang, L., Mi, N., Yu, D., and Li, H., 2011. Lithospheric structure of the Ordos Block and its boundary areas inferred from Rayleigh wave dispersion. *Tectonophysics* 499, 132-141.
- Berg, S.S. and Skar, T., 2005. Controls on damage zone asymmetry of a normal fault zone: outcrop analyses of a segment of the Moab fault, SE Utah. *Journal of Structural Geology* 27, 1803–1822.
- CENC (China Earthquake Networks Center), 2007, The 1556 Huaxian great earthquake, Shaanxi, China: the largest total of fatalities ever claimed (in Chinese). Available online at: http://www.csi.ac.cn/manage/html/4028861611c5c2ba0111c5c558b00001/_history/hxz/qyzhenhai/zh20060609002.htm (Last accessed 10 October 2013)
- Carbonel, D., Gutiérrez, F., Linares, R., Roqué, C., Zarroca, M., McCalpin, J., Guerrero, J., and Rodríguez, V., 2013. Differentiating between gravitational and tectonic faults by means of geomorphological mapping, trenching and geophysical surveys. The case of the Zenzano Fault (Iberian Chain, N Spain). *Geomorphology* 189, 93-108.
- Crone, A.J., Machette, M.N., Bonilla, M.G., Lienkaemper, J.J., Pierce, K.L., Scott, W.E., Bucknam, R.C., 1987. Surface faulting accompanying the Borah Peak earthquake and segmentation of the lost river fault, central Idaho. *Bulletin of the*

- Seismological Society of America 77, 739-770.
- Cowie, P.A., and Roberts, G.P., 2001. Constraining slip rates and spacings for active normal faults. *Journal of Structural Geology* 23, 1901-1915.
- Deng, Q., 2007. Active Tectonics Map of China, Seismological Press (in Chinese).
- Deng, Q., Zhang, P., Ran, Y., Yang, X., Min, W., and Chu, Q., 2003. Basic characteristics of active tectonics of China. *Science In China Series D* 46, 356-372.
- Ding, Z.L., Yu, Z., Rutter, N., Liu, T., 1994. Towards an orbital time scale for Chinese loess deposits. *Quaternary Science Reviews*, 31, 39-70.
- Doser, D.I. and Smith, R.B., 1989. An assessment of source parameters of earthquakes in the cordillera of the western United States. *Bulletin of the Seismological Society of America* 79, 1383 -1409.
- Friedmann, S.J. and Burbank, D.W., 1995. Rift basins and supradetachment basins: intracontinental extensional end-members. *Basin Research* 7, 109-127.
- Gibbs, A.D., 1984. Structural evolution of extensional basin margins. *Journal of the Geological Society* 141, 609-620
- Guo, J., Zheng, J., Guan, B., Fu, B., Shi, P., Du, J., Xie, C., and Liu, L., 2012. Coseismic Surface Rupture Structures Associated with 2010 M_s 7.1 Yushu Earthquake, China. *Seismological Research Letters* 83, 109 -118.
- Harrison, T.M., Copeland, P., Kidd, W.S.F. and Yin, A., 1992. Raising Tibet. *Science* 255, 1663-1670.
- Harvard University (2014). Global CMT Catalog Search. Available online at:

- 1 <http://www.globalcmt.org/CMTsearch.html> (last accessed 5 April 2014).
- 2
- 3 He, J., Cai, D., Li, Y. and Gong, Z., 2004. Active extension of the Shanxi rift, north
- 4
- 5
- 6 China: does it result from anticlockwise block rotations? *Terra Nova* 16, 38-42.
- 7
- 8
- 9 Hou, J., Han, M., Chai, B. and Han, H., 1998. Geomorphological observations of
- 10
- 11 active faults in the epicentral region of the Huaxian large earthquake in 1556 in
- 12
- 13 Shaanxi Province, China. *Journal of Structural Geology* 20, 549-557.
- 14
- 15
- 16
- 17 Huang, Z., Xu, M., Wang, L., Mi, N., Yu, D., and Li, H., 2008. Shear wave splitting in
- 18
- 19 the southern margin of the Ordos Block, north China. *Geophysical Research*
- 20
- 21 *Letters* 35, L19301. <http://dx.doi.org/10.1029/2008GL035188> .
- 22
- 23
- 24
- 25 Jackson, J.A., 1987. Active normal faulting and crustal extension. Geological Society,
- 26
- 27 London, Special Publications 28, 3-17.
- 28
- 29
- 30
- 31 Jackson, J.A. and White, N.J., 1989. Normal faulting in the upper continental crust:
- 32
- 33 observations from regions of active extension. *Journal of Structural Geology* 11,
- 34
- 35
- 36 15-36.
- 37
- 38
- 39 Jia, D., Li, Y., Lin, A., Wang, M., Chen, W., Wu, X., Ren, Z., Zhao, Y., and Luo, L.,
- 40
- 41
- 42 2010. Structural model of 2008 M_w 7.9 Wenchuan earthquake in the rejuvenated
- 43
- 44 Longmen Shan thrust belt, China. *Tectonophysics* 491, 174-184.
- 45
- 46
- 47
- 48 Kim, Y.-S., Peacock, D.C.P., Sanderson, D.J., 2004. Fault damage zones. *Journal of*
- 49
- 50 *Structural Geology* 26, 503–517.
- 51
- 52
- 53
- 54
- 55
- 56 Kuo, T., 1957. On the Shensi earthquake of January 23, 1556. *Acta Geophysica Sinica*,
- 57
- 58 6, 59-68 (in Chinese with English abstract).
- 59
- 60
- 61
- 62
- 63
- 64
- 65
- Lecomte, E., Le Pourhiet, L. and Lacombe, O., 2012. Mechanical basis for slip along

1 low-angle normal faults. Geophysical Research Letters 39, L03307.

2
3 <http://dx.doi.org/10.1029/2011GL050756>.

4
5
6 Li and Ran, 1983. Active faults along the north margins of Huashan and Weinan
7
8 Loess Tableland. North China Earthquake Science 1, 10-18 (in Chinese with
9
10 English abstract).

11
12
13 Li, Y., Jia, D., Shaw, J.H., Hubbard, J., Lin, A., Wang, M., Luo, L., Li, H., and Wu, L.,
14
15 2010. Structural interpretation of the coseismic faults of the Wenchuan
16
17 earthquake: Three-dimensional modeling of the Longmen Shan fold-and-thrust
18
19 earthquake: Three-dimensional modeling of the Longmen Shan fold-and-thrust
20
21 belt. Journal of Geophysical Research 115, B04317.
22
23 <http://dx.doi.org/10.1029/2009JB006824>

24
25
26
27 Lin, A., Fu, B., Guo, J., Zeng, Q., Dang, G., He, W., and Zhao, Y., 2002. Co-seismic
28
29 strike-slip and rupture length produced by the 2001 M_s 8.1 central Kunlun
30
31 earthquake. Science 296, 2015-2017.

32
33
34
35 Lin, A., Rao, G., Jia, D., Wu, X., Yan, B., and Ren, Z., 2011. Co-seismic strike-slip
36
37 surface rupture and displacement produced by the 2010 M_w 6.9 Yushu
38
39 earthquake, China, and implications for Tibetan tectonics. Journal of
40
41 Geodynamics 52, 249-259.

42
43
44
45 Lin, A., Ren, Z., Jia, D. and Wu, X., 2009. Co-seismic thrusting rupture and slip
46
47 distribution produced by the 2008 M_w 7.9 Wenchuan earthquake, China.
48
49 Tectonophysics 471, 203-215.

50
51
52
53 Lin, A., Shin, J., and Kano, K., 2012. Fluidized Cataclastic Veins along the
54
55 Itoigawa-Shizuoka Tectonic Line Active Fault System, Central Japan, and Its
56
57

- Seismotectonic Implications. *The Journal of Geology* 120, 453-465.
- Lin, A., Toda, S., Rao, G., Tsuchihashi, S. and Yan, B., 2013a. Structural Analysis of Coseismic Normal Fault Zones of the 2011 M_w 6.6 Fukushima Earthquake, Northeast Japan. *Bulletin of the Seismological Society of America* 103, 1603-1613.
- Lin, A., Yamashita, K., and Tanaka, M., 2013b. Repeated seismic slips recorded in ultracataclastic veins along active faults of the Arima–Takatsuki Tectonic Line, southwest Japan. *Journal of Structural Geology* 48, 3-13.
- Liu, J., Zhang, P., Lease, R., Zheng, D., Wan, J., Wang, W., and Zhang, H., 2013. Eocene onset and late Miocene acceleration of Cenozoic intracontinental extension in the North Qinling range–Weihe graben: Insights from apatite fission track thermochronology. *Tectonophysics* 584, 281-296.
- Liu, M., Cui, X. and Liu, F., 2004. Cenozoic rifting and volcanism in eastern China: a mantle dynamic link to the Indo-Asian collision? *Tectonophysics* 393, 29-42.
- Ma, X. and Wu, D., 1987. Cenozoic extensional tectonics in China. *Tectonophysics* 133, 243-255.
- Ma, X., 1989. *Lithospheric Dynamics Atlas of China* (in Chinese). China Cartographic Publishing House, Beijing, 548 pp.
- McCalpin, J.P., 2009. *Paleoseismology*, Second Edition. International Geophysics Series, vol. 95, Academic Press 613 pp.
- Meng, Q. and Zhang, G., 2000. Geologic framework and tectonic evolution of the Qinling orogen, central China. *Tectonophysics* 323, 183-196.

- Mercier, J.L., Vergely, P., Zhang, Y.Q., Hou, M.J., Bellier, O., and Wang, Y.M., 2013. Structural records of the Late Cretaceous–Cenozoic extension in Eastern China and the kinematics of the Southern Tan-Lu and Qinling Fault Zone (Anhui and Shaanxi provinces, PR China): *Tectonophysics* 582, 50-75.
- Molnar, P. and Deng, Q., 1984. Faulting associated with large earthquakes and the average rate of deformation in central and eastern Asia. *Journal of Geophysical Research* 89, 6203-6227.
- Molnar, P. and Lyon-Caen, H., 1989. Fault plane solutions of earthquakes and active tectonics of the Tibetan Plateau and its margins. *Geophysical Journal International* 99, 123-154.
- Moro, M., Saroli, M., Gori, S., Falcucci, E., Galadini, F., and Messina, P., 2012. The interaction between active normal faulting and large scale gravitational mass movements revealed by paleoseismological techniques: A case study from central Italy. *Geomorphology* 151–152, 164-174.
- Myers, W.B., and Hamilton, W., 1964. Deformation accompanying the Hebgen Lake earthquake of August 17, 1959. U.S. Geological Survey Professional Paper 435, 55-98.
- Peltzer, G., Tapponnier, P., Zhitao, Z. and Qin, X.Z., 1985. Neogene and Quaternary faulting in and along the Qinling Shan. *Nature* 317, 500-505.
- Rao, G., Lin, A., Yan, B., Jia, D., Wu, X., and Ren, Z., 2011. Co-seismic Riedel shear structures produced by the 2010 M_w 6.9 Yushu earthquake, central Tibetan Plateau, China. *Tectonophysics* 507, 86-94.

- Ratschbacher, L., Hacker, B., Calvert, A., Webb, L., Grimmer, J., McWilliams, M., Ireland, T., Dong, S., and Hu, J., 2003. Tectonics of the Qinling (Central China): tectonostratigraphy, geochronology, and deformation history. *Tectonophysics* 366, 1-53.
- Shelton, J.W., 1984. Listric normal faults; an illustrated summary. *AAPG Bulletin* 68, 801-815.
- State Seismological Bureau (SSB), 1988. Active fault system around Ordos Massif (in Chinese). Seismological Press, Beijing, 352 pp.
- Steven, D.K., Beach, A., Brockbank, P.J., Brown, J.L., McCallum, J.E., Welbon, A.I., 1996. Spatial and mechanical controls on normal fault populations. Journal of Structural Geology 18, 359-372.
- Stewart, I.S. and Hancock, P.L., 1991. Scales of structural heterogeneity within neotectonic normal fault zones in the Aegean region. *Journal of Structural Geology* 13, 191-204.
- Stuiver, M., Reimer, P.J., Reimer, R., 2005. CALIB radiocarbon calibration version 7.0. <http://radiocarbon.pa.qub.ac.uk/calib/> (Last accessed, 10 October 2013).
- Sun, J., Johnson, K.M., Cao, Z., Shen, Z., Bürgmann, R., and Xu, X., 2011. Mechanical constraints on inversion of coseismic geodetic data for fault slip and geometry: Example from InSAR observation of the 6 October 2008 M_w 6.3 Dangxiong-Yangyi (Tibet) earthquake. *Journal of Geophysical Research*, 116(B1), B01406. <http://dx.doi.org/10.1029/2010JB007849> .
- Tapponnier, P. and Molnar, P., 1977. Active faulting and tectonics in China. *Journal of*

- Geophysical Research 82, 2905-2930.
- Tian, X., Teng, J., Zhang, H., Zhang, Z., Zhang, Y., Yang, H., and Zhang, K., 2011. Structure of crust and upper mantle beneath the Ordos Block and the Yinshan Mountains revealed by receiver function analysis. *Physics of the Earth and Planetary Interiors* 184, 186-193.
- Toda, S. and Tsutsumi, H., 2013. Simultaneous Reactivation of Two, Subparallel, Inland Normal Faults during the M_w 6.6 11 April 2011 Iwaki Earthquake Triggered by the M_w 9.0 Tohoku-Oki, Japan, Earthquake. *Bulletin of the Seismological Society of America* 103, 1584 -1602.
- Wallace, R.E., 1984. Fault scarps formed during the earthquake of October 2, 1915, in Pleasant Valley, Nevada, and some tectonic implications. U.S. Geological Survey Professional Paper 1274-A, 1-33.
- Wang, J., 1980. Ground ruptures during the large earthquake of 1556, Huaxian County, Shanxi. *Acta Seismologica Sinica* 2, 430-437 (in Chinese with English abstract).
- Wang, J., 1987. The Fenwei rift and its recent periodic activity. *Tectonophysics* 133, 257-275.
- Wang, Q., Zhang, P., Freymueller, J.T., Bilham, R., Larson, K.M., Lai, X., You, X., Niu, Z., Wu, J., Li, Y., Liu, J., Yang, Z., and Chen, Q., 2001. Present-Day Crustal Deformation in China Constrained by Global Positioning System Measurements. *Science* 294, 574-577.
- West, M.W., 1993. Extensional reactivation of thrust faults accompanied by coseismic surface rupture, southwestern Wyoming and north-central Utah. *Geological*

- 1 Society of America Bulletin 105, 1137-1150.
- 2
- 3 Xiao, H. and Suppe, J., 1992. Origin of rollover. AAPG Bulletin 76, 509-529.
- 4
- 5
- 6 Xie, Z., 2011. Structural Model of Earthquake Preparation in Weihe Extensional Basin.
- 7
- 8
- 9 Journal of Catastrophology 26, 18-21 (in Chinese with English abstract).
- 10
- 11 Xie, Y., 1992. On magnitude of 1556 Guanzhong great earthquake. Journal of
- 12
- 13
- 14 Catastrophology 7, 10-13 (in Chinese with English abstract)..
- 15
- 16
- 17 Ye, H., Zhang, B. and Mao, F., 1987. The Cenozoic tectonic evolution of the Great
- 18
- 19
- 20 North China: two types of rifting and crustal necking in the Great North China
- 21
- 22
- 23 and their tectonic implications. Tectonophysics 133, 217-227.
- 24
- 25
- 26 Yeats, R., Seih, K., Allen, C., 1997. The Geology of earthquakes. Oxford University
- 27
- 28
- 29 Press, Oxford, 568 pp.
- 30
- 31 Yuan, T. and Feng X., 2010, The 1556 Huaxian great earthquake (in Chinese),
- 32
- 33
- 34 Seismological Press, Beijing, 386 pp.
- 35
- 36
- 37 Yin, A. and Harrison, T.M., 2000. Geologic Evolution of the Himalayan-Tibetan
- 38
- 39
- 40 Orogen. Annual Review of Earth and Planetary Sciences 28, 211-280.
- 41
- 42
- 43 Zhang, A., Yang, Z., Zhong, J. and Mi, F., 1995. Characteristics of late quaternary
- 44
- 45
- 46 activity along the Southern Border Fault Zone of Weihe Graben Basin.
- 47
- 48
- 49 Quaternary International 25, 25-31.
- 50
- 51
- 52 Zhang, B., Jia, S., Wang, T. and Zheng, B., 1985. Intraplate seismotectonic features of
- 53
- 54
- 55 North China. Tectonophysics 117, 177-191.
- 56
- 57
- 58 Zhang, Y., Mercier, J.L. and Vergély, P., 1998. Extension in the graben systems around
- 59
- 60
- 61 the Ordos (China), and its contribution to the extrusion tectonics of south China
- 62
- 63
- 64
- 65

with respect to Gobi-Mongolia. Tectonophysics 285, 41-75.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure captions

Figure 1. (a) Location map of the Weihe Graben, showing the distribution of the major active faults and large historical earthquakes of the graben systems around the Ordos Block [modified from Deng (2007), focal mechanism data from Harvard University (2014)]. The inset map (b) shows the tectonic background. ATF, Altyn Tagh Fault; HYF, Haiyuan Fault; KLF, Kunlun Fault; XSHF, Xianshuihe Fault; SCB, South China Block; QLF, Qinling Fault; LMS, Longmenshan. (c) The color-shaded relief map shows the location and topographic characters of the study area. The red star indicates the epicenter of the 1556 M ~8.5 Huaxian earthquake (SSB, 1988; CENC, 2007). HPF, Huashan Piedmont Fault; NMF–WLT, North Margin Fault of the Weinan Loess Tableland; LPF, Lishan Piedmont Fault; QPF, Qinling Piedmont Fault; KZ–GSF, Kouzhen–Guanshan Fault.

Figure 2. Perspective view, looking towards the SE, of a 1 m WorldView image, reveals the active faults developed along the piedmont of the Huashan Mountains. The active fault zones are characterized by stepped normal fault scarps that are restricted to a narrow zone less than 500 m wide, whereas the lengths of individual fault traces are generally less than 200 m. The faults displace the alluvial fans, the ages of which are constrained to the Holocene by radiocarbon dating (Table 1), indicating the recent activity of the faults. The topographic profiles measured in the field are shown in Fig. 5.

Figure 3. (a) Map showing the distribution of active normal faults in the region

1 between Weinan and Huayin cities, the epicentral area of the 1556 M 8.5
2
3 Huaxian earthquake. The base map is a south perspective view of the
4
5 color-shaded relief map based on 30-m resolution ASTER GDEM data. (b–c)
6
7
8 Distribution of active fault traces mapped along the northern margin of the
9
10 Weinan Loess Tableland.
11
12
13

14 **Figure 4.** Map showing the distribution of active fault traces along the piedmont of
15
16 the Huashan Mountains, identified from the 0.5 m WorldView (a) and Google
17
18 Earth images (b–c).
19
20
21

22 **Figure 5.** (a–f) Topographic profiles across the active normal fault zones,
23
24 measured in the field using a laser rangefinder. By summing the individual
25
26 fault offsets, the cumulative vertical offset amounts for the whole zone of
27
28 active faulting are estimated, ranging from 15 to 80 m (generally 30–40 m).
29
30
31
32

33 **Figure 6.** (a) Stepped fault scarps developed along the Huashan piedmont (Loc. 1).
34
35 The person (circled) gives the scale. (b) The height of an individual scarp is as
36
37 much as ~19 m. (c–d) The main shear zone of the fault is composed mainly of
38
39 a fault breccia cataclasite, derived from the basement rocks, and disturbed
40
41 loess and alluvial deposits. The shear zone material is bounded by distinct
42
43 fault planes that dip NW at angles of 40°–54°. (e) Some brownish gray veins
44
45 composed of a fine-grained microbreccia were injected into both the loess and
46
47 the sandy gravel deposits.
48
49
50
51
52
53
54

55 **Figure 7.** (a) Close-up view of the 0.5-m resolution WorldView image of Loc. 2. (b)
56
57 The shear zone of the fault is well exposed in four parallel quarry pits (T1–T4)
58
59
60
61
62
63
64
65

that cut across the 10-m-high fault scarp along the piedmont of the Huashan Mountains. (c) Sketches of seven exposure walls clearly reveal fault planes that dip N at angles of 40°–71°, as well as other fault-related structures such as colluvial wedges and in-filled fissures, which are probably associated with paleoearthquakes. The ¹⁴C ages suggest the deformed landforms formed during the late Pleistocene–Holocene (see Table 1 for details).

Figure 8. (a–e) Field photographs show representative outcrops of the fault exposed on the trench walls at Loc. 2, where clear fault planes with high dip angles can be observed. The sand gravels and loess deposits were disturbed, and most were dragged into positions parallel or subparallel to the fault planes (d–e), and some were injected to form the wedge structures and fracture fillings (c). (a): T1-E; (b): T2-W; (c): T3-W; (d-e): T3-E.

Figure 9. (a) The shear zone of the fault observed at Loc. 3 (which is on the topographic boundary between basin and mountain, approximately 600 m east of Loc. 2). The fault plane dips at a high angle of 56°, and it separates unconsolidated alluvial deposits in the hanging wall from basement rocks in the footwall. (b) The shear zone of the fault is characterized by a narrow fault gouge and breccia zone less than 30 cm in thickness, in which most of the fault breccias were dragged into positions parallel or subparallel to the fault plane, suggesting a dominantly normal sense of slip, consistent with the observed fault steps and striations (c–e).

Figure 10. (a) 1-m resolution IKONOS image shows the fault scarps developed on

1 the terrace risers of the Chishuihe River along the northern margin of the
2
3 Weinan Loess Tableland. (b) The fault crops out on both banks of the
4
5 Chishuihe River, and the alluvial deposits of sand–pebble and cobble are
6
7 faulted against silts and sandstones. Fault gouges and striations were observed
8
9 on the fault plane. More details of the outcrop are shown in Fig. 11.
10
11
12
13

14 **Figure 11.** (a) Field photograph and corresponding sketch (b) of the fault outcrop at
15
16 Loc. 4. The pebbly deposits that bound the main fault plane were mostly
17
18 dragged into positions parallel or subparallel to the fault plane. (c) The
19
20 equal-area projection of measured striation data and topographic features
21
22 indicate an almost pure normal slip-sense at this site. The ^{14}C age dating
23
24 shows that the alluvial sediments in the hanging wall were deposited at
25
26 $18,270 \pm 60$ yr B.P. (Table 1), indicating recent activity on the fault since the
27
28 late Pleistocene.
29
30
31
32
33
34
35

36 **Figure 12.** Mode of tectonic deformation in the Weihe Graben. (a) The landscape of
37
38 the Weihe Graben is controlled by intracontinental normal faulting, resulting
39
40 in large amounts of subsidence and the accumulation of a thick pile of
41
42 sediments in the rift basin. Subsurface structures are modified from Wang
43
44 (1987) and SSB (1988). (b) Active normal faults in the study area formed in a
45
46 regime of ongoing extension that is probably related to the pre-existing
47
48 spreading and rifting of the continental crust in this area (in contrast to the
49
50 Ordos Block and other neighboring orogenic regions). The lithospheric
51
52 structures are schematic and based on the geophysical data of Bao et al.
53
54
55
56
57
58
59
60
61
62
63
64
65

(2011). The vertical depths are not precisely scaled.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1 Results of the ¹⁴C age dating

Sample code	Laboratory ID [*]	Material	¹⁴ C age (yr B.P.) [†]	Calendar year (2σ) [§]
C38	Beta-290421*	Shell	19,450 ± 80	BC 21,734-BC 21,153
C39	Beta-290422*	Shell	5,500 ± 40	BC 4,449-BC 4,319
C40	Beta-290423*	Shell	6,330 ± 40	BC 5,379-BC 5,217
C48	Beta-290425*	Shell	6,880 ± 40	BC 5,846-BC 5,671
Shell01	IAAA-100055**	Shell	14,050 ± 50	BC 15,386-BC 14,895
Shell11	IAAA-100065**	Shell	16,270 ± 50	BC 17,912-BC 17,529
C12	IAAA-120529**	Calcareous soil	18,270 ± 50	BC 20,386-BC 19,966

^{*} Samples were analyzed at the BETA Analysis Inc., USA.

^{**} Samples were analyzed at the Institute of Accelerator Analysis Ltd., Japan.

[†] Using Accelerator Mass Spectrometry (AMS) method, referenced to the year A.D. 1950.

[§] Dendrochronologically calibrated calendar age by Calibration Version 7.0 ([Stuiver et al., 2005](#)).

Table 2 Estimated fault dip-slip rates in Weihe Graben

Site	Vertical offset [*] (m)	Ages of landforms [#] (yr. BP)	Dip-slip rate (mm/yr)	Reference	Figure site
1	15.0	~5,500-6,880	~2.2-2.7	Alluvial fan	Figs. 2 and 5a
2	34.6	~14,050-16,270 [†]	2.1-2.5	Loess plateau surface	Fig. 5b
3	37	~14,050-16,270 [†]	2.3-2.6	Loess plateau surface	Figs. 2 and 5c
4	43	~14,050-16,270 [†]	2.6-3.1	Loess plateau surface	Figs. 2 and 5d
5	80	~14,050-16,270 [†]	4.9-5.7	Loess plateau surface	Fig. 5e
6	38	~14,050-16,270 [†]	2.3-2.7	Loess plateau surface	Fig. 5f

^{*} Cumulative vertical offset value for the whole fault zone.
[#] The ¹⁴C ages referenced to the year A.D. 1950 (see Table 1 for the details).
[†] The ages of regional loess plateau surfaces.

Figure1
[Click here to download high resolution image](#)




Figure-2
[Click here to download high resolution image](#)




Figure-3
[Click here to download high resolution image](#)




Figure-4
[Click here to download high resolution image](#)




Figure-5
[Click here to download high resolution image](#)




Figure-6
[Click here to download high resolution image](#)




Figure7
[Click here to download high resolution image](#)




Figure-8
[Click here to download high resolution image](#)



Figure-9
[Click here to download high resolution image](#)




Figure-10
[Click here to download high resolution image](#)



Figure-11
[Click here to download high resolution image](#)




Figure-12
[Click here to download high resolution image](#)

