

Weak dissipativity in derivative nonlinear Schrödinger equations

Hideaki Sunagawa
Osaka Metropolitan University

This short article is an announcement of the forthcoming paper [20] with Chunhua Li, Yoshinori Nishii and Yuji Sagawa, which concerns L^2 -decay properties of small solutions to a class of cubic derivative nonlinear Schrödinger equations in one space dimension. Throughout this article, we denote by $\mathcal{L}$ the standard free Schrödinger operator $i\partial_t + \frac{1}{2}\partial_x^2$ for $(t, x) \in \mathbb{R} \times \mathbb{R}$ with $i = \sqrt{-1}$. The function space H^k stands for the L^2 -based Sobolev space of order k equipped with the norm $\|\phi\|_{H^k} = \sum_{j \leq k} \|\partial_x^j \phi\|_{L^2}$, and the weighted Sobolev space $H^{k,m}$ is defined by $\{\phi \in L^2 \mid \langle \cdot \rangle^m \phi \in H^k\}$ with $\langle x \rangle = \sqrt{1+x^2}$.

1 Backgrounds

First of all, let us recall some of well-known results on large-time behavior of small data solutions to the cubic power-type nonlinear Schrödinger equation in the form

$$\mathcal{L}u = \lambda|u|^2u, \quad t > 0, \ x \in \mathbb{R}, \quad (1.1)$$

where λ is a constant. What is interesting in (1.1) is that the large-time behavior of the solution is actually affected by the coefficient λ even if the initial data is sufficiently small, smooth and decaying fast as $|x| \rightarrow \infty$. If $\lambda \in \mathbb{R}$, it is shown in [5] that the solution to (1.1) with small initial data behaves like

$$u(t, x) = \frac{1}{\sqrt{it}} \alpha(x/t) e^{i\{t \frac{x^2}{2t} - \lambda|\alpha(x/t)|^2 \log t\}} + o(t^{-1/2}) \quad \text{as } t \rightarrow \infty$$

with a suitable $\mathbb{C}$ -valued function $\alpha(y)$. An important consequence of this asymptotic expression is that the solution decays like $O(t^{-1/2})$ in $L^\infty(\mathbb{R}_x)$, while it does not behave like the free solution unless $\lambda = 0$. In other words, the additional logarithmic factor in the phase reflects the long-range character of the cubic nonlinear Schrödinger equations in one space dimension. If $\lambda \in \mathbb{C}$ in (1.1), another kind of long-range effect can be observed. For instance, according to [26] (see also [16], [9], [3], etc.), the small data solution $u(t, x)$ to (1.1) decays like $O(t^{-1/2}(\log t)^{-1/2})$ in $L^\infty(\mathbb{R}_x)$ as $t \rightarrow +\infty$ if $\text{Im } \lambda < 0$. This gain of additional logarithmic time decay should be interpreted as another kind of long-range effect (see also [1], [2], [3], [4], [6], [7], [8], [9], [10], [11], [13], [14], [16], [17], [18], [21], [22], [24], [25], and so on). Time decay in L^2 -norm is also investigated by several authors. Among others, it is pointed out by Kita-Sato [15] that the optimal L^2 -decay rate is $O((\log t)^{-1/2})$ in the case of (1.1) with $\text{Im } \lambda < 0$. We intend to extend this kind of L^2 -decay results to the case where the nonlinear term depends also on $\partial_x u$.

2 Derivative nonlinear Schrödinger equations

2.1 Weak dissipativity

From now on, we turn our attention to the initial value problem in the form

$$\mathcal{L}u = N(u, \partial_x u), \quad t > 0, \quad x \in \mathbb{R} \quad (2.1)$$

with

$$u(0, x) = \varphi(x), \quad x \in \mathbb{R}, \quad (2.2)$$

where φ is a prescribed $\mathbb{C}$ -valued function on $\mathbb{R}$. The nonlinear term $N(u, \partial_x u)$ is a cubic homogeneous polynomial in $(u, \bar{u}, \partial_x u, \bar{\partial}_x u)$ with complex coefficients. If φ is $O(\varepsilon)$ in $H^3 \cap H^{2,1}$ with $0 < \varepsilon \ll 1$, what we can expect for general cubic nonlinear Schrödinger equations in $\mathbb{R}$ is the lower estimate for the lifespan T_ε in the form $T_\varepsilon \geq \exp(c/\varepsilon^2)$ with some $c > 0$ not depending on ε , and this is best possible in general (see [12] for an example of small data blow-up). More precise information on the lifespan is available under the restriction

$$N(e^{i\theta}, 0) = e^{i\theta} N(1, 0), \quad \theta \in \mathbb{R} \quad (2.3)$$

and the initial condition

$$u(0, x) = \varepsilon \psi(x), \quad x \in \mathbb{R}, \quad (2.4)$$

instead of (2.2), where $\psi \in H^3 \cap H^{2,1}$ is independent of ε . In fact we have the following.

Theorem 2.1 ([23], [27]). *Assume that $\psi \in H^3 \cap H^{2,1}(\mathbb{R})$. Suppose that the nonlinear term N satisfies (2.3). Let T_ε be the supremum of $T > 0$ such that the initial value problem (2.1)–(2.4) admits a unique solution in $C([0, T]; H^3 \cap H^{2,1}(\mathbb{R}))$. Then it holds that*

$$\liminf_{\varepsilon \rightarrow +0} \varepsilon^2 \log T_\varepsilon \geq \frac{1}{2 \sup_{\xi \in \mathbb{R}} (|\hat{\psi}(\xi)|^2 \operatorname{Im} \nu(\xi))} \quad (2.5)$$

with the convention $1/0 = +\infty$, where the function $\nu : \mathbb{R} \rightarrow \mathbb{C}$ is defined by

$$\nu(\xi) = \frac{1}{2\pi i} \oint_{|z|=1} N(z, i\xi z) \frac{dz}{z^2}, \quad \xi \in \mathbb{R}, \quad (2.6)$$

and $\hat{\psi}$ denotes the Fourier transform of ψ , i.e.,

$$\hat{\psi}(\xi) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{-iy\xi} \psi(y) dy, \quad \xi \in \mathbb{R}.$$

In view of the right-hand side in (2.5), it may be natural to expect that the sign of $\operatorname{Im} \nu(\xi)$ has something to do with global behavior of small data solutions to (2.1). In fact, it has been pointed out in [23] that typical results on small data global existence and large-time asymptotic behavior for (2.1) under (2.3) can be summarized in terms of $\operatorname{Im} \nu(\xi)$ as follows:

- Small data global existence holds in $C([0, \infty); H^3 \cap H^{2,1})$ under the condition

$$\operatorname{Im} \nu(\xi) \leq 0, \quad \xi \in \mathbb{R}, \quad (\mathbf{A})$$

- The global solution has (at most) logarithmic phase correction if

$$\operatorname{Im} \nu(\xi) = 0, \quad \xi \in \mathbb{R}. \quad (\mathbf{A}_0)$$

Also it is not difficult to see that there is no L^2 -decay under $(\mathbf{A}_0)$ for generic initial data of small amplitude.

- L^2 -decay of the global solution occurs under the condition

$$\sup_{\xi \in \mathbb{R}} \operatorname{Im} \nu(\xi) < 0. \quad (\mathbf{A}_+)$$

Note that $\nu(\xi) = \lambda$ if $N = \lambda|u|^2u$. So these results cover the results in the power-type nonlinearity case mentioned in Section 1. However, it is pointed out in [19] that an interesting case is not covered by these classifications, that is the case where $(\mathbf{A})$ is satisfied but $(\mathbf{A}_0)$ and $(\mathbf{A}_+)$ are violated (for example, if $N = -i|u_x|^2u$, we can easily check that $\operatorname{Im} \nu(\xi) = -\xi^2 \leq 0$, while the inequality is not strict because of vanishing at $\xi = 0$). This is what we are interested in.

Before going further, let us remember the fact that, if $(\mathbf{A})$ is satisfied but $(\mathbf{A}_0)$ and $(\mathbf{A}_+)$ are violated, then there exist $c_0 > 0$ and $\xi_0 \in \mathbb{R}$ such that $\operatorname{Im} \nu(\xi) = -c_0(\xi - \xi_0)^2$. The converse is also true. This fact naturally leads us to the following definition of *the weak dissipativity*.

Definition 2.1. We say that a cubic nonlinear term N is *weakly dissipative* if the following two conditions (i) and (ii) are satisfied:

- (i) $N(e^{i\theta}, 0) = e^{i\theta} N(1, 0)$ for $\theta \in \mathbb{R}$.
- (ii) There exist $c_0 > 0$ and $\xi_0 \in \mathbb{R}$ such that $\operatorname{Im} \nu(\xi) = -c_0(\xi - \xi_0)^2$.

2.2 Upper and lower L^2 -decay bounds in the weakly dissipative case

The following two results are due to [20], which reveal the L^2 -decay property in the weakly dissipative case.

Theorem 2.2 ([20]). *Suppose that N is weakly dissipative and that $\varepsilon = \|\varphi\|_{H^3 \cap H^{2,1}}$ is sufficiently small. Then there exists a positive constant C , not depending on ε , such that the global solution u to (2.1)–(2.2) satisfies*

$$\|u(t)\|_{L_x^2} \leq \frac{C\varepsilon}{(1 + \varepsilon^2 \log(t+1))^{1/4}}$$

for $t \geq 0$.

Theorem 2.3 ([20]). *Suppose that N is weakly dissipative and that $\hat{\psi}$ does not vanish at the point ξ_0 coming from (ii) in Definition 2.1. Then we can choose $\varepsilon_0 > 0$ such that the global solution u to (2.1)–(2.4) satisfies*

$$\liminf_{t \rightarrow +\infty} ((\log t)^{1/4} \|u(t)\|_{L_x^2}) > 0$$

for $\varepsilon \in (0, \varepsilon_0]$.

Remark 2.1. According to [15], the optimal L^2 -decay rate is $O((\log t)^{-1/2})$ in the case where $N = \lambda|u|^2u$ with $\text{Im } \lambda < 0$. This should be contrasted with Theorems 2.2 and 2.3, because these tell us that the optimal L^2 -decay rate in the weakly dissipative case is $O((\log t)^{-1/4})$.

Now, let us explain heuristically why L^2 -decay rate should be $O((\log t)^{-1/4})$ if $\hat{\psi}(\xi_0) \neq 0$. For this purpose, let us first remember the fact that the solution u^0 to the free Schrödinger equation (i.e., the case of $N = 0$) with (2.2) behaves like

$$\partial_x^k u^0(t, x) \sim \left(\frac{ix}{t}\right)^k \frac{e^{-i\pi/4}}{\sqrt{t}} \hat{\varphi}\left(\frac{x}{t}\right) e^{i\frac{x^2}{2t}} + \dots$$

as $t \rightarrow +\infty$ for $k = 0, 1, 2, \dots$. Viewing it as a rough approximation of the solution u for (2.1), we may expect that $\partial_x^k u(t, x)$ could be better approximated by

$$\left(\frac{ix}{t}\right)^k \frac{1}{\sqrt{t}} A\left(\log t, \frac{x}{t}\right) e^{i\frac{x^2}{2t}}$$

with a suitable function $A(\tau, \xi)$, where $\tau = \log t$, $\xi = x/t$ and $t \gg 1$. Note that

$$A(0, \xi) = e^{-i\pi/4} \hat{\varphi}(\xi)$$

and that the extra variable $\tau = \log t$ is responsible for possible long-range nonlinear effect. Substituting the above expression into (2.1) and keeping only the leading terms, we can see (at least formally) that $A(\tau, \xi)$ should satisfy the ordinary differential equation

$$i\partial_\tau A = \nu(\xi)|A|^2 A + \dots$$

under (2.3). If N is weakly dissipative, we see that

$$\partial_\tau |A|^2 = -2c_0(\xi - \xi_0)^2 |A|^4 + \dots$$

Then it follows that

$$|A(\tau, \xi)|^2 = \frac{|\hat{\varphi}(\xi)|^2}{1 + 2c_0(\xi - \xi_0)^2 |\hat{\varphi}(\xi)|^2 \tau} + \dots,$$

whence

$$\|u(t)\|_{L_x^2} \sim \|A(\log t)\|_{L_\xi^2} \sim \left(\int_{\mathbb{R}} \frac{|\hat{\varphi}(\xi)|^2}{1 + 2c_0(\xi - \xi_0)^2 |\hat{\varphi}(\xi)|^2 \log t} d\xi \right)^{1/2} \quad (t \rightarrow +\infty)$$

up to harmless remainder. By considering the behavior as $t \rightarrow +\infty$ of this integral carefully, we see that L^2 -decay rate in the weakly dissipative case should be just $O((\log t)^{-1/4})$. Indeed, we have the following lemma.

Lemma 2.1. *Let $\theta \in L^\infty(\mathbb{R})$, $\xi_0 \in \mathbb{R}$ and*

$$S(\tau) = \int_{\mathbb{R}} \frac{|\theta(\xi)|^2}{1 + (\xi - \xi_0)^2 |\theta(\xi)|^2 \tau} d\xi, \quad \tau \geq 1. \quad (2.7)$$

(1) *We have*

$$S(\tau) \leq 4\|\theta\|_{L^\infty} \tau^{-1/2}, \quad \tau \geq 1.$$

(2) *Assume that there exists an open interval I with $I \ni \xi_0$ such that $\inf_{\xi \in I} |\theta(\xi)| > 0$. Then we can choose a positive constant C_* , which is independent of $\tau \geq 1$ (but may depend on θ and ξ_0), such that*

$$S(\tau) \geq C_* \tau^{-1/2}, \quad \tau \geq 1.$$

Our strategy of the proof of Theorems 2.2 and 2.3 is to justify the above heuristic argument. For the details, see the forthcoming paper [20].

Acknowledgments

The work is supported by JSPS KAKENHI (21K03314), and Osaka Central Advanced Mathematical Institute, Osaka Metropolitan University (MEXT Joint Usage/Research Center on Mathematics and Theoretical Physics JPMXP0619217849).

References

- [1] T. Cazenave and Z. Han, *Asymptotic behavior for a Schrödinger equation with nonlinear subcritical dissipation*, Discrete Contin. Dyn. Syst., **40** (2020), no.8, 4801–4819.
- [2] T. Cazenave, Z. Han and I. Naumkin, *Asymptotic behavior for a dissipative nonlinear Schrödinger equation*, Nonlinear Anal., **205** (2021), Paper No.112243, 37 pp.
- [3] N. Hayashi, C. Li and P. I. Naumkin, *Time decay for nonlinear dissipative Schrödinger equations in optical fields*, Adv. Math. Phys., **2016**, Article ID 3702738.
- [4] N. Hayashi, C. Li and P. I. Naumkin, *Upper and lower time decay bounds for solutions of dissipative nonlinear Schrödinger equations*. Commun. Pure Appl. Anal. **16** (2017), no.6, 2089–2104.
- [5] N. Hayashi and P. I. Naumkin, *Asymptotics for large time of solutions to the nonlinear Schrödinger and Hartree equations*, Amer. J. Math., **120** (1998), no.2, 369–389.
- [6] N. Hayashi, P. I. Naumkin and H. Sunagawa, *On the Schrödinger equation with dissipative nonlinearities of derivative type*, SIAM J. Math. Anal., **40** (2008), no.1, 278–291.

- [7] G. Hoshino, *Asymptotic behavior for solutions to the dissipative nonlinear Schrödinger equations with the fractional Sobolev space*, J. Math. Phys., **60** (2019), no.11, 111504, 11 pp.
- [8] G. Hoshino, *Dissipative nonlinear Schrödinger equations for large data in one space dimension*. Commun. Pure Appl. Anal. **19** (2020), no.2, 967–981.
- [9] G. Jin, Y. Jin and C. Li, *The initial value problem for nonlinear Schrödinger equations with a dissipative nonlinearity in one space dimension*, J. Evol. Equ., **16** (2016), no.4, 983–995.
- [10] S. Katayama, C. Li and H. Sunagawa, *A remark on decay rates of solutions for a system of quadratic nonlinear Schrödinger equations in 2D*, Differential Integral Equations **27** (2014), no.3–4, 301–312.
- [11] D. Kim, *A note on decay rates of solutions to a system of cubic nonlinear Schrödinger equations in one space dimension*, Asymptot. Anal., **98** (2016), no.1–2, 79–90.
- [12] N. Kita, *Existence of blowing-up solutions to some Schrödinger equations including nonlinear amplification with small initial data*, OCAMI Preprint Series 20-2 (2020).
- [13] N. Kita and C. Li, *Decay estimate of solutions to dissipative nonlinear Schrödinger equations*, OCAMI Preprint Series 20-5 (2020).
- [14] N. Kita and Y. Nakamura, *Decay estimate and asymptotic behavior of small solutions to Schrödinger equations with subcritical dissipative nonlinearity*, Adv. Stud. Pure Math., **81** (2019), 121–138.
- [15] N. Kita and T. Sato, *Optimal L^2 -decay of solutions to a cubic dissipative nonlinear Schrödinger equation*, to appear in Asymptotic Analysis.
- [16] N. Kita and A. Shimomura, *Large time behavior of solutions to Schrödinger equations with a dissipative nonlinearity for arbitrarily large initial data*, J. Math. Soc. Japan, **61** (2009), no.1, 39–64.
- [17] C. Li, Y. Nishii, Y. Sagawa and H. Sunagawa, *Large time asymptotics for a cubic nonlinear Schrödinger system in one space dimension*, Funkcialaj Ekvacioj, **64** (2021), no.3, 361–377.
- [18] C. Li, Y. Nishii, Y. Sagawa and H. Sunagawa, *Large time asymptotics for a cubic nonlinear Schrödinger system in one space dimension, II*, Tokyo J. Math., **44** (2021), no.2, 411–416.
- [19] C. Li, Y. Nishii, Y. Sagawa and H. Sunagawa, *On the derivative nonlinear Schrödinger equation with weakly dissipative structure*, J. Evol. Equ., **21** (2021), 1541–1550.
- [20] C. Li, Y. Nishii, Y. Sagawa and H. Sunagawa, *Upper and lower L^2 -decay bounds for a class of derivative nonlinear Schrödinger equations*, to appear in Discrete Contin. Dyn. Syst. [arXiv:2204.07320].

- [21] C. Li and H. Sunagawa, *On Schrödinger systems with cubic dissipative nonlinearities of derivative type*, Nonlinearity, **29** (2016), no.5, 1537–1563; Corrigendum, *ibid.*, no.12, C1–C2.
- [22] T. Ogawa and T. Sato, *L^2 -decay rate for the critical nonlinear Schrödinger equation with a small smooth data*. NoDEA Nonlinear Differential Equations Appl. **27** (2020), no. 2, Paper No.18, 20 pp.
- [23] Y. Sagawa and H. Sunagawa, *The lifespan of small solutions to cubic derivative nonlinear Schrödinger equations in one space dimension*, Discrete Contin. Dyn. Syst., **36** (2016), no.10, 5743–5761; Corrigendum, *ibid.*, **40** (2020), no.7, 4577–4578.
- [24] T. Sato, *L^2 -decay estimate for the dissipative nonlinear Schrödinger equation in the Gevrey class*. Arch. Math. (Basel) **115** (2020), no.5, 575–588.
- [25] T. Sato, *Lower bound estimate for the dissipative nonlinear Schrödinger equation*, Partial Differ. Equ. Appl., **2** (2021), no.5, 66.
- [26] A. Shimomura, *Asymptotic behavior of solutions for Schrödinger equations with dissipative nonlinearities*, Comm. Partial Differential Equations, **31** (2006), no.7–9, 1407–1423.
- [27] H. Sunagawa, *Lower bounds of the lifespan of small data solutions to the nonlinear Schrödinger equations*, Osaka J. Math., **43** (2006), no.4, 771–789.

Department of Mathematics
 Graduate School of Science
 Osaka Metropolitan University
 Sumiyoshi-ku, Osaka 558-8585
 JAPAN
 E-mail address: sunagawa@omu.ac.jp