

Editorial Manager(tm) for Japanese Journal of Radiology
Manuscript Draft

Manuscript Number: RMED-273R3

Title: Influence of manipulating hypoxia in solid tumors on radiation dose-rate effect in vivo, referring to that in quiescent cell population

Article Type: Original Article

Keywords: Dose-rate effect; Manipulating hypoxia; Quiescent cell; Carbon-ion beams; Gamma-rays

Corresponding Author: Dr. Shin-ichiro Masunaga, M.D., Ph.D.

Corresponding Author's Institution: Research Reactor Institute, Kyoto University

First Author: Shin-ichiro Masunaga, M.D., Ph.D.

Order of Authors: Shin-ichiro Masunaga, M.D., Ph.D.; Ryoichi Hirayama, Ph.D.; Akiko Uzawa, Ph.D.; Genro Kashino, Ph.D.; Takushi Takata, Ph.D.; Hiroki Tanaka, Ph.D.; Minoru Suzuki, M.D., Ph.D.; Yuko Kinashi, M.D., Ph.D.; Yong Liu, M.D., Ph.D.; Sachiko Koike, Ph.D.; Koichi Ando, Ph.D.; Koji Ono, M.D., Ph.D.

Influence of manipulating hypoxia in solid tumors on radiation dose-rate effect *in vivo*, referring to that in quiescent cell population

Shin-ichiro Masunaga¹, Ryoichi Hirayama², Akiko Uzawa², Genro Kashino¹, Takushi Takata³, Hiroki Tanaka⁴, Minoru Suzuki¹, Yuko Kinashi⁵, Yong Liu¹, Sachiko Koike², Koichi Ando⁶ and Koji Ono¹

¹Particle Radiation Oncology Research Center, ⁴Radiation Medical Physics and ⁵Radiation Safety and Control, Research Reactor Institute, Kyoto University, Kumatori, Osaka, Japan.

²Heavy-Ion Radiobiology Research Group, Research Center for Charged Particle Therapy, National Institute of Radiological Sciences, Chiba, Japan.

³Wakasawan Energy Research Center, Tsuruga, Fukui, Japan.

⁶Heavy Ion Medical Center, Gunma University, Gunma, Japan.

Running head:

Influence of manipulating hypoxia on radiation dose-rate effect

All correspondence to:

Shin-ichiro Masunaga, M.D., Ph.D.

Particle Radiation Oncology Research Center

Research Reactor Institute, Kyoto University

2-1010, Asashiro-nishi, Kumatori-cho,

Sennan-gun, Osaka 590-0494, Japan.

Tel: +81-72-451-2406, 2487, Fax: +81-72-451-2627

E-mail: smasuna@rri.kyoto-u.ac.jp

Acknowledgments

This study was supported, in part, by a Grant-in-aid for Scientific Research (C) (20591493) from the Japan Society for the Promotion of Science.

Original article

The authors declare no conflicts of interest concerning this study.

Abstract

Purpose: To clarify the effect of manipulating intratumor hypoxia on radiosensitivity under reduced dose-rate (RDR) irradiation.

Methods: Tumor-bearing mice were continuously given 5-bromo-2'-deoxyuridine (BrdU) to label all proliferating (P) cells. They received γ -rays or accelerated carbon-ion beams at high dose-rate (HDR) or RDR with or without tumor clamping to induce hypoxia. Some mice without clamping received nicotinamide, an acute hypoxia-releasing agent or misonidazole, a hypoxic cell radio-sensitizer before irradiation. The responses of quiescent (Q) and total (= P+Q) cells were assessed by the micronucleus frequency using immunofluorescence staining for BrdU.

Results: The clearer decrease in radiosensitivity in Q than total cells after RDR γ -ray irradiation was suppressed with carbon-ion beams, especially with a higher linear energy transfer value. Repressing the decrease in the radiosensitivity under RDR irradiation through keeping tumors hypoxic during irradiation and enhancing the decrease in the radiosensitivity by nicotinamide were clearer with γ -rays and in total cells than with carbon-ion beams and in Q cells, respectively. Inhibiting the decrease in the radiosensitivity by misonidazole was clearer with γ -rays and in Q cells than with carbon-ion beams and in total cells, respectively.

Conclusion: Manipulating hypoxia during RDR as well as HDR irradiation influences tumor radiosensitivity, especially with γ -rays.

Key words:

Dose-rate effect; Manipulating hypoxia; Quiescent cell; Carbon-ion beams; γ -Rays

Introduction

Intensity modulated radiotherapy (IMRT) and stereotactic irradiation have become common as a new radiotherapy technique for treatment of malignancies. Both modalities generally use multiple arc or fixed-portal radiation beams, and radiation beams are exposed intermittently. These techniques often require 30 min or a longer time in one treatment session for precise positioning of patients.^{1,2} Prolongation of irradiation time may reduce a radiation effect, and evokes a major concern for the dose rate effect. Thus, it is needed to clarify the effect of the reduction of dose rate on the radiosensitivity of tumors *in vivo*.

When using low linear energy transfer (LET) radiation, lowering the dose rate is thought to reduce late effects in normal tissue much more than it decreases tumor control. Thus, the "therapeutic ratio" increases as the dose rate decreases, because the therapeutic ratio is equal to the ratio of tumor control to normal tissue complications. Further, the difference between early and late effects for low dose-rate radiotherapy, as well as improving the therapeutic ratio, allows complete treatment in a short period of time, minimizing the effects of tumor repopulation. In other words, decreasing the dose rate increases the therapeutic ratio, limited only by tumor cell repopulation.³ This is the primary rationale for low dose-rate radiotherapy using low LET radiation. High LET radiation is more effective than low LET X- or γ -radiation at inducing biologic damage. High LET radiation results in a greater relative biological effectiveness (RBE) value for cell killing, a reduced oxygen effect, and a reduced dependence on the cell cycle and the irradiation dose

1 rate.⁴

2
3 Manipulating hypoxia in solid tumors during irradiation apparently
4 influences tumor radiosensitivity under high dose-rate (HDR)
5 irradiation using low LET radiation.⁵ However, its significance in solid
6 tumors irradiated at a reduced dose rate (RDR) is less clear *in vivo*,
7 whether low or high LET radiation is employed.
8
9

10 Many cells in solid tumors are quiescent *in situ* but still
11 clonogenic.⁶ The quiescent (Q) tumor cells are more resistant to low-LET
12 radiation because of their larger hypoxic fraction and greater capacity
13 to recover from potentially lethal damage (PLD) than proliferating (P)
14 tumor cells. The rationale for low dose-rate radiotherapy does not take
15 into account the response of Q tumor cells at all.
16
17

18 Thus, in this study, we tried to elucidate the effect of
19 manipulating hypoxia in irradiated solid tumors at a RDR with low-LET
20 γ -rays or high-LET 290 MeV/u accelerated carbon-ion beams *in vivo*,
21 compared with HDR irradiation. Further, the responses of the total (=
22 P + Q) and Q cell populations in irradiated solid tumors were separately
23 detected with the method for selectively detecting the response of Q
24 cells within solid tumors.⁷ This is the first attempt to clarify the
25 direct relationship between the irradiation dose rate effect and the
26 oxygen effect *in vivo*, referring to the response of the Q cell population
27 in irradiated solid tumors.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Materials and Methods

Mice and tumors

SCC VII squamous cell carcinoma cell line derived from C3H/He mice was maintained *in vitro* in Eagle's minimum essential medium supplemented with 12.5 % fetal bovine serum. The tumor cells (1.0×10^5) were inoculated subcutaneously into the left hind leg of 9-week-old syngeneic female C3H/He mice (Japan Animal Co., Ltd., Osaka, Japan). Fourteen days later, the tumors, approximately 1 cm in diameter, were employed for experimental treatment, and the body weight of the tumor-bearing mice was 22.1 ± 2.3 (Mean $\pm$ SD) g. Mice were handled according to the Recommendations for Handling of Laboratory Animals for Biomedical Research, compiled by the Committee on Safety Handling Regulations for Laboratory Animal Experiments. Incidentally, the p53 of SCC VII tumor cells is the wild type.'

Labeling with 5-bromo-2'-deoxyuridine (BrdU)

Nine days after the inoculation, mini-osmotic pumps (Durect Corporation, Cupertino, CA) containing BrdU dissolved in physiological saline (250 mg/ml) were implanted subcutaneously to label all P cells for 5 days. The percentage of labeled cells after continuous labeling with BrdU was 55.3 ± 4.5 %, and reached a plateau at this stage. Therefore, tumor cells not incorporating BrdU after continuous labeling were regarded as Q cells.

Treatment

After the labeling with BrdU, tumor-bearing mice received γ -ray or accelerated carbon-ion whole-body irradiation, with the animal held in a specially designed device made of acrylic resin with the tail or all four legs firmly fixed with adhesive tape with no anesthetic. Some tumors were made totally hypoxic by clamping the proximal end 15 min

1 before irradiation.⁷ This clamping for 15 min did not influence
2
3 clonogenic cell survival or the level of micronucleation.

4
5 γ -Rays were delivered with a cobalt-60 γ -ray irradiator at a dose
6
7 rate of 2.5 or 0.039 Gy/min.

8
9 Carbon-12 ions were accelerated up to 290 MeV/u by the synchrotron
10
11 of the Heavy Ion Medical Accelerator installed at National Institute
12
13 of Radiological Sciences in Chiba, Japan. The dose rate was regulated
14
15 through a beam attenuation system, and irradiation was conducted using
16
17 horizontal carbon beams with a dose rate of 1.0 or 0.035 Gy/min. The
18
19 LET of the carbon ion beam with the 6-cm spread-out Bragg peak (SOBP)
20
21 ranges from 14 keV/ μ m to greater than 200 keV/ μ m, depending on depth.
22
23 A desired LET beam was obtained by selecting the depth along the beam
24
25 path using a Lucite range shifter. An LET of 18 and 50 keV/ μ m at the
26
27 middle of the plateau and the SOBP were employed here, respectively.⁸
28
29

30
31 Irradiated tumor-bearing mice were divided into 4 groups. I) Tumors
32
33 were excised immediately after irradiation only under aerated
34
35 conditions. II) Tumors were kept totally hypoxic during irradiation,
36
37 then excised immediately after irradiation. III) The tumor-bearing mice
38
39 received an intraperitoneal administration of an acute
40
41 hypoxia-releasing agent, nicotinamide (1000 mg/kg of mouse weight)
42
43 dissolved in physiological saline 60 min before irradiation, then the
44
45 tumors were excised immediately after irradiation under aerated
46
47 conditions. IV) The tumor-bearing mice received an intraperitoneal
48
49 administration of a hypoxic cell radio-sensitizer, misonidazole (1000
50
51 mg/kg of mouse weight) dissolved in physiological saline 30 min before
52
53 irradiation, then the tumors were excised immediately after irradiation
54
55 under aerated conditions. All the doses employed, sequences and timing
56
57 for nicotinamide, misonidazole and irradiation were appropriate enough
58
59
60
61
62
63
64
65

1 to function completely^{9,10}.

2 Each treatment group also included mice not pretreated with BrdU.

3
4 ***Immunofluorescence staining of BrdU-labeled cells and micronucleus (MN)***
5
6
7 ***assay***
8

9 Tumors excised from the mice given BrdU were minced and trypsinized
10 [0.05% trypsin and 0.02% ethylenediamine-tetraacetic acid (EDTA) in
11 phosphate-buffered saline (PBS), 37 °C, 20 min]. Tumor cell suspensions
12 were incubated for 72 h in tissue culture dishes containing complete
13 medium and 1.0 µg/ml of cytochalasin-B to inhibit cytokinesis while
14 allowing nuclear division, and the cultures were then trypsinized and
15 cell suspensions were fixed. After the centrifugation of fixed cell
16 suspensions, the cell pellet was resuspended with cold Carnoy's fixative
17 (ethanol:acetic acid = 3:1 in volume). The suspension was placed on
18 a glass microscope slide and the sample was dried at room temperature.
19 The slides were treated with 2 M hydrochloric acid for 60 min at room
20 temperature to dissociate the histones and partially denature the DNA.
21 The slides were immersed in borax-borate buffer (pH 8.5) to neutralize
22 the acid. BrdU-labeled tumor cells were detected by indirect
23 immunofluorescence staining using a monoclonal anti-BrdU antibody
24 (Becton Dickinson, San Jose, CA) and a fluorescein isothiocyanate
25 (FITC)-conjugated antimouse IgG antibody (Sigma, St. Louis, MO). To
26 observe the double staining of tumor cells with green-emitting FITC
27 and red-emitting propidium iodide (PI), cells on the slides were treated
28 with PI (2 µg/ml in PBS) and monitored under a fluorescence microscope.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52

53 The MN frequency in cells not labeled with BrdU could be examined
54 by counting the micronuclei in the binuclear cells that showed only
55 red fluorescence. The MN frequency was defined as the ratio of the number
56 of micronuclei in the binuclear cells to the total number of binuclear
57
58
59
60
61
62
63
64
65

cells observed.⁷ The MN frequency has already been shown to be a tool for detecting radiosensitivity to carbon-ion beams.¹¹

The ratios obtained in tumors not pretreated with BrdU indicated the MN frequency at all phases in the total tumor cell population. More than 400 binuclear cells were counted to determine the MN frequency.

Clonogenic cell survival assay

The clonogenic cell survival assay was also performed in the mice given no BrdU using an *in vivo-in vitro* assay method. Tumors were disaggregated by stirring for 20 min at 37 °C in PBS containing 0.05 % trypsin and 0.02% EDTA. The cell yield was $(4.5 \pm 1.1) \times 10^7$ /g tumor weight.

As stated above, the MN frequencies for Q cells were obtained from non-labeled tumor cells after continuous BrdU labeling. The MN frequencies and surviving fractions for the total cell population were obtained from cells in tumors not pretreated with BrdU. Thus, there was no effect of interaction between BrdU and irradiation on the values of MN frequency and SF. More than 3 tumor-bearing mice were used to assess each set of conditions and each experiment was repeated at least twice. To examine the differences between pairs of values, Student's *t*-test was used when variances of the two groups could be assumed to be equal; otherwise the Welch *t*-test was used.

Results

The plating efficiency and MN frequency at 0 Gy are shown in **Table 1**. The plating efficiency was significantly smaller for the combination with MISO than for any other condition in the total cell population. In both the total and Q cell populations, the MN frequency significantly increased in the following order: absolute control < totally hypoxic conditions < combination with nicotinamide < combination with misonidazole. Further, the MN frequency was significantly higher for Q cell population than the total cell population under all conditions.

Cell survival curves for the total tumor cells as a function of radiation dose after γ -ray irradiation are shown in **Figure 1**. For both HDR and RDR irradiation, the surviving fractions (SFs) decreased in the following order: irradiation under totally hypoxic conditions > aerobic irradiation without any drug > irradiation after nicotinamide loading > irradiation after misonidazole loading. The SFs under all conditions increased as the dose rate of radiation decreased, especially after nicotinamide loading.

Cell survival curves for the total tumor cells as a function of radiation dose after accelerated carbon-ion beam irradiation with an LET of 18 and 50 keV/ μ m are shown in the left and right panels of **Figure 2**, respectively. For both HDR and RDR irradiation, the SFs decreased in the same order as for γ -ray irradiation, but the degree of change was reduced, especially with 50 keV/ μ m carbon-ion beams. The increases in the SF with the decrease in dose rate were suppressed compared with γ -ray irradiation, again especially with 50 keV/ μ m carbon-ion beams.

For baseline correction, we used the normalized MN frequency to exclude the MN frequency in non-irradiated tumors. The normalized MN

frequency was the MN frequency in the irradiated tumors minus that in the non-irradiated tumors. Dose response curves of the normalized MN frequency for total and Q tumor cell populations as a function of radiation dose after γ -ray irradiation are shown in the left and right panels of **Figure 3**, respectively. Overall, the normalized MN frequencies were significantly smaller in Q cells than the total cells. In both total and Q cells, under RDR as well as HDR irradiation, the normalized MN frequencies increased in the following order: irradiation under totally hypoxic conditions < aerobic irradiation without drugs < irradiation after nicotinamide loading < irradiation after misonidazole loading. The normalized MN frequencies decreased with the decrease in dose rate under all conditions, especially irradiation after nicotinamide loading in the total cell population.

Dose response curves of the normalized MN frequency for the total and Q tumor cells as a function of radiation dose after accelerated carbon-ion beam irradiation with an LET of 18 (**Figure 4**) or 50 keV/ μ m (**Figure 5**) are shown in the left and right panels, respectively. Overall, the normalized MN frequencies were significantly smaller in Q cells than total cells, but the differences in radiosensitivity between the total and Q cells were reduced compared with γ -ray irradiation, especially with 50 keV/ μ m carbon-ion beams. In both the total and Q cells, under RDR as well as HDR irradiation, the normalized MN frequencies increased in the same order as for γ -ray irradiation, but the degree of change was reduced, especially with 50 keV/ μ m carbon-ion beams. The decreases in the normalized MN frequency with the decrease in radiation dose rate were suppressed compared with γ -ray irradiation, again especially with 50 keV/ μ m carbon-ion beams.

To estimate the effect of aerobic irradiation compared with hypoxic irradiation in both the total and Q cells, the data for aerobic irradiation without any drug and irradiation under totally hypoxic conditions were used (**Table 2**). Following γ -ray irradiation, the values were significantly smaller for Q cells than the total cells ($P < 0.05$), and in both the total and Q cells, the values were significantly smaller for RDR than HDR irradiation ($P < 0.05$). In both populations, carbon-ion beams produced significantly smaller values than γ -rays ($P < 0.05$), and the values approached to 1.0 as the LET values increased.

To assess the radio-enhancing effect of nicotinamide or misonidazole under aerobic conditions in both the total and Q cells compared with aerobic irradiation without any drug, the data for aerobic irradiation with and without drugs were used (**Table 3**). The enhancing effect of nicotinamide was more marked in the total cell population, especially with HDR γ -ray irradiation. The effect was little observed for RDR γ -ray irradiation and accelerated carbon-ion irradiation especially with a higher LET value. In contrast, the enhancing effect of misonidazole was more marked in the Q cells. The enhancing effect was also attenuated for accelerated carbon-ion irradiation especially with a higher LET value.

To investigate the reduction in radiosensitivity caused by a decrease in radiation dose rate, dose-modifying factors were calculated using the data for all irradiation conditions given in **Figures 1** through **5** (**Table 4**). On the whole, the reduction in radiosensitivity was more marked in Q than the total cells, especially under γ -ray irradiation. In the total cells, the degree of the reduction of radiosensitivity was reduced in the following order: aerobic irradiation with nicotinamide

1 > aerobic irradiation without any drug > aerobic irradiation with
 2
 3 misonidazole > irradiation under totally hypoxic conditions. In Q cells,
 4
 5 the degree of the reduction of radiosensitivity was reduced in the
 6
 7 following order: aerobic irradiation with nicotinamide = aerobic
 8
 9 irradiation without any drug > irradiation under totally hypoxic
 10
 11 conditions > aerobic irradiation with misonidazole. This order of the
 12
 13 reduction in radiosensitivity and the difference in the reduction in
 14
 15 radiosensitivity between Q and the total cells became more indistinct
 16
 17 with the use of accelerated carbon-ion beams, especially at a higher
 18
 19 LET value, than with the use of γ -rays.
 20
 21
 22

23
 24 To examine the difference in radiosensitivity between the total
 25
 26 and Q cells, dose-modifying factors, which allow us to compare the dose
 27
 28 of radiation necessary to obtain a normalized MN frequency of 0.2 in
 29
 30 Q cells with that in the total cells, were calculated using the data
 31
 32 in **Figures 2 and 3 (Table 5)**. Overall, the difference in radiosensitivity
 33
 34 was greater under RDR than HDR irradiation, especially with γ -rays. The
 35
 36 difference in radiosensitivity increased in the following order:
 37
 38 irradiation under totally hypoxic conditions < aerobic irradiation with
 39
 40 misonidazole < aerobic irradiation without any drug $\leq$ aerobic
 41
 42 irradiation with nicotinamide. This order of the increase in the
 43
 44 difference in radiosensitivity and the difference itself in
 45
 46 radiosensitivity became more indistinct with the use of accelerated
 47
 48 carbon-ion beams, especially at a higher LET value, than with the use
 49
 50 of γ -rays.
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65

Discussion

Solid tumors, especially human tumors, are thought to contain a high proportion of Q cells.⁶ The presence of Q cell is probably due, in part, to hypoxia and the depletion of nutrients in the tumor core, another consequence of poor vascular supply.⁶ This might promote the formation of micronuclei at 0 Gy in Q tumor cells (**Table 1**). Q cells were shown to have significantly less radiosensitivity than the total cells here (**Figs. 3 through 5**). This means that more Q cells survive radiation therapy than P cells. Thus, the control of Q cells has a great impact on the outcome of radiation therapy. In both the total and Q cell populations, carbon ion irradiation was less dependent on oxygenation status with little recovery from radiation-induced DNA damage, leading to high RBE values compared with γ -ray irradiation.¹¹ In terms of the tumor cell-killing effect as a whole, including intratumor Q cell control, carbon-ion beam radiotherapy can be a promising treatment for refractory tumors.

Q cell population had been shown to include a significantly larger hypoxic fraction than the total cell population,¹² resulting in a significantly smaller effect of carbon-ion beams under aerobic irradiation compared with hypoxic irradiation in Q cells (**Table 2**). Since tumor radiosensitivity was significantly less dependent on intratumor oxygenation status and the irradiation dose rate effect when carbon-ion beams, especially with a higher LET value, were used, a much smaller effect of aerobic irradiation compared with hypoxic irradiation was observed here. This was partly because the frequency of closely spaced DNA lesions forming a cluster of DNA damage produced by high LET carbon-ion irradiation is much less dependent on oxygenation status

1 at the time of irradiation than that of DNA damage produced by low LET
2
3 γ -ray irradiation.¹³
4

5 Tumor hypoxia is a direct consequence of structural abnormalities
6 of the microvasculature and functional abnormalities of the
7 microcirculation in solid tumors and results from either limited oxygen
8 diffusion (chronic hypoxia) or limited perfusion (acute hypoxia,
9 transient hypoxia, or ischemic hypoxia). Large intercapillary distances
10 resulting from rapid tumor cell proliferation lead to chronically
11 hypoxic cells existing at the rim of the oxygen diffusion distance.¹⁴
12 Factors such as vessel plugging by blood cells or circulating tumor
13 cells, the collapse of vessels in regions of high tumor interstitial
14 pressure, or spontaneous vasomotor activity in normal tissue vessels
15 incorporated into the tumor which subsequently affects flow in
16 downstream tumor microvessels cause intermittent blood flow in tumors,
17 which results in acute hypoxia.¹⁵ Thus, acute hypoxic areas are
18 distributed throughout the tumor depending on these causative factors
19 and can occur sporadically in large areas of a solid tumor. Nicotinamide,
20 a vitamin B₃ analogue, is known to prevent these transient fluctuations
21 in tumor blood flow that lead to the development of acute hypoxia.⁹
22 Misonidazole is a typical 2-nitro-imidazole hypoxic cell sensitizer
23 which is thought to function as an oxygen-mimicking agent in intratumor
24 hypoxic areas under irradiation.¹⁰ In SCC VII tumors, it had been shown
25 that the hypoxic fraction of the total cell population is predominantly
26 made up of acute hypoxic areas and that of the hypoxia rich-Q cell
27 population is mainly made up of chronic hypoxic areas.¹² Therefore, under
28 HDR γ -ray irradiation, the enhancement ratio of nicotinamide was higher
29 in the total cells than in Q cells, and that of misonidazole was higher
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

in the Q cells (**Table 3**). When carbon-ion beams, especially with a higher LET value, were employed, these differences in the enhancement ratio were reduced because radiosensitivity was much less dependent on intratumor oxygenation status.¹¹ However, under RDR γ -ray irradiation, acute hypoxic areas appear and disappear throughout a solid tumor during long periods of irradiation.¹⁵ As a result, RDR irradiation even without nicotinamide could make it possible to irradiate all acute hypoxic areas under oxic conditions, leading to no radio-enhancing effect of nicotinamide. The radio-enhancing effect of misonidazole, which depends on the size of the hypoxic fraction in solid tumors, however, still could be observed under RDR as well as HDR irradiation.

Enhancement of the irradiation dose rate effect on the normalized frequency of micronuclei by γ -ray irradiation in the presence of nicotinamide compared with γ -ray irradiation alone was observed in the acute hypoxia-rich total cells rather than the chronic hypoxia-rich Q cells (**Table 4**). Suppression of the dose rate effect by inducing total hypoxia during irradiation was slightly more clearly observed in normoxia-rich total cell than hypoxia-rich Q cells. Some recent studies *in vitro* found that hypoxia-induced translational repression can explain the decreased homologous recombination (HR) repair of radiation-induced DNA double-stranded breaks (dsbs),¹⁶ a more important mechanism for the repair of dsbs in late-S and G2, and that HR plays a greater role in determining hypoxic radiosensitivity than normoxic radiosensitivity.¹⁷ Thus, the use of nicotinamide or tumor clamping to induce total hypoxia influenced repair more in the total cells including late-S and G2 phase cells than in Q cells. In contrast, the repression of the dose rate effect by the hypoxic cell radio-sensitizer

misonidazole was slightly more clearly observed in the hypoxia-rich Q cells than normoxia-rich total cells. It had already been shown that the loading of misonidazole after low LET HDR irradiation to solid tumors inhibited the recovery from radiation-induced PLD *in vivo*, especially in Q cells.¹⁸ In this study, it was shown that misonidazole combined with low LET RDR irradiation could suppress the dose rate effect, especially in Q cells. According to the previous finding that the recovery from PLD and the decrease in radiosensitivity through a reduction in the irradiation dose rate under low LET irradiation are mainly due to non-homologous end-joining (NHEJ) repair, which is the predominant DNA dsbs repair process for cells in G₀, G₁ or early-S phase,¹⁹ misonidazole itself or protein adducts of reductively-activated misonidazole in hypoxic areas in irradiated tumors may inhibit NHEJ more efficiently than HR.¹⁰ Again, when carbon-ion beams, especially with a higher LET value, were employed, the effects of combined treatment on the irradiation dose rate effect became indistinct because radiosensitivity was much less dependent on intratumor oxygenation status and the irradiation dose rate.¹¹

Overall, the difference in radiosensitivity between the total and Q cells was increased by reducing the dose rate, especially for γ -ray irradiation, because of the greater reduction in radiosensitivity caused by a decrease in the dose rate in Q cells than in the total cells⁹ (Table 5). Nicotinamide enhanced the radiosensitivity of the total cells, leading to a widening of the difference in radiosensitivity between the total and Q cell populations compared with aerobic HDR γ -ray irradiation without any drug. But, for RDR γ -ray irradiation, the effect of nicotinamide disappeared due to the characteristics of acute hypoxia

1 in solid tumors, resulting in no change in the difference in
2 radiosensitivity between the total and Q cells compared with γ -ray
3 irradiation only. Meanwhile, under totally hypoxic conditions, the
4 difference in radiosensitivity was smaller than for aerobic γ -ray
5 irradiation because of the radio-resistance induced by total hypoxia
6 in both the total and Q cell populations in solid tumors. Misonidazole
7 enhanced the radiosensitivity of the hypoxia-rich Q cells much more
8 than that of the total cells under both HDR and RDR γ -ray irradiation,
9 leading to a decrease in the difference in radiosensitivity between
10 the two cells compared with aerobic γ -ray irradiation only. When
11 carbon-ion beams, especially with a higher LET value, were employed,
12 the effects of combined treatment on the difference in radiosensitivity
13 between the total and Q cells became indistinct because tumor
14 radiosensitivity was much less dependent on intratumor oxygenation
15 status and the irradiation dose rate.¹¹ Anyway, at least in this study,
16 it was elucidated that manipulating hypoxia during RDR irradiation,
17 especially with γ -rays, influences tumor radiosensitivity as well as
18 HDR irradiation in both total and Q cell populations. In radiotherapy,
19 irradiation dose rate also should be taken into account when intratumor
20 hypoxia is manipulated.

References

1. Ahmed RS, Kim RY, Duan J, Meleth S, De Los Santos JF, Fiveash JB. IMRT dose escalation for positive para-aortic lymph nodes in patients with locally advanced cervical cancer while reducing dose to bone marrow and other organs at risk. *Int J Radiat Oncol Biol Phys* 2005;60:505-512.
2. Wulf J, Haedinger U, Oppitz U, Thiele W, Mueller G, Flentje M. Stereotactic radiotherapy for primary lung cancer and pulmonary metastases: a noninvasive treatment approach in medically inoperable patients. *Int J Radiat Oncol Biol Phys* 2004;60:186-196.
3. Hall EL, Giaccia AJ. Repair of radiation damage and the dose-rate effect. In: Hall EJ, Giaccia AJ, editors. *Radiobiology for the Radiologist*, 6th ed. Philadelphia, Lippincott Williams & Wilkins; 2006 p 60-84.
4. Hall EL, Giaccia AJ. Linear energy transfer and relative biological effectiveness. In: Hall EJ, Giaccia AJ, editors. *Radiobiology for the Radiologist*, 6th ed. Philadelphia, Lippincott Williams & Wilkins; 2006 p 106-116.
5. Hall EL, Giaccia AJ. Oxygen effect and reoxygenation. In: Hall EJ, Giaccia AJ, editors. *Radiobiology for the Radiologist*, 6th ed. Philadelphia, Lippincott Williams & Wilkins; 2006 p 85-105.
6. Vaupel P. Tumor microenvironmental physiology and its implications for radiation oncology. *Semin Radiat Oncol* 2004;14:197-275.
7. Masunaga S, Ono K. Significance of the response of quiescent cell populations within solid tumors in cancer therapy. *J Radiat Res* 2002;43:11-25.
8. Torikoshi M, Minohara S, Kanematsu N, Komori M, Kanazawa M, Noda

K, et al. Irradiation system for HIMAC. J Radiat Res
2007;48(Suppl):A15-A25.

9. Chaplin DJ, Horsman MR, Trotter MJ. Effect of nicotinamide on the microregional heterogeneity of oxygen delivery within a murine tumor. J Natl Cancer Inst 1990;82:672-676.
10. Wardman P. Chemical radiosensitizers for use in radiotherapy. Clin Oncol 2007;19:397-417.
11. Masunaga S, Ando K, Uzawa A, Hirayama R, Furusawa Y, Sakurai Y, et al. Radiobiologic significance of the response of intratumor quiescent cells *in vivo* to accelerated carbon ion beams compared with γ -rays and reactor neutron beams. Int J Radiat Oncol Biol Phys 2008;70:221-228.
12. Masunaga S, Ono K, Suzuki M, Nishimura Y, Hiraoka M, Kinashi Y, et al. Alteration of the hypoxic fraction of quiescent cell populations by hyperthermia at mild temperatures. Int J Hyperthermia 1997;13:401-411.
13. Hada M, Georgakilas AG. Formation of clustered DNA damage after high-LET irradiation: a Review. J Radiat Res 2008;49:203-210.
14. Thomlinson RH, Gray LH. The histological structure of some human lung cancers and the possible implications for radiotherapy. Br J Cancer 1955;9:539-549.
15. Brown JM. Evidence of acutely hypoxic cells in mouse tumours, and a possible mechanism of reoxygenation. Br J Radiol 1979;52:650-656.
16. Chan N, Koritzinsky M, Zhao H, Bindra R, Glazer PM, Powell S, et al. Chronic hypoxia decreases synthesis of homologous recombination proteins to offset chemoresistance and radioresistance. Cancer Res 2008;68:605-614.

- 1 17. Sprong D, Janssen HL, Vens C, Begg AC. Resistance of hypoxic cells
2 to ionizing radiation is influenced by homologous recombination
3 status. Int J Radiat Oncol Biol Phys 2006;64:562-572.
4
5
6
- 7 18. Masunaga S, Ono K, Abe M. Potentially lethal damage repair by
8 quiescent cells in murine solid tumors. Int J Radiat Oncol Biol Phys
9
10
11
12 1992;22:973-978.
13
- 14 19. Masunaga S, Nagata K, Suzuki M, Kashino G, Kinashi Y, Ono K.
15
16 Inhibition of repair from radiation-induced damage by mild
17
18 temperature hyperthermia, referring to the effect on quiescent cell
19
20
21 populations. Radiat Med 2007;25:417-425.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Captions for Illustrations

Fig. 1. Cell survival curves for the total tumor cell population as a function of radiation dose after γ -ray irradiation. Open and solid symbols represent the surviving fractions after high dose-rate and reduced dose-rate γ -ray irradiation, respectively. Circles, reversed triangles, triangles, and squares represent the surviving fractions after aerobic irradiation without any drug, irradiation under totally hypoxic conditions, aerobic irradiation after nicotinamide (NA) loading, and aerobic irradiation after misonidazole (MISO) loading, respectively. Bars represent standard errors.

Fig. 2. Cell survival curves for the total tumor cell population as a function of radiation dose after accelerated carbon-ion beam irradiation with an LET of 18 and 50 keV/ μ m are shown in the left and right panels, respectively. Open and solid symbols represent the surviving fractions after high dose-rate and reduced dose-rate accelerated carbon-ion beam irradiation, respectively. Circles, reversed triangles, triangles, and squares represent the surviving fractions after aerobic irradiation without any drug, irradiation under totally hypoxic conditions, aerobic irradiation after nicotinamide (NA) loading, and aerobic irradiation after misonidazole (MISO) loading, respectively. Bars represent standard errors.

Fig. 3. Dose response curves of the normalized micronucleus frequency for the total and quiescent tumor cell populations as a function of radiation dose after γ -ray irradiation are shown in the left and right panels, respectively. Open and solid

symbols represent the normalized micronucleus frequencies after high dose-rate and reduced dose-rate γ -ray irradiation, respectively. Circles, reversed triangles, triangles, and squares represent the normalized micronucleus frequencies after aerobic irradiation without any drug, irradiation under totally hypoxic conditions, aerobic irradiation after nicotinamide (NA) loading, and aerobic irradiation after misonidazole (MISO) loading, respectively. Bars represent standard errors.

Fig. 4. Dose response curves of the normalized micronucleus frequency for the total and quiescent tumor cell populations as a function of radiation dose after accelerated carbon-ion beam irradiation with an LET of 18 keV/ μ m are shown in the left and right panels, respectively. Open and solid symbols represent the normalized micronucleus frequencies after high dose-rate and reduced dose-rate accelerated carbon-ion beam irradiation, respectively. Circles, reversed triangles, triangles, and squares represent the normalized micronucleus frequencies after aerobic irradiation without any drug, irradiation under totally hypoxic conditions, aerobic irradiation after nicotinamide (NA) loading, and aerobic irradiation after misonidazole (MISO) loading, respectively. Bars represent standard errors.

Fig. 5. Dose response curves of the normalized micronucleus frequency for the total and quiescent tumor cell populations as a function of radiation dose after accelerated carbon-ion beam irradiation with an LET of 50 keV/ μ m are shown in the left and

right panels, respectively. Open and solid symbols represent the normalized micronucleus frequencies after high dose-rate and reduced dose-rate accelerated carbon-ion beam irradiation, respectively. Circles, reversed triangles, triangles, and squares represent the normalized micronucleus frequencies after aerobic irradiation without any drug, irradiation under totally hypoxic conditions, aerobic irradiation after nicotinamide (NA) loading, and aerobic irradiation after misonidazole (MISO) loading, respectively. Bars represent standard errors.

Table 1.

Plating efficiency and micronucleus frequency at 0 Gy

	Total tumor cells	Quiescent cells
<Plating efficiency (%)>		
Absolutely control	53.7 ± 8.5 ^a	----
Totally hypoxic	50.1 ± 7.0	----
+ Nicotinamide	46.8 ± 0.6	----
+ Misonidazole	31.2 ± 7.7	
<Micronucleus frequency>		
Absolutely control	0.043 ± 0.005	0.063 ± 0.009
Totally hypoxic	0.058 ± 0.006	0.073 ± 0.010
+ Nicotinamide	0.073 ± 0.009	0.119 ± 0.014
+ Misonidazole	0.093 ± 0.011	0.152 ± 0.018

^a; Mean ± standard deviation

Table 2.

Irradiation under aerobic conditions
compared with irradiation under hypoxic conditions^a

γ -Rays	Carbon-ion beams (18 keV/ μ m)	Carbon-ion beams (50 keV/ μ m)
<Surviving fraction = 0.3>		
<u>Total cells</u>		
High dose-rate irradiation		
2.1 (2.0-2.2) ^b	1.4 (1.3-1.5)	1.15 (1.1-1.2)
Reduced dose-rate irradiation		
1.8 (1.7-2.0)	1.25 (1.15-1.35)	1.1 (1.05-1.15)
<Normalized micronucleus frequency = 0.2>		
<u>Total cells</u>		
High dose-rate irradiation		
1.9 (1.8-2.0)	1.35 (1.25-1.45)	1.2 (1.1-1.3)
Reduced dose-rate irradiation		
1.7 (1.6-1.8)	1.25 (1.15-1.35)	1.15 (1.1-1.2)
<u>Quiescent cells</u>		
High dose-rate irradiation		
1.5 (1.4-1.6)	1.25 (1.15-1.35)	1.15 (1.1-1.2)
Reduced dose-rate irradiation		
1.3 (1.2-1.4)	1.1 (1.05-1.15)	1.05 (1.0-1.1)

^a; The ratio of the dose of radiation necessary to obtain each end-point under hypoxic conditions to that needed to obtain each end-point under aerobic conditions.

^b; Values in parentheses are 95% confidence limits, determined using standard errors. If the ranges of 95 % confidence limits showed no overlap between any two values, the difference between the two values was considered significant ($p < 0.05$).

Table 3.

Enhancement ratios^a due to combination
with nicotinamide or misonidazole

	High dose rate irradiation	Reduced dose rate irradiation
<Surviving fraction = 0.3>		
<u>Total cells</u>		
γ -Rays		
+ Nicotinamide	1.3 (1.2-1.4) ^b	1.05 (1.0-1.1)
+ Misonidazole	1.8 (1.6-2.0)	1.85 (1.7-2.0)
Carbon-ion beams (18 keV/ μ m)		
+ Nicotinamide	1.15 (1.1-1.2)	1.05 (1.0-1.1)
+ Misonidazole	1.3 (1.2-1.4)	1.35 (1.3-1.4)
Carbon-ion beams (50 keV/ μ m)		
+ Nicotinamide	1.05 (1.0-1.1)	1.05 (1.0-1.1)
+ Misonidazole	1.1 (1.05-1.15)	1.15 (1.1-1.2)
<Normalized micronucleus frequency = 0.2>		
<u>Total cells</u>		
γ -Rays		
+ Nicotinamide	1.2 (1.1-1.3)	1.05 (1.0-1.1)
+ Misonidazole	1.3 (1.2-1.4)	1.35 (1.25-1.45)
Carbon-ion beams (18 keV/ μ m)		
+ Nicotinamide	1.15 (1.1-1.2)	1.05 (1.0-1.1)
+ Misonidazole	1.2 (1.1-1.3)	1.25 (1.15-1.35)
Carbon-ion beams (50 keV/ μ m)		
+ Nicotinamide	1.05 (1.0-1.1)	1.05 (1.0-1.1)
+ Misonidazole	1.05 (1.0-1.1)	1.05 (1.0-1.1)
<u>Quiescent cells</u>		
γ -Rays		
+ Nicotinamide	1.1 (1.05-1.15)	1.05 (1.0-1.1)
+ Misonidazole	1.45 (1.35-1.55)	1.5 (1.4-1.6)
Carbon-ion beams (18 keV/ μ m)		
+ Nicotinamide	1.05 (1.0-1.1)	1.0 (0.95-1.05)
+ Misonidazole	1.25 (1.15-1.35)	1.3 (1.2-1.4)
Carbon-ion beams (50 keV/ μ m)		
+ Nicotinamide	1.05 (1.0-1.1)	1.0 (0.95-1.05)
+ Misonidazole	1.1 (1.05-1.15)	1.1 (1.05-1.15)

^a; The ratio of the dose of radiation necessary to obtain each end-point without the drug to that needed to obtain each end-point with the drug.

^b; As in Table 2.

Table 4.

Dose-modifying factors due to the reduction in **radiosensitivity** caused by a decrease in radiation dose rate^a

γ -Rays	Carbon-ion beams (18 keV/ μ m)	Carbon-ion beams (50 keV/ μ m)
<Surviving fraction = 0.3>		
<u>Total cells</u>		
Radiation only		
1.3 (1.2-1.4) ^b	1.2 (1.1-1.3)	1.15 (1.1-1.2)
Totally hypoxic		
1.2 (1.1-1.3)	1.15 (1.05-1.25)	1.1 (1.05-1.15)
+ Nicotinamide		
1.55 (1.4-1.7)	1.35 (1.25-1.45)	1.2 (1.15-1.25)
+ Misonidazole		
1.25 (1.15-1.35)	1.2 (1.1-1.3)	1.15 (1.1-1.2)
<Normalized micronucleus frequency = 0.2>		
<u>Total cells</u>		
Radiation only		
1.25 (1.15-1.35)	1.2 (1.1-1.3)	1.1 (1.05-1.15)
Totally hypoxic		
1.15 (1.1-1.2)	1.1 (1.05-1.15)	1.05 (1.0-1.1)
+ Nicotinamide		
1.4 (1.25-1.55)	1.3 (1.2-1.4)	1.2 (1.1-1.3)
+ Misonidazole		
1.2 (1.1-1.3)	1.15 (1.05-1.25)	1.1 (1.05-1.15)
<u>Quiescent cells</u>		
Radiation only		
1.4 (1.3-1.5)	1.3 (1.2-1.4)	1.25 (1.15-1.35)
Totally hypoxic		
1.25 (1.15-1.35)	1.15 (1.1-1.2)	1.1 (1.05-1.15)
+ Nicotinamide		
1.4 (1.3-1.5)	1.3 (1.2-1.4)	1.25 (1.15-1.35)
+ Misonidazole		
1.15 (1.1-1.2)	1.1 (1.05-1.15)	1.1 (1.05-1.15)

^a; The ratio of the dose of radiation necessary to obtain each end-point with reduced dose-rate irradiation to that needed to obtain each end-point with high dose-rate irradiation.

^b; As in Table 2.

Table 5.

**Dose-modifying factors for quiescent cells
relative to total tumor cells^a**

γ -Rays	Carbon-ion beams (18 keV/ μ m)	Carbon-ion beams (50 keV/ μ m)
<Normalized micronucleus frequency = 0.2>		
<u>High dose rate irradiation</u>		
Radiation only ^b		
1.65 (1.5-1.8) ^b	1.45 (1.35-1.55)	1.35 (1.25-1.45)
Totally hypoxic		
1.4 (1.15-1.55)	1.3 (1.2-1.4)	1.25 (1.15-1.35)
+ Nicotinamide		
1.75 (1.6-1.9)	1.5 (1.4-1.6)	1.4 (1.3-1.5)
+ Misonidazole		
1.5 (1.35-1.65)	1.35 (1.25-1.45)	1.3 (1.2-1.4)
<u>Reduced dose rate irradiation</u>		
Radiation only		
1.85 (1.7-2.0)	1.6 (1.5-1.7)	1.5 (1.4-1.6)
Totally hypoxic		
1.45 (1.35-1.55)	1.35 (1.25-1.45)	1.3 (1.2-1.4)
+ Nicotinamide		
1.85 (1.7-2.0)	1.6 (1.5-1.7)	1.5 (1.4-1.7)
+ Misonidazole		
1.55 (1.45-1.65)	1.4 (1.3-1.5)	1.35 (1.25-1.45)

^a; The ratio of the dose of radiation necessary to obtain each normalized micronucleus frequency in quiescent cell population to that needed to obtain each normalized micronucleus frequency in total tumor cell population.

^b; As in Table 2.

Figure 1
[Click here to download high resolution image](#)




Figure 2
[Click here to download high resolution image](#)




Figure 3
[Click here to download high resolution image](#)




Figure 4
[Click here to download high resolution image](#)





Figure 5
[Click here to download high resolution image](#)



Acknowledgments

This study was supported, in part, by a Grant-in-aid for Scientific Research (C) (20591493) from the Japan Society for the Promotion of Science.