

END INVARIANTS OF HECKOID GROUPS FOR 2-BRIDGE LINKS

DONGHI LEE AND MAKOTO SAKUMA

1. INTRODUCTION

By extending the concept of a geometrically infinite end of a Kleinian group, Bowditch [4] introduced the notion of the end invariants of a type-preserving $SL(2, \mathbb{C})$ -representation of the fundamental group $\pi_1(\mathbf{T})$ of the once-punctured torus $\mathbf{T}$. Tan, Wong and Zhang [23, 24] extended this notion (with slight modification) to an arbitrary $SL(2, \mathbb{C})$ -representation of $\pi_1(\mathbf{T})$. In [12], we gave an explicit description of the sets of end invariants of the $SL(2, \mathbb{C})$ -characters of the once-punctured torus corresponding to the holonomy representation of hyperbolic 2-bridge link groups. The purpose of this note is to announce the result obtained in [14] which explicitly describes the sets of end invariants of the $SL(2, \mathbb{C})$ -characters of the once-punctured torus corresponding to the holonomy representation of Heckoid groups (Theorem 4.1).

2. BOWDITCH, TAN-WONG-ZHANG END INVARIANTS

Motivated by the definition of the end of a geometrically infinite end of a Kleinian group, Bowditch [4] introduced the notion of the end invariants of an arbitrary type-preserving $PSL(2, \mathbb{C})$ -representation of $\pi_1(\mathbf{T})$. Tan, Wong and Zhang [23, 24] extended this notion (with slight modification) to an arbitrary $PSL(2, \mathbb{C})$ -representation of $\pi_1(\mathbf{T})$. To describe this, let $\mathcal{C}$ be the set of free homotopy classes of essential simple loops on $\mathbf{T}$. Then $\mathcal{C}$ is identified with $\hat{\mathbb{Q}}$, the vertex set of the Farey tessellation $\mathcal{D}$, by the following rule $s \mapsto \beta_s$, where β_s is the image of a line in $\mathbb{R}^2 - \mathbb{Z}^2$ of slope s in $\mathbf{T} = (\mathbb{R}^2 - \mathbb{Z}^2)/\mathbb{Z}^2$. The projective lamination space $\mathcal{PL}$ of $\mathbf{T}$ is then identified with $\hat{\mathbb{R}} := \mathbb{R} \cup \{\infty\}$ and contains $\mathcal{C}$ as the dense subset of rational points.

Definition 2.1. Let ρ be a $PSL(2, \mathbb{C})$ -representation of $\pi_1(\mathbf{T})$.

- (1) An element $X \in \mathcal{PL}$ is an *end invariant* of ρ if there exists a sequence of distinct elements $X_n \in \mathcal{C}$ such that $X_n \rightarrow X$ and that $\{|\mathrm{tr}\rho(X_n)|\}_n$ is bounded from above.
- (2) $\mathcal{E}(\rho)$ denotes the set of end invariants of ρ .

In the above definition, it should be noted that $|\mathrm{tr}\rho(X_n)|$ is well-defined though $\mathrm{tr}\rho(X_n)$ is defined only up to sign. Note also that the condition that $\{|\mathrm{tr}\rho(X_n)|\}_n$ is bounded from above is equivalent to the condition that the (real) hyperbolic translation lengths of the isometries $\rho(X_n)$ of $\mathbb{H}^3$ are bounded from above. So, if ρ is a faithful discrete type-preserving representation and ν is the end invariant of a geometrically infinite end of the quotient hyperbolic manifold, then ν is an end invariant of ρ in the sense of the above definition.

Tan, Wong and Zhang [23, 24] showed that $\mathcal{E}(\rho)$ is a closed subset of $\mathcal{PL}$ and proved various interesting properties of $\mathcal{E}(\rho)$, including a characterization of those representations ρ with $\mathcal{E}(\rho) = \emptyset$ or $\mathcal{PL}$, generalizing results of Bowditch [4]. They also proposed an

Received December 8, 2012.

interesting conjecture [24, Conjecture 1.8] concerning possible homeomorphism types of $\mathcal{E}(\rho)$. The following is a modified version of the conjecture which Tan [22] informed to the authors.

Conjecture 2.2. Suppose $\mathcal{E}(\rho)$ has at least two accumulation points. Then either $\mathcal{E}(\rho) = \mathcal{PL}$ or a Cantor set of $\mathcal{PL}$.

They constructed a family of representations ρ which have Cantor sets as $\mathcal{E}(\rho)$, and proved the following supporting evidence to the conjecture (see [24, Theorem 1.7]).

Theorem 2.3. Let $\rho : \pi_1(\mathbf{T}) \rightarrow \mathrm{SL}(2, \mathbb{C})$ be discrete in the sense that the set $\{\mathrm{tr}(\rho(X)) \mid X \in \mathcal{C}\}$ is discrete in $\mathbb{C}$. Then if $\mathcal{E}(\rho)$ has at least three elements, then $\mathcal{E}(\rho)$ is either a Cantor set of $\mathcal{PL}$ or all of $\mathcal{PL}$.

However, the above theorem does not describe the set $\mathcal{E}(\rho)$ explicitly. In [12], we gave an explicit description of the sets of end invariants of the $\mathrm{SL}(2, \mathbb{C})$ -characters of the once-punctured torus corresponding to the holonomy representation of hyperbolic 2-bridge link groups. In this note, we announce a result obtained in [14] which explicitly describes the sets of end invariants of the $\mathrm{SL}(2, \mathbb{C})$ -characters of the once-punctured torus corresponding to the holonomy representation of Heckoid groups (Theorem 4.1). These give an infinite family of representations ρ for which $\mathcal{E}(\rho)$ are explicitly described Cantor sets.

3. HECKOID ORBIFOLD $\mathcal{S}(r; n)$ AND HECKOID GROUP $G(r; n)$

For a rational number $r \in \hat{\mathbb{Q}} := \mathbb{Q} \cup \{\infty\}$, let $K(r)$ be the 2-bridge link of slope r , which is defined as the sum $(S^3, K(r)) = (B^3, t(\infty)) \cup (B^3, t(r))$ of rational tangles of slope ∞ and r . The common boundary $\partial(B^3, t(\infty)) = \partial(B^3, t(r))$ of the rational tangles is identified with the *Conway sphere* $(\mathcal{S}^2, \mathbf{P}) := (\mathbb{R}^2, \mathbb{Z}^2)/H$, where H is the group of isometries of the Euclidean plane $\mathbb{R}^2$ generated by the π -rotations around the points in the lattice $\mathbb{Z}^2$. Let $\mathcal{S}$ be the 4-punctured sphere $\mathcal{S}^2 - \mathbf{P}$ in the link complement $S^3 - K(r)$. Any essential simple loop in $\mathcal{S}$, up to isotopy, is obtained as the image of a line of slope $s \in \hat{\mathbb{Q}}$ in $\mathbb{R}^2 - \mathbb{Z}^2$ by the covering projection onto $\mathcal{S}$. The (unoriented) essential simple loop in $\mathcal{S}$ so obtained is denoted by α_s . We also denote by α_s the conjugacy class of an element of $\pi_1(\mathcal{S})$ represented by (a suitably oriented) α_s . The loops α_∞ and α_r bound disks in $B^3 - t(\infty)$ and $B^3 - t(r)$, respectively. Thus the *link group* $G(K(r)) = \pi_1(S^3 - K(r))$ is obtained as follows:

$$G(K(r)) = \pi_1(S^3 - K(r)) \cong \pi_1(\mathcal{S}) / \langle\langle \alpha_\infty, \alpha_r \rangle\rangle \cong \pi_1(B^3 - t(\infty)) / \langle\langle \alpha_r \rangle\rangle.$$

For each rational number r and an integer $n \geq 2$, the *even Heckoid orbifold of index n for the 2-bridge link $K(r)$* is the 3-orbifold $\mathcal{S}(r; n)$, such that the underlying space $|\mathcal{S}(r; n)|$ is the exterior, $E(K(r)) = S^3 - \mathrm{int} N(K(r))$, of $K(r)$, and that the singular set is the lower tunnel of $K(r)$ (i.e., the core tunnel of $(B^3, t(\infty))$ in the sense of [10, p.360]), where the index of the singularity is n (see Figure 1). We call the orbifold fundamental group $\pi_1(\mathcal{S}(r; n))$ the *Heckoid group of index n for $K(r)$* , and denote it by $G(r; n)$. Since the loop α_r is isotopic to a meridional loop around the lower tunnel, the even Heckoid group $G(r; n) = \pi_1(\mathcal{S}(r; n))$ ($n \geq 2$) is obtained as follows:

$$G(r; n) = \pi_1(\mathcal{S}(r; n)) \cong \pi_1(\mathcal{S}) / \langle\langle \alpha_\infty, \alpha_r^n \rangle\rangle \cong \pi_1(B^3 - t(\infty)) / \langle\langle \alpha_r^n \rangle\rangle.$$

The announcement by Agol [1] and the announcement made in the second author's joint work with Akiyoshi, Wada and Yamashita in [2, Section 3 of Preface] suggest that




FIGURE 1. The Heckoid orbifold $S(r; n)$. The labels ∞ indicate the parabolic loci. Here $(S^3, K(r)) = (B^3, t(\infty)) \cup (B^3, t(r))$ with $r = [4, 2] = 2/9$, where $(B^3, t(r))$ and $(B^3, t(\infty))$, respectively, are the inside and the outside of the bridge sphere S^2 . The lower tunnel is the core tunnel of $(B^3, t(r))$.

the group $G(r; n)$ makes sense even when n is a half-integer greater than 1. We refer to [14, Definition 3.2] for the definition of the group $G(r; n)$ and the corresponding orbifold $S(r; n)$ when n is a non-integral half-integer greater than 1. Roughly speaking, $S(r; n)$ is defined to be a $\mathbb{Z}/2\mathbb{Z}$ -covering of a certain orbifold $O(r; m)$, with $m = 2n$, which is obtained from the quotient of $K(r)$ by the natural $(\mathbb{Z}/2\mathbb{Z})^2$ -symmetry (see Figure 2 for the case when $K(r)$ is a knot). We call them the *odd Heckoid orbifold* and the *odd Heckoid group*, respectively, of index n for $K(r)$. A topological description of an odd Heckoid orbifold is given by [14, Proposition 5.3 and Figures 5 and 6].

Remark 3.1. Our terminology is slightly different from that of Riley [20], where $G(r; n)$ is called the Heckoid group of index “ m ” for $K(r)$ with $m = 2n$. The Heckoid orbifold $S(r; n)$ and the Heckoid group $G(r; n)$ are *even* or *odd* according to whether Riley’s index $m = 2n$ is even or odd.

The following theorem was anticipated in [20] and is contained in [1] without proof.

Theorem 3.2. *For a rational number r and an integer or a half-integer $n > 1$, the Heckoid group $G(r; n)$ is isomorphic to a geometrically finite Kleinian group generated by two parabolic transformations.*

A proof of this theorem is given in [14, Section 6] by using the orbifold theorem for pared orbifolds [3, Theorem 8.3.9] (cf. [5, 8]). As noted in [1], the proof is analogous to the arguments in [7, Proof of Theorem 9].

By this theorem and the topological description of odd Heckoid orbifolds ([14, Proposition 5.3]), we obtain the following proposition, which shows a significant difference between odd and even Heckoid groups (see [14, Section 6]).

Proposition 3.3. *Any odd Heckoid group is not a one-relator group.*

4. END INVARIANTS OF EVEN HECKOID GROUPS

For a rational number r and an integer $n \geq 2$, let $\rho_{r,n}$ be the $\mathrm{PSL}(2, \mathbb{C})$ -representation of $\pi_1(S)$ obtained as the composition

$$\pi_1(S) \rightarrow \pi_1(S)/\langle\langle \alpha_\infty, \alpha_r^n \rangle\rangle \cong G(r; n) \rightarrow \mathrm{Isom}^+(\mathbb{H}^3) \cong \mathrm{PSL}(2, \mathbb{C}),$$




FIGURE 2. The case when $K(r)$ is a knot and $m = 2n > 1$ is an odd integer. Here $r = 2/9 = [4, 2]$. The odd Heckoid orbifold $S(r; n)$ (middle right) is a $\mathbb{Z}/2\mathbb{Z}$ -covering of $O(r; m)$ (lower left). The upper left figure is not an orbifold, but is a hyperbolic cone manifold. The odd Heckoid orbifold $S(r; n)$ is the quotient of the cone manifold by the π -rotation around the axis containing the singular set.

where the last homomorphism is the holonomy representation of the pared hyperbolic orbifold $S(r; n)$.

Now, let O be the orbifold $(\mathbb{R}^2 - \mathbb{Z}^2)/\hat{H}$ where $\hat{H}$ is the group generated by π -rotations around the points in $(\frac{1}{2}\mathbb{Z})^2$. Note that O is the orbifold with underlying space a once-punctured sphere and with three cone points of cone angle π . The surfaces T and S , respectively, are $\mathbb{Z}/2\mathbb{Z}$ -covering and $(\mathbb{Z}/2\mathbb{Z})^2$ -covering of O , and hence their fundamental groups are identified with subgroups of the orbifold fundamental group $\pi_1(O)$ of indices 2 and 4, respectively. The $\mathrm{PSL}(2, \mathbb{C})$ -representation $\rho_{r,n}$ of $\pi_1(S)$ extends, in a unique way, to that of $\pi_1(O)$ (see [2, Proposition 2.2]), and so we obtain, in a unique way, a

$\mathrm{PSL}(2, \mathbb{C})$ -representation of $\pi_1(\mathcal{T})$ by restriction. We continue to denote it by $\rho_{r,n}$. The following theorem, which determines the set $\mathcal{E}(\rho_{r,n})$, is obtained by [16].

Theorem 4.1. *For a non-integral rational number r and an integer $n \geq 2$, the set $\mathcal{E}(\rho_{r,n})$ of end invariants of $\rho_{r,n}$ is equal to the limit set $\Lambda(\Gamma(r;n))$ of the group $\Gamma(r;n)$.*

5. SIMPLE LOOPS ON BRIDGE SPHERES OF HECKOID ORBIFOLDS

Let $\mathcal{D}$ be the *Farey tessellation*, that is, the tessellation of the upper half space $\mathbb{H}^2$ by ideal triangles which are obtained from the ideal triangle with the ideal vertices $0, 1, \infty \in \hat{\mathbb{Q}}$ by repeated reflection in the edges. Then $\hat{\mathbb{Q}}$ is identified with the set of the ideal vertices of $\mathcal{D}$. For each $r \in \hat{\mathbb{Q}}$, let Γ_r be the group of automorphisms of $\mathcal{D}$ generated by reflections in the edges of $\mathcal{D}$ with an endpoint r . It should be noted that Γ_r is isomorphic to the infinite dihedral group and that the region bounded by two adjacent edges of $\mathcal{D}$ with an endpoint r is a fundamental domain for the action of Γ_r on $\mathbb{H}^2$. For an integer m , let $C_r(m)$ be the group of automorphisms of $\mathcal{D}$ generated by the parabolic transformation, centered on the vertex r , by m units in the clockwise direction.

For r a rational number and n an integer or a half-integer greater than 1, let $\Gamma(r;n)$ be the group generated by Γ_∞ and $C_r(2n)$. Suppose that r is not an integer. Then $\Gamma(r;n)$ is the free product $\Gamma_\infty * C_r(2n)$ having a fundamental domain, R , shown in Figure 3. Here, R is obtained as the intersection of fundamental domains for Γ_∞ and $C_r(2n)$, and so R is bounded by the following two pairs of Farey edges:

- (1) the pair of adjacent Farey edges with an endpoint ∞ which cuts off a region in $\bar{\mathbb{H}}^2$ containing r , and
- (2) a pair of Farey edges with an endpoint r which cuts off a region in $\bar{\mathbb{H}}^2$ containing ∞ , such that one edge is the image of the other by a generator of $C_r(2n)$.

Let $\bar{I}(n;r)$ be the union of two closed intervals in $\partial\mathbb{H}^2 = \hat{\mathbb{R}}$ obtained as the intersection of the closure of R with $\partial\mathbb{H}^2$. Note that there is a pair $\{r_1, r_2\}$ of boundary points of $\bar{I}(n;r)$ such that r_2 is the image of r_1 by a generator of $C_r(2n)$. Set $I(n;r) = \bar{I}(n;r) - \{r_i\}$ with $i = 1$ or 2 . Note that $I(n;r)$ is the disjoint union of a closed interval and a half-open interval, except for the special case when $r \equiv \pm 1/p \pmod{\mathbb{Z}}$.




FIGURE 3. A fundamental domain of $\Gamma(r;n)$ in the Farey tessellation (the shaded domain) for $r = 3/10 = \frac{1}{3 + \frac{1}{3}}$ and $n = 2$

The following theorem proved in [14] is the starting point of all the results which we announce in this note.

Theorem 5.1. *Suppose that r is a non-integral rational number and that n is an integer or a half integer greater than 1. Then, for any $s \in \hat{\mathbb{Q}}$, there is a unique rational number $s_0 \in I(r; n) \cup \{\infty, r\}$ such that s is contained in the $\Gamma(r; n)$ -orbit of s_0 . Moreover α_s is homotopic to α_{s_0} in $S(r; n)$. In particular, if $s_0 = \infty$, then α_s is null-homotopic in $S(r; n)$.*

Theorem 5.2 is proved in [14], and Theorems 5.3 and 5.4 will be proved in [15].

Theorem 5.2. *Suppose that r is a non-integral rational number and that n is an integer with $n \geq 2$. Then the loop α_s is null-homotopic in $S(r; n)$ if and only if s belongs to the $\Gamma(r; n)$ -orbit of ∞ . In other words, if $s \in I(r; n) \cup \{r\}$, then α_s is not null-homotopic in $S(r; n)$.*

Theorem 5.3. *Suppose that r is a non-integral rational number and that n is an integer with $n \geq 2$. For two rational numbers s and s' , the simple loops α_s and $\alpha_{s'}$ are homotopic in $S(r; n)$ if and only if s and s' belong to the same $\Gamma(r; n)$ -orbit. In other words, for distinct $s, s' \in I(r; n) \cup \{\infty, r\}$, the simple loops α_s and $\alpha_{s'}$ are not homotopic in $S(r; n)$.*

Theorem 5.4. *Suppose that r is a non-integral rational number and that n is an integer with $n \geq 2$. Then the following hold.*

- (1) *The loop α_s is peripheral in $S(r; n)$ if and only if s belongs to the $\Gamma(r; n)$ -orbit of ∞ .*
- (2) *The loop α_s is torsion in $S(r; n)$ if and only if s belongs to the $\Gamma(r; n)$ -orbit of ∞ or r .*

In other words, there is no rational number $s \in I(r; n)$ for which the simple loop α_s is peripheral or torsion in $S(r; n)$.

In the above theorem, we say that α_s is *peripheral* or *torsion* if the conjugacy class α_s is represented by a (possibly trivial) parabolic or elliptic transformation, respectively, when we identify $G(r; n)$ with a Kleinian group generated by two parabolic transformations.

These theorems are proved by using the small cancellation theory [17]. Please see [13] for basic ideas of the proof. Theorem 4.1 is proved by using these theorems, Bowditch's results [4] and the discreteness of marked length spectrum of geometrically finite hyperbolic 3-manifolds, as in [12, Section 8].

REFERENCES

- [1] I. Agol, *The classification of non-free 2-parabolic generator Kleinian groups*, Slides of talks given at Austin AMS Meeting and Budapest Bolyai conference, July 2002, Budapest, Hungary.
- [2] H. Akiyoshi, M. Sakuma, M. Wada, and Y. Yamashita, *Punctured torus groups and 2-bridge knot groups (I)*, Lecture Notes in Mathematics **1909**, Springer, Berlin, 2007.
- [3] M. Boileau, and J. Porti, *Geometrization of 3-orbifolds of cyclic type*, Appendix A by Michael Heusener and Porti, Astérisque No. 272 (2001).
- [4] B. H. Bowditch, *Markoff triples and quasifuchsian groups*, Proc. London Math. Soc. **77** (1998), 697–736.
- [5] D. Cooper, C. D. Hodgson, and S. P. Kerckhoff *Three-dimensional orbifolds and cone-manifolds*, MSJ Memoirs, **5**, Mathematical Society of Japan, Tokyo, 2000.
- [6] E. Hecke, *Über die Bestimmung Dirichletscher Reihen durch ihre Funktionalgleichung*, Math. Ann. **112** (1936), 664–699.

- [7] K. N. Jones and A. W. Reid, *Minimal index torsion-free subgroups of Kleinian groups.* Math. Ann. **310** (1998), 235–250.
- [8] M. Kapovich, *Hyperbolic manifolds and discrete groups*, Progress in Mathematics, **183**, Birkhäuser Boston, Inc., Boston, MA, 2001.
- [9] D. Lee and M. Sakuma, *Simple loops on 2-bridge spheres in 2-bridge link complements*, Electron. Res. Announc. Math. Sci. **18** (2011), 97–111.
- [10] D. Lee and M. Sakuma, *Epimorphisms between 2-bridge link groups: Homotopically trivial simple loops on 2-bridge spheres*, Proc. London Math. Soc. **104** (2012), 359–386.
- [11] D. Lee and M. Sakuma, *Homotopically equivalent simple loops on 2-bridge spheres in 2-bridge link complements (I), (II) and (III)*, arXiv:1010.2232, arXiv:1103.0856, arXiv:1111.3562.
- [12] D. Lee and M. Sakuma, *A variation of McShane’s identity for 2-bridge links*, arXiv:1112.5859.
- [13] D. Lee and M. Sakuma, *Simple loops on 2-bridge spheres in Heckoid orbifolds for 2-bridge links*, Electron. Res. Announc. Math. Sci. **19** (2012), 97–111.
- [14] D. Lee and M. Sakuma, *Epimorphisms from 2-bridge link groups onto Heckoid groups (I) and (II)*, to appear in Hiroshima J. Math..
- [15] D. Lee and M. Sakuma, *Homotopically equivalent simple loops on 2-bridge spheres in even Heckoid orbifold for 2-bridge links*, preliminary notes.
- [16] D. Lee and M. Sakuma, *A variation of McShane’s identity for even Heckoid orbifolds for 2-bridge links*, in preparation.
- [17] R. C. Lyndon and P. E. Schupp, *Combinatorial group theory*, Springer-Verlag, Berlin, 1977.
- [18] T. Ohtsuki, R. Riley, and M. Sakuma, *Epimorphisms between 2-bridge link groups*, Geom. Topol. Monogr. **14** (2008), 417–450.
- [19] R. Riley, *Parabolic representations of knot groups. I*, Proc. London Math. Soc. **24** (1972), 217–242.
- [20] R. Riley, *Algebra for Heckoid groups*, Trans. Amer. Math. Soc. **334** (1992), 389–409.
- [21] R. Riley, *A personal account of the discovery of hyperbolic structures on some knot complements. With a postscript by M. B. Brin, G. A. Jones and D. Singerman*, preprint.
- [22] S. P. Tan, *Private communication*, May, 2011.
- [23] S. P. Tan, Y. L. Wong, and Y. Zhang, *$SL(2, \mathbb{C})$ character variety of a one-holed torus*, Electron. Res. Announc. Amer. Math. Soc. **11** (2005), 103–110.
- [24] S. P. Tan, Y. L. Wong, and Y. Zhang, *End invariants for $SL(2, \mathbb{C})$ characters of the one-holed torus*, Amer. J. Math. **130** (2008), 385–412.

DEPARTMENT OF MATHEMATICS, PUSAN NATIONAL UNIVERSITY, SAN-30 JANGJEON-DONG,
GEUMJUNG-GU, PUSAN, 609-735, KOREA
E-mail address: donghi@pusan.ac.kr

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, HIROSHIMA UNIVERSITY,
HIGASHI-HIROSHIMA, 739-8526, JAPAN
E-mail address: sakuma@math.sci.hiroshima-u.ac.jp