

Comparing *Mimiviridae* communities using meta-barcoding
during a red tide period in an enclosed bay

京都大学 化学研究所 バイオインフォマティクスセンター 化学生命科学領域 Xia Jun

研究成果概要

This study is using the supercomputer of Institute for Chemical Research, Kyoto University, to do an analysis for comparing *Mimiviridae* communities during a red tide period in an enclosed bay, west Japan.

“*Mimiviridae*” is a proposed family of giant viruses infecting unicellular eukaryotes. They are very abundant and ubiquitous in the sea water and affect marine microbial community by their lytic infection cycle. However, only a few *Mimiviridae* members have been explored in the marine environments, and their ecological roles in marine ecosystems remains unknown.

For studying the differences of *Mimiviridae* community in two different kind of water, surface sea water samples (0.22-3 μm) were collected from an enclosed bay (Uranouchi Inlet, Kochi) during a red tide period. A set of degenerated primers (called “MEGAPRIMER”), which target family B DNA polymerase genes of *Mimiviridae*, was used to assess the composition of *Mimiviridae* community. Bioinformatic analysis was applied after sequencing. As a result, hundreds of operational taxonomic units (OTUs) were identified in each sample. When we compared the *Mimiviridae* structures between within and without red-tide waters, there was a clear difference between these two samples in a community variation analysis. The difference was also visible in dominant OTU proportions between the two types of samples. Phylogenetic analysis based on a maximum-likelihood method showed that almost all of the OTUs belong to unknown *Mimiviridae* branches. Future study will mainly focus on the host-virus interactions and the relationship between viral community and environmental factors, that may reveal ecological functions of *Mimiviridae* in a coastal ecosystem.