

Organic Materials Chemistry

- Organic Structural Chemistry -

<http://133.3.40.20/>



Prof
KOMATSU, Koichi
(D Eng)



Assoc Prof
KITAGAWA, Toshikazu
(D Eng)



Instr
NISHINAGA, Tohru
(D Eng)



Instr
MURATA, Yasujiro
(D Eng)



Guest Res Assoc
CHENG, Fuyong
(D Sc)

Students

WAKAMIYA, Atsushi (D3)
IDOMOTO, Yuichi (M2)
MASAOKA, Naoki (M1)
YAMAZAKI, Tetsuya (UG)

LEE, Yangsoo (D2)
MIYATA, Yasuo (M2)
YAMAZAKI, Daisuke (M1)

OGAWA, Kohei (D1)
MURATA, Michihisa (M2)
UTO, Takayuki (UG)

Scope of Research

Fundamental studies are being conducted for creation of new functional materials with novel structures and properties. The major subjects are: synthetic and structural studies on novel cyclic π -conjugated systems, particularly the positively charged species stabilized by σ - π interaction; synthesis of new redox-active and supramolecular π -systems; organo-chemical transformation of fullerene C_{60} , specifically the synthesis of fullerene dimers and trimers by the use of mechanochemical solid-state reactions; synthesis and reactions of open-cage fullerene derivatives; generation of alkylated C_{60} cation and its application for synthesis of functional materials.

Research Activities (Year 2002)

Presentations

Synthesis of New Fullerene Dimers and Open-Cage Fullerenes by Solid-State and Liquid-Phase Reactions of C_{60} , Komatsu K, Murata Y, et al., The 201st Meeting of the Electrochemical Society, 16 May, Philadelphia, USA.

The Solvent-Free, Solid-State Reaction of Fullerenes to Give Novel Fullerene Dimers, Komatsu K, The 5th International Symposium on Green Chemistry in China, 23 May, Hefei, China.

Radical Cation and Dication of Sulfur-Containing π -Electron Systems Stabilized by Rigid σ -Frameworks, Nishinaga T, Komatsu K, et al., The 5th International Symposium on Functional π -Electron Systems, 1 June, Ulm, Germany.

Fullerene-Based Nanomaterials, Komatsu K, The 12th International Symposium on Supramolecular Chemistry, 7 October, Eilat, Israel.

Structure and Properties of a Novel Fullerene Dimer $RC_{60}-C_{60}R$ and Facile Generation of Fullerenyl Cation RC_{60}^+ , Cheng F, Murata Y, Kitagawa T, Komatsu K, Gordon Research Conference, 29 July, Aioi, Japan.

Grants

Komatsu K, Appearance of biradical character in π -conjugated hydrocarbons by annelation with strained bicyclic frameworks, Grant-in-Aid for Scientific Research (B) (2), Apr. 2002 - Mar. 2004.

Kitagawa T, Construction of fullerene-coated nanostructures using alkylfullerenyl cations, Grant-in-Aid for Scientific Research (C) (2), Apr. 2002 - Mar. 2004.

Nishinaga T, Design and synthesis of π -conjugated oligomers folded in helical structure by electron transfer, Grant-in-Aid for Encouragement of Young Scientists (A), Apr. 2002 - Mar. 2004.


Murata Y, Synthesis of endohedral fullerenes by the use of organic chemistry, Grant-in-Aid for Encouragement of Young Scientists (A), Apr. 2002 - Mar. 2004.

Kitagawa T, Construction of nano-electronic devices based on precise molecular design, CREST, Japan Science and Technology Corporation, Nov. 2002 - Oct. 2007.

Murata Y, Komatsu K, Application of functionalized fullerenes for fabrication of solar cells, Academic-Industrial Cooperative Research Fund, Aug. 2002 - Jul. 2003.

Formation of the First Stable Radical Cation Having a Tetrathia 4-Center/7-Electron Bond


Annulation of cyclic π -conjugated systems with rigid bicycloalkene frameworks has been shown to stabilize the π -system when it is positively charged, both by the σ - π conjugation and steric protection effects. Thus, the radical cation and dication of 1,2-dithiin (1,2-dithia-3,5-cyclohexadiene) were generated as stable species by this structure modification for the first time. Furthermore, the radical cation was found to disproportionate to give a totally new radical cation having the tetrathiabicyclic structure, which is remarkably stabilized by a strong transannular interaction among four sulfur atoms forming a novel 4-center/7-electron system, together with a thiophene derivative.[1]



1. Wakamiya A, Nishinaga T, Komatsu K, *J. Am. Chem. Soc.*, **2002**, 124, 15038.

Generation and Thermodynamic Stabilities of Alkylfullerenyl Cations


The generation and the direct observation of functionalized fullerenyl cations have been considered a challenging task, owing to the electronegative nature of the C_{60} cage. We could generate monoalkylated fullerenyl cations (RC_{60}^+ , R = chloroalkyl group) as long-lived ions by treating the corresponding fullereneol precursors with a strong acid. Spectral analysis showed that these cations have structures with C_s symmetry and their positive charge is mainly located on C-2. Kinetic studies of the S_N1 solvolysis of the corresponding chlorides and the equilibrium constants of the hydrolysis to form fullereneols indicated that the cations RC_{60}^+ have stabilities comparable to that of the *tert*-butyl cation.[2]



2. Kitagawa T, Lee Y, Komatsu K, et al., *Chem. Commun.*, **2002**, 3062.

Synthesis of a Novel Open-Cage Fullerene Derivative with a 13-Membered-Ring Orifice

As the first step in a novel approach to endohedral fullerene complexes, which are expected as nano-scale materials with unique properties, organic synthesis of open-cage fullerenes is quite important. When a thermal reaction of C_{60} with a 1,2,4-triazine derivative was conducted, a [4+2] cycloaddition followed by nitrogen extrusion took place to give a derivative with an 8-membered-ring orifice. This orifice was further enlarged by photochemical oxidation followed by sulfur insertion to the rim of the orifice to give a totally new open-cage fullerene derivative with a 13-membered-ring opening, which is the largest orifice reported so far. A preliminary experiment indicated that it is quite promising to incorporate small molecules into the inside of this fullerene cage.[3]



3. Murata Y, Murata M, Komatsu K, *Chem. Eur. J.*, **2003**, 9, in press.