

Bounded cohomology of subgroups of mapping class groups

東北大学 藤原耕二 (Koji Fujiwara)

Math Institute, Tohoku University

I give a talk on a joint work with Mladen Bestvina [3].

When G is a discrete group, a *quasi-homomorphism* on G is a function $h : G \rightarrow \mathbf{R}$ such that

$$\Delta(h) := \sup_{\gamma_1, \gamma_2 \in G} |h(\gamma_1 \gamma_2) - h(\gamma_1) - h(\gamma_2)| < \infty.$$

The number $\Delta(h)$ is the *defect* of h . We denote by $QH(G)$ the vector space of all quasi-homomorphisms $G \rightarrow \mathbf{R}$ modulo the subspace of bounded functions, and by $\widetilde{QH}(G)$ the vector space of all quasi-homomorphisms $G \rightarrow \mathbf{R}$ modulo the subspace of functions within uniform distance to a homomorphism.

Let S be a compact orientable surface of genus g and p punctures. We consider the associated mapping class group $Mod(S)$ of S . This group acts on the *curve complex* X of S defined by Harvey [7] and successfully used in the study of mapping class groups by Harer [6], [5]. For our purposes, we will restrict to the 1-skeleton of Harvey's complex, so that X is a graph whose vertices are isotopy classes of essential, non-parallel, nonperipheral, pairwise disjoint simple closed curves in S (also called *curve systems*) and two distinct vertices are joined by an edge if the corresponding curve systems can be realized simultaneously by pairwise disjoint curves. In certain sporadic cases X as defined above is 0-dimensional (this happens when there are no curve systems consisting of two curves, i.e. when $g = 0, p \leq 4$ and when $g = 1, p \leq 1$). In the theorem below these cases are excluded. The mapping class group $Mod(S)$ acts on X by $f \cdot a = f(a)$.

H. Masur and Y. Minsky proved the following remarkable result.

Theorem 1 [9] *The curve complex X is δ -hyperbolic. An element of $Mod(S)$ acts hyperbolically on X if and only if it is pseudo-Anosov.*

Using their result, we show the following theorem. H. Endo and D. Kotschick [2] have shown using 4-manifold topology and Seiberg-Witten invariants that $\widetilde{QH}(Mod(S)) \neq 0$ when S is hyperbolic.

Theorem 2 [3] *Let G be a subgroup of $Mod(S)$ which is not virtually abelian. Then $\dim \widetilde{QH}(G) = \infty$.*

The following is a version of superrigidity for mapping class groups. It was conjectured by N.V. Ivanov and proved by Kaimanovich and Masur [8] in the case when the image group contains independent pseudo-Anosov homeomorphisms and it was extended to the general case by Farb and Masur [4] using the classification of subgroups of $Mod(S)$ as above. Our proof is different in that it does not use random walks on mapping class groups, but instead uses the work of M. Burger and N. Monod [1] on bounded cohomology of lattices.

Corollary 3 [3] *Let Γ be an irreducible lattice in a connected semi-simple Lie group G with no compact factors and of rank > 1 . Then every homomorphism $\Gamma \rightarrow Mod(S)$ has finite image.*

参考文献

- [1] M. Burger and N. Monod. Bounded cohomology of lattices in higher rank Lie groups. *J. Eur. Math. Soc. (JEMS)*, 1(2):199–235, 1999.
- [2] H. Endo and D. Kotschick. Bounded cohomology and non-uniform perfection of mapping class groups. Preprint, October 2000, <http://front.math.ucdavis.edu/math.GT/0010300>.
- [3] K. Fujiwara and M. Bestvina. Bounded cohomology of subgroups of mapping class groups. Preprint, December 2000, <http://front.math.ucdavis.edu/math.GT/0012115>.
- [4] Benson Farb and Howard Masur. Superrigidity and mapping class groups. *Topology*, 37(6):1169–1176, 1998.

- [5] John L. Harer. Stability of the homology of the mapping class groups of orientable surfaces. *Ann. of Math. (2)*, 121(2):215–249, 1985.
- [6] John L. Harer. The virtual cohomological dimension of the mapping class group of an orientable surface. *Invent. Math.*, 84(1):157–176, 1986.
- [7] W. J. Harvey. Boundary structure of the modular group. In *Riemann surfaces and related topics: Proceedings of the 1978 Stony Brook Conference (State Univ. New York, Stony Brook, N.Y., 1978)*, pages 245–251, Princeton, N.J., 1981. Princeton Univ. Press.
- [8] Vadim A. Kaimanovich and Howard Masur. The Poisson boundary of the mapping class group. *Invent. Math.*, 125(2):221–264, 1996.
- [9] Howard A. Masur and Yair N. Minsky. Geometry of the complex of curves. I. Hyperbolicity. *Invent. Math.*, 138(1):103–149, 1999.