

Viral nature of the aquatic ecosystems

Authors and affiliations

Daichi Morimoto¹, Kento Tominaga¹, Hiroaki Takebe¹, Sigita Šulčius², Takashi Yoshida¹

¹Graduate School of Agriculture, Kyoto University, Kitashirakawa-Oiwake, Sakyo-ku, Kyoto 606-8502, Japan

²Laboratory of Algology and Microbial Ecology, Nature Research Centre, Akademijos 2, 08412 Vilnius, Lithuania

Correspondence

Takashi Yoshida

Tel: +81-75-753-6217

Fax: +81-75-753-6226

E-mail: yoshiten@kais.kyoto-u.ac.jp

Abstract

Viruses infecting microorganisms are ubiquitous and highly abundant in aquatic environments. They considerably affect the dynamics, diversity, and evolution of their host microorganisms. In this review, we discuss the ecological implications of viruses from the perspectives of the biogeochemical cycles, microbial diversity, and virus–host coevolutionary dynamics in aquatic environments. Generally, viruses redirect host metabolism toward reproduction through molecular host–virus interactions characterized by the compositional and stoichiometric changes in intracellular metabolites, which are eventually released into the environment when the infected host cells are lysed, thus also changing the chemical composition of the water. Therefore, the modulation of metabolite biosynthesis and promotion of their recycling are major viral functions. Viruses also maintain microbial community diversity via increased infection and lysis rates of the dominant taxa and genotypes in a frequency-dependent manner, thereby allowing the co-existence of members with various competitive abilities. Finally, viruses can expand their own genotypic diversity and that of the host through complex defense and counter-defense interactions, including loss of host fitness due to the cost of resistance and the possible need for antiviral defense-specific (e.g., intra- vs. extracellular) changes in the hosts genome diversification. Continuous interactions drive

the coevolution of hosts and viruses, thereby increasing both the host and viral micro-diversity. Hence, these fundamental functions are viral “raison d’etre” and are essential for the functioning of aquatic ecosystems and its components.

Keywords

Virus-host interactions, Biogeochemical cycle, Microbial diversity, Coevolution, Lytic infection, Lysogenic infection

1.1 Introduction

Viruses infecting microorganisms are ubiquitous and abundant in aquatic ecosystems (Suttle 2005, 2007). They typically are small particles (generally 20–200 nm in length) comprised of nucleic acids (single- or double-stranded DNA or RNA) and structural proteins and have no intrinsic metabolism. Thus, their reproduction depends entirely on host cellular metabolism and replication machinery. Viral reproduction can be classified as lytic or lysogenic (Guttman et al. 2004). During lytic infection, viruses inject their genomes into host microorganisms, redirect host metabolism for efficient viral genomic nucleic acid replication and protein synthesis, and are finally released through host cell lysis (Guttman et al. 2004). In contrast, in lysogenic infection, the viral genome is integrated into the host genome as a provirus (also called prophage if the

virus integrates into the bacterial chromosome) and is propagated vertically within the host lineage until the induction of the lytic cycle under specific conditions (e.g., depending on host cell density or environmental conditions) (Howard-Varona et al. 2017a).

Both types of viral infections, lytic and lysogenic, have great potential to affect microbial communities in aquatic ecosystems. For example, viral-mediated cell lysis releases nutrients and organic matter from cells to the environment, thus stimulating biogeochemical cycling (Fuhrman 1999; Suttle 2005, 2007). In addition, viruses affect host microbial diversity in at least three different ways (Marston et al. 2012). First, viruses contribute to the maintenance of host microbial diversity by frequency-dependent infection, often seeming to have a greater effect upon those microbial taxa and genotypes that either are highly abundant or most metabolically active in the environment (Thingstad 2000). Second, viruses increase host genetic diversity via the reciprocal co-evolution of host resistance and viral infectivity (Buckling and Rainey 2002a). Lastly, viruses affect the genomic evolution and the fitness of microbial hosts through horizontal gene transfer (HGT) including the presence and movement of auxiliary metabolic genes (Breitbart et al. 2007; Hurwitz and U'Ren 2016), generalized transduction during lytic infection (Thierauf et al. 2009; Touchon et

al. 2017), and specialized transduction during lysogenic infection (Gottesman and Yarmolinsky 1968; Fernandes et al. 1989; Campos et al. 2003; Touchon et al. 2017).

The ever-growing number of culturable viral isolates, together with recent advances in sequencing technologies and bioinformatics, provides us with a deeper understanding of the viral nature in aquatic ecosystems. This chapter summarizes current understanding of viral effects on biogeochemical cycles, microbial diversity, and virus–host evolutionary dynamics in aquatic ecosystems. The viral role in the promotion of HGT that affects host genomic evolution and fitness has been reviewed extensively elsewhere and is therefore not discussed here (e.g. Balcázar 2018; Yoshida et al. 2019; Chapter E and therein).

1.2 Viral influence on biogeochemical cycle


1.2.1 Viral modulation on patterns of geochemical cycling in the ocean

In the ocean, approximately 10^{29} cells of different microorganisms form the basis of the marine food web (Whitman et al. 1998). Photosynthetic eukaryotes and prokaryotes contribute to up to 50% of the total net primary production on Earth (Field et al. 1998). Approximately half of the resultingly fixed carbon is released into the environment, re-mineralized by heterotrophic prokaryotes, and then incorporated into higher trophic levels of the aquatic food web (Azam et al. 1983). This process of

recycling of photosynthetic products is called “microbial loop” and is an essential pathway of biogeochemical cycling in the ocean (Azam et al. 1983).

Viruses outnumber prokaryotes, and up to 20% of marine microorganisms are thought to be infected and lysed by viruses daily (according to a certain view in Suttle 2007). Lysis of infected cells leads to the release of organic matter and nutrients, which would otherwise be incorporated into higher trophic levels by grazing (Fuhrman 1999; Wilhelm and Suttle 1999). This pathway of carbon flux regulated by viruses is called “viral shunt” (Wilhelm and Suttle 1999) (Fig. 1.2.1). Calculations based on the presumed microbial biomass, its turnover rates, and the predicted daily lysis suggest that viral shunts are responsible for the release of approximately 25% of primary production in the surface ocean, which amounts to up to 3 gigatons of carbon into the oceans per year (Wilhelm and Suttle 1999; Suttle 2005). However, the quantification of virus-mediated carbon flux in natural environments remains challenging owing to methodological limitations and ecosystem complexity. Recently, more advanced nutrient–phytoplankton–zooplankton (NPZ) models, including heterotrophic bacteria and viruses, have proposed that viral shunts accelerate organic matter recycling and increase net primary productivity while reducing transfer to higher trophic levels (Weitz et al. 2015). Moreover, the effect of virus-mediated carbon flux may depend on the

110 trophic status of the system and the limiting nutrients (Pourtois et al. 2020).



112 **Fig 1.2.1. Overview of virus-mediated biogeochemical cycling in aquatic microbial**

113 **ecosystem.** When viruses lyse their host, intracellular organic matter is released from

114 host cells to the particulate organic matter (POM) and dissolved organic matter (DOM)

115 pools, a process that has been termed a Viral shunt. This process is accompanied by

116 compositional and stoichiometric changes in chemical properties of intracellular

117 metabolites via viral metabolic redirection, i.e., hijacking host transcription-translation

118 systems and expression of viral auxiliary metabolic genes (AMG). Viral particles are

119 also a source of phosphorus-rich DOM (compared with host debris) and consumed by

120 direct grazing. Viral infection facilitates carbon export to the deeper layer (biological

121 pump) through particle aggregation driven by the release of lysis products and

virus-induced alterations in host physiology (Viral shuttle).

On the other hand, viruses can also contribute to carbon removal from the surface ocean (Weinbauer 2004; Sullivan et al. 2017; Laber et al. 2018). Cellular debris, including cytoplasmic material and components of the cell wall, released via viral lysis can easily aggregate and sink to the deeper layers leading to carbon sequestration (Weinbauer 2004; Laber et al. 2018). This alternative viral influence on the geochemical flux that promotes the biological carbon pump is referred to as the “viral shuttle” (Sullivan et al. 2017) (Fig. 1.2.1). For example, virus-induced carbon transportation has been extensively studied in *Emiliania huxleyi* (haptophyte) and its known virus (Laber et al. 2018). Several laboratory studies have reported that viral infection stimulates the production of transparent exopolymer particles (TEP), which increases the stickiness of cells, thus promoting aggregation (Rosenwasser et al. 2014; Nissimov et al. 2018). Field studies monitoring *E. huxleyi* blooms in the North Atlantic have revealed that TEP concentrations increase during the early stages of infection and that infected cells are preferentially transported to the deep ocean (Sheyn et al. 2018). Considering the huge abundance of *E. huxleyi* (reaching 10^7 cells/mL during their bloom period; Silkin et al. 2020), the virus-mediated sinking of its cells would thus have a significant impact on the available carbon in the surface ocean. Moreover, laboratory experiments have

reported that the virus-infected culture of *Chaetoceros tenuissimus* (Diatomea) is up to 59-fold enriched in particulate organic matter compared with uninfected controls (Yamada et al. 2018). At the global scale, a metagenomic study based on data from the Tara Ocean has suggested that the infection and lysis of the widespread and abundant cyanobacteria *Synechococcus* significantly contribute to carbon export compared with other microorganisms (Guidi et al. 2016). Further studies using quantitative methods are required for a better understanding of the effect of viral shuttles in the global ocean.

In addition, viral particles themselves contribute to the biogeochemical cycling of carbon and other nutrients. For example, approximately 0.03 Gt C per year (Bar-On and Milo 2019), and most of the viral biomass is attributed to the dissolved organic matter (DOM) fraction ($< 0.45 \mu\text{m}$) due to the size of the virion (e.g., bacterial viruses generally range between 20 and 200 nm) (Zsolnay 2003; Leenheer and Croué 2003; Findlay and Parr 2017). Recently, a relatively large number of marine viruses have been identified to attach to non-host organisms and particles (Yamada et al. 2020). Thus, if or when these non-host organisms or particles are predated, their attached viruses indirectly contribute to classical marine food webs even when they are not infecting any organisms. Altogether, viral particles could contribute to both DOM and the particulate organic matter (POM) pools in the ocean.

A previous study investigating the elemental composition of both virus particles and viral lysates eluted from their host debris revealed that viral lysates tend to be more depleted in phosphorus than are uninfected cells because the amount of genomic nucleic acids contained in progeny viral particles results in those viruses being relatively phosphorus-rich as compared with their amounts of carbon and nitrogen (Jover et al. 2014). By extrapolating this model to the ecosystem scale, marine viruses are predicted to constitute a comparatively high proportion (> 5%) of the total DOP pool in the surface ocean (Jover et al. 2014). Thus, viruses themselves can be regarded as an abundant nutrient source.

Predation of viral particles by predators has been demonstrated in several studies using culture experiments (Suttle and Chen 1992; Bettarel et al. 2005; Lawrence et al. 2018; Welsh et al. 2020). For example, a co-cultivation study exposing *Phaeocystis globosa* and its virus to various predators has demonstrated that viruses can be effectively removed (up to 98% within 24 h) from the water column by non-host organisms, including sea anemones, polychaete larvae, sea squirts, crabs, cockles, oysters, and sponges (Welsh et al. 2020). Therefore, although the rate of viral particle removal by aquatic protists may vary depending on both the virus and predator strains (Suttle and Chen 1992; Gonzalez and Suttle 1993; Lawrence et al. 2018; Welsh et al.

2020), the ingestion of viruses can serve as a possible source of nutrients, especially phosphorus.

These viral influences on the biogeochemical cycles fluctuate across short- and long-time scales. On a long-term scale, for instance, the microbial community exhibits seasonal compositional changes (Cram et al. 2015; Parada and Fuhrman 2017; Needham et al. 2018; Choi et al. 2020), which are followed by the seasonal dynamics of their viruses (Needham et al. 2017; Ignacio-Espinoza et al. 2020). Furthermore, diverse taxa of photosynthetic microorganisms and even some heterotrophic ones show diel activity in culture and environmental studies, which is partly explained by the viral infection cycle (Morimoto et al. 2020 and references therein). Both the seasonality and the diel cycle activity of microorganisms and their viruses suggest that host–virus interactions could generate temporal fluctuations in geochemical cycles in aquatic environments.

1.2.2 Virus–host interactions-mediated modification of host cell metabolism

Viruses switch their host metabolism from cellular replication to progeny production. Compared with non-infected hosts, metabolically reprogrammed cells can be generally distinguished based on changes in the host transcription program,

eventually leading to distinctiveness in the proportion of end-point products between infected and uninfected cells (Ankrah et al. 2014; Jover et al. 2014; Rosenwasser et al. 2014; Ma et al. 2018) (Fig. 1.2.1). For instance, disproportioning of phosphorus between infected and uninfected hosts (as discussed in the previous section) could be attributed to a viral reprogramming mechanism in which viruses degrade host DNA and utilize the resultant nucleic acids for the synthesis of viral DNA (Wikner et al. 1993; Kutter et al. 2018).

Currently, cell metabolic reprogramming by viruses has been studied using both transcriptomic and metabolomic analyses and is found to be a highly regulated process. For instance, the infection strategy of T4-like viruses follows the three temporal expression classes of early, middle, and late genes, corresponding to host takeover, replication, and virion morphogenesis, respectively, and occurs in accordance with the downregulation of genes related to host replication (Roucourt and Lavigne 2009). Such transcriptional regulation by viruses has also been investigated in several lineages of marine and freshwater prokaryotic as well as eukaryotic phytoplankton (Lindell et al. 2007; Rosenwasser et al. 2014; Bachy et al. 2018; Moniruzzaman et al. 2018; Morimoto et al. 2018; Ku et al. 2020) and heterotrophic bacteria (Ankrah et al. 2014; Howard-Varona et al. 2017b, 2020). The metabolic regulation of host cells may

also depend on host taxonomy and thus differ between various species (e.g., cyanoviruses with broad and narrow host range seem to have different infection strategies; *E. huxleyi* virus EhV possesses five gene expression phases during infection) (Lindell et al. 2005; Clokie et al. 2006; Doron et al. 2016; Morimoto et al. 2018; Ku et al. 2020). In addition, it may be influenced by host physiological states, which in turn depends on nutrient availability (e.g. phosphate) (Kelly et al. 2013; Lin et al. 2016; Bachy et al. 2018). Thus, viruses could affect the proportion of end-point products in the infected cells while those hosts either directly or indirectly are responding to environmental conditions.

Viruses can also possess host-derived genes (also called as auxiliary metabolic genes, AMGs) that are expressed during infection, thus altering host metabolism, and increasing the efficiency of viral reproduction (Breitbart et al. 2007; Hurwitz and U'Ren 2016). These AMGs can largely be classified into two classes (Class I and II) based on their function according to the Kyoto Encyclopedia of Genes and Genomes database (Hurwitz and U'Ren 2016). Viral-encoded AMGs not only maintain cellular functions necessary for viral DNA replication and virion production (e.g., ATP production and nucleotide synthesis) during infection (Lindell et al. 2004, 2005), but also both down and up-regulate a range of targeted metabolic pathways that can substantially alter cell

stoichiometry and nutrient metabolism (De Smet et al. 2016, 2017). Most AMG that have been identified to date are directly involved in either the utilization and uptake of limiting nutrients or energy production (Enav et al. 2014; Hurwitz and U'Ren 2016), which in turn may have (at least temporarily) a positive feedback effect on the host cell by improving its fitness during infection (Zeng and Chisholm 2012): the acquisition and metabolism of carbon (e.g., *psbA* and *psbD*; Lindell et al. 2004, 2005; Thompson et al. 2011), nitrogen (e.g., *amt*; Monier et al. 2017), and phosphorus (e.g., *pstS* and *phoA*; Zeng and Chisholm 2012). New putative AMGs are continuously being discovered in bacterial (Breitbart 2011; Crummett et al. 2016; Breitbart et al. 2018; Warwick-Dugdale et al. 2019), eukaryotic (Schvarcz and Steward 2018; Needham et al. 2019), and archaea viruses (Ahlgren et al. 2019) or the more broadly defined environmental viromes (Williamson et al. 2008; Anantharaman et al. 2014; Hurwitz et al. 2015; Moniruzzaman et al. 2020; Schulz et al. 2020; Kieft et al. 2020). Thus, considering that viral-encoded AMGs are abundant and widespread in aquatic environments (Williamson et al. 2008), AMG-mediated metabolic reprogramming can substantially contribute to major biogeochemical cycles and ecosystem functioning at the global scale (Sieradzki et al. 2019).

Indeed, changes in metabolites mediated by virus–host interactions have been

observed in both eukaryotes and prokaryotes using a metabolomic approach (Rosenwasser et al. 2014; Ma et al. 2018). For example, *E. huxleyi* virus EhV downregulates host *de novo* sphingolipid genes and simultaneously promotes the induction of a viral-encoded homologous pathway, resulting in the metabolic shift toward viral sphingolipid production (Rosenwasser et al. 2014). A recent culture-based study on *Synechococcus* and its viruses revealed that the composition of chemical compounds (proteins, carbohydrates, and lipids) of organic matter differs between infected and uninfected cells (Ma et al. 2018). Such compositional changes induced by viruses have been detected during viral infection of *Sulfitobacter* (C:N ratio of host cell shifted to nitrogen-rich state compared with uninfected cells) (Ankrah et al. 2014).

Thus, from an ecological perspective, the virus-induced metabolic reprogramming depends on both the virus-host pair (and therefore on the diversity and composition of microbial assemblages) and the host physiological state during infection and can modulate the generation as well as the diversity of end-point products, thus leading to altered biogeochemical cycling. It is also speculated that halting the viral reproduction at various infection stages using antiviral responses, including signal transduction, cell cycle regulation (Moniruzzaman et al. 2018), and metabolic pathway (Rosenwasser et al. 2014), might establish metabolite diversity in the infected cells via

the generation of intermediate products of viral progeny (Zborowsky and Lindell 2019)
or unusual proteins in the uninfected cells.

1.3 Viral infection shaping microbial community diversity

1.3.1 Contribution of lytic infection to microbial diversity maintenance

In aquatic ecosystems, diverse microorganisms compete for nutrient resources
but can co-exist and account for a large proportion of total aquatic biodiversity, a
concept that has been known as the classical question “paradox of the plankton”
(Hutchinson 1961). Viruses are currently thought to contribute to the co-existence of
microbial species and genotypes.

Basically, viruses are believed to infect their specific microbial hosts in a
frequency-dependent manner (Fuhrman and Suttle 1993). Therefore, viral infection
checkmates microbial species that become dominant through the competition among
co-existing microorganisms that possess different substrate affinity, and thereby enables
the co-existence of multiple competing microbial species (“Kill the Winner” hypothesis)
(Thingstad 2000) (Fig. 1.3.1). Indeed, several culture and environmental studies have
demonstrated that viral top-down control modulates microbial abundance, which is
consistent with the results expected from a Kill the Winner hypothesis (Tarutani et al.
2000; Schwalbach et al. 2004; Bouvier and Del Giorgio 2007; Yoshida et al. 2008a;

Rodriguez-Brito et al. 2010; Kuno et al. 2012; Parsons et al. 2012; Kimura et al. 2013; Needham et al. 2013; Cram et al. 2016).

The viral-mediated co-existence mechanism, by which viruses are expected to affect host diversity in a frequency dependent manner according to the Kill the Winner hypothesis, could provide a mechanism that explains the continuing co-existence of diverse genotypes within a single microbial species (or closely related lineage) rather than the co-existence of diverse microbial species. Conventionally, the philosophy has been that phenotypic and genotypic diversity within a microbial population was expected to become homogenized to a greater level of fitness in the environment. The microbial population (species or closely related lineage) that are genetically cohesive and ecologically distinct are called an “ecotype” (ecotype hypothesis) (Maharjan et al. 2006; Cohan and Koeppel 2008). The ecotype had been believed to be periodically replaced as fitter ecotypes emerged after profitable mutation or preferable environmental changes (Maharjan et al. 2006; Cohan and Koeppel 2008). However, metagenome sequence alignment analyses have demonstrated that several genomic regions (metagenomic islands; MGIs) are underrepresented even within a single microbial population in similar environments (Coleman and Chisholm 2007; Cuadros-Orellana et al. 2007; Kettler et al. 2007; Wilhelm et al. 2007; Frias-Lopez et al.

2008; Rodriguez-Valera et al. 2009; Rodriguez-Valera and Ussery 2012). This suggests that diverse genotypes coexist within a single microbial population. Furthermore, these MGIs include diverse accessory genes, such as extracellular structure-related genes that can be viral recognition sites (Reva and Tümmler 2008; Sharma et al. 2008; Rodriguez-Valera et al. 2009; Rodriguez-Valera and Ussery 2012) and antiviral response-related genes such as CRISPR in addition to genes that affect restriction-modification (Sorek et al. 2008; Wilmes et al. 2009). Therefore, MGIs are thought to play an important role in the ability of hosts to escape or survive from viral infection, in which host genotypic diversity is driven by viral predation pressure, thereby leading to the co-existence of multiple competing microbial genotypes (“Constant Diversity dynamics” model) (Rodriguez-Valera et al. 2009) (Fig. 1.3.1). Indeed, multiple genotypes of *Microcystis aeruginosa* possessing different CRISPR arrays and its virus Ma-LMM01 coexist and oscillate during the massive bloom of this nuisance and toxic species (Kuno et al. 2012, 2014; Kimura et al. 2013). Hence, lytic viruses play important roles in not only the co-existence of microbial species but also the maintenance of high diversity within a single microbial population.




Fig 1.3.1. Schematic diagram of mechanisms in virus-mediated maintenance of microbial community diversity. Preferential viral infection of abundant species enables the co-existence of diverse competing microbial species by preventing the dominance of only few species (“Kill the Winner” hypothesis). Similar viral top-down control is proposed as a mechanism to maintain high genotypic diversity within a single microbial population (“Constant Diversity dynamics” model). Lastly, the prevalence of lysogeny in dominant microbial population may be another potential mechanism that allows abundant host species to be dominant by taking advantages benefit from lysogenic conversion such as superinfection exclusion (“Piggyback-the-Winner” model).

1.3.2 Potential contribution of lysogenic infection to microbial diversity maintenance

So far, we have focused on lytic viruses and described their contribution to the maintenance of microbial diversity in aquatic environments. Another key question related to viral impact on microbial diversity is whether temperate viruses also contribute to shaping microbial community diversity.

Provirus integration can occur either through repeated random transposition events or at specific integration sites (e.g., host tRNA genes; also called site-specific recombination) and is associated with the immediate transcriptional suppression (e.g. via specific virus repressors) of lytic promoters and genes associated with virion production (Casjens and Hendrix 2015). This mode of viral infection that generates lysogenic cells (Hobbs and Abedon 2016) is considered as an adaptive strategy of temperate viruses to ensure their persistence in the environment, in which novel phenotypic or metabolic advantages are sometimes conferred to host microorganisms via concomitant effects by mechanisms such as HGT (Hendrix et al. 2000; Howard-Varona et al. 2017a), the capability of up and down-regulation of host genes (Argov et al. 2017), and integration-driven gene disruption (Feiner et al. 2015).

Although one study estimated that approximately half of 100 marine bacterial isolates harbored temperate viruses in their genomes (Paul 2008), proviruses are rarely

found in the marine-dominant bacterial lineages (e.g., only one provirus was recently reported in SAR11 clades; see Morris et al. 2020) possibly due to genome streamlining, in which a bacterial genome is minimized to a highly constrained gene set that confers maximum fitness (Touchon et al. 2016). Therefore, the ecological significance of lysogenic infection on microbial diversity maintenance in marine ecosystems remains under debate. Traditionally, it was believed that bacterial viruses control their host abundances in a frequency-dependent manner as described above, and thus viral abundance is typically 10-folds higher than that of prokaryotes (Wommack and Colwell 2000; Weinbauer 2004). Therefore, lysogenic infection is presumed to be the preferred viral strategy under conditions of reduced host cell number and activity (Stewart and Levin 1984; Sime-Ngando 2014; Brum et al. 2016). However, viral metagenomic and metadata approaches have revealed that viral particles are relatively less abundant at high microbial densities (Knowles et al. 2016; Wigington et al. 2016). Likewise, it was demonstrated that the virus/host genome abundance ratio was negatively correlated with the host abundance at the genus or phylum levels (Coutinho et al. 2017). Additionally, the relative abundance of hallmark genes encoded by temperate viruses increased with microbial density in a coral reef (Knowles et al. 2016). These findings suggest that lysogenic infection may become dominant at high-cell densities because proviruses can

replicate quickly in a way that will keep pace with their fast-growing host and provirus-mediated superinfection resistance might become increasingly important at high cell densities (called “Piggyback-the-Winner” model) (Knowles et al. 2016, 2017; Coutinho et al. 2017) (Fig. 1.3.1).

Footnotes

Paradox of the plankton: The concept arguing paradoxical situation of coexistence of various plankton species competing for identical resources in homogeneous and resource limited environment (Hutchinson 1961).

Kill the Winner hypothesis: A model proposing the dynamics of virus-host interactions in which an increase of host population (winner) is accompanied with increasing of its infectious viruses, and thereby viruses prevent their hosts from becoming dominant through increased mortality of the winner (Thingstad 2000).

Constant Diversity dynamics model: A hypothetical model proposing that the diversity of prokaryotic populations is maintained by viral predation, because the best-adapted populations are selected by viral predation. The hypothesis assumes that each microbial population has distinct viral receptor (Rodriguez-Valera et al. 2009).

Piggyback-the-Winner model: A hypothetical model proposing that lysogeny would be

388 favored when a bacterial host is dominated in the environment because a provirus can
389 replicate rapidly together with host DNA replication (Knowles et al. 2016).

391 **1.4 Evolutionary roles of viruses that generate genotype-level microbial** 392 **diversification**

393 In short-term laboratory experiments, host–virus coevolution appeared to be
394 suppressed by the emergence of a viral-resistant genotype, which the virus could not
395 evolve to overcome (Dennehy 2012). In particular, *de novo* mutations that cause
396 changes in those bacterial cell-surface structures which serve as viral receptor sites are
397 one of the major factors that prevent viral attachment, and thereby confer the potential
398 host with resistance against viral infection (Lenski and Levin 1985). Therefore,
399 host–virus coevolution has been considered to be constrained by the asymmetry of
400 evolutionary potential between hosts and viruses (Cannon et al. 1971; Cowlshaw and
401 Mrsa 1975; Barnet et al. 1981; Lenski and Levin 1985; Waterbury and Valois 1993;
402 Middelboe et al. 2001; Wei et al. 2010, 2011).

403 On the other hand, long-term evolutionary laboratory studies have indicated
404 that the host and its virus undergo persistent coevolution over a prolonged period, as
405 evidenced with the soil bacterium *Pseudomonas fluorescens* (Buckling and Rainey
406 2002a, b; Brockhurst et al. 2007; Hall et al. 2011a, b) and *E. coli* (Mizoguchi et al.

2003). Similar co-evolutionary dynamics have also been observed in the marine bacteria *Prochlorococcus* (Avrani et al. 2011), *Synechococcus* (Marston et al. 2012), and *Cellulophaga baltica* (Middelboe et al. 2009).

The above-mentioned co-evolution scenario, based on the one-virus-to-one-bacteria relationship, predicts that the genetic contents of both bacteria and virus would converge over repeated interactions. As described above, however, genotypic diversity is observed in MGIs even within a single microbial population (Coleman and Chisholm 2007; Cuadros-Orellana et al. 2007; Kettler et al. 2007; Wilhelm et al. 2007; Frias-Lopez et al. 2008; Rodriguez-Valera et al. 2009; Rodriguez-Valera and Ussery 2012). Additionally, under-represented genomic regions have been found in marine viruses (metaviromic islands) with a large fraction of their identified genes (e.g., 59 out of 138) associated with host recognition of viruses (Mizuno et al. 2014). Thus, the recent understanding of host–virus coevolution is based on multiple genotype (strain)-level interactions within both the microbial host and viral species which seem to represent defense and counter-defense strategies (e.g., CRISPR and mutation in protospacer) or resistance and overcome of resistance (e.g., mutation in cell surface and viral tail gene) (Fig. 1.4.1).




Fig. 1.4.1 Proposed mechanisms in host pan-genome expansion via virus–host interactions. (a) Under the viral top-down control toward dominant species and genotypes, “Red Queen” like host–virus co-diversification can be established in abundant host and abundant viruses. Continuous arms race via intracellular defense (e.g., CRISPR-Cas system) and extracellular resistance (e.g., viral recognition sites) plays a part in genomic diversification of both host and virus. (b) A complicated balance between trade-off and specificities of each defense mechanism can affect the arms race. For example, extracellular resistance (e.g., surface modification) is preferable to intercellular defense (e.g., CRISPR-Cas) under conditions of high viral genetic diversity

because the former promotes resistance toward against broader viral genotypes. However, because extracellular resistance may result in a higher cost, such as the impairment of nutrient uptake ability, intercellular defense is possibly preferable under competitive situations with host competitors.

In freshwater ecosystems, host-virus coevolution have been intensively studied in the bloom-forming cyanobacterium *M. aeruginosa* and its viruses (Yoshida et al. 2005, 2008a, b; Kimura et al. 2012, 2013, 2018; Kuno et al. 2012, 2014; Yoshida-Takashima et al. 2012; Morimoto et al. 2019). Interestingly, the most abundant *Microcystis* CRISPR genotype is known to coexist with that derived by novel spacer acquisitions from cyanoviruses in the environment. This finding suggested that both abundant host and viral genotypes have diversified in the bloom without a complete selective sweep (Kimura et al. 2018). Thus, the Red Queen like dynamics could be established, to some extent, between the abundant host genotype and its cyanoviruses under high viral contact rate; with reciprocal adaptation via defense and counter-defense continuously occurring in multiple-to-multiple relationships (Koskella and Brockhurst 2014), thus subsequently increasing the diversity of host organisms and viruses (Fig. 1.4.1). Also, a recent metagenomic survey revealed the co-existence of highly host-specific (narrow host range) and broad host range *Microcystis* viruses and the high

co-expression of antiviral defense and viral genes in the environment (Morimoto et al. 2019). Considering that they often induce antiviral responses, broad host range viruses might be important for host genotype diversification (Morimoto et al. 2019).

In contrast with freshwater cyanobacteria typically having the greatest overall numbers of defense genes (Makarova et al. 2011), marine prokaryotes rarely possess distinctive defense genes such as CRISPR-Cas due to their genome streamlining (Touchon et al. 2016). However, instances of co-existence between dominant marine prokaryotes and their viruses were observed, which are presumably sustained by Red Queen-like co-evolution dynamics. For example, the dominance of the SAR11 clade bacteria and their viruses was predicted to be maintained by host rapid adaptation to viruses. The rapid adaptation was achieved by high recombination rates among SAR11 in their variable genomic region that encoded genes involved in synthesis of cell surface proteins, and this hypothesis is possibly supported by their high host cell density (King of the Mountain hypothesis) (Zhao et al. 2013). Furthermore, constant turnover of single-nucleotide polymorphism variants in relatively abundant marine viruses also suggests that Red Queen-like virus–host coexistence could be established by perpetually changing minor variants.

As described in this section, one of the major forces driving host and viral

genotypic diversification is the reciprocal defense and counter-defense that occur through extracellular (e.g., *de novo* mutation of cell-surface structure) and intracellular resistance (e.g., acquisition of novel CRISPR spacers). Comparative antiviral resistance analyses in *Synechococcus* and *Prochlorococcus* provided new insights into host-favored antiviral defenses in narrow and broad host range viruses (Doron et al. 2016; Zborowsky and Lindell 2019). Host cyanobacteria resist against narrow host range viruses irrespective of the viral family by preventing viral entry into the cell, whereas intracellular resistance arrests the infection cycle of broad host range viruses at various infection stages (Doron et al. 2016; Zborowsky and Lindell 2019) (Fig. 1.4.1). These differences in antiviral responses that seemingly occur according to viral host range are speculated to be associated with fitness trade-offs in extracellular and intracellular antiviral responses. In extracellular antiviral responses, mutations in cellular surface structures can impair nutrient uptake and utilization but can protect against diverse viral attacks, leading to an increase in host fitness trade-off favoring the host (Winter et al. 2010; Avrani et al. 2012) (Fig. 1.4.1). On the other hand, choosing to modify intracellular antiviral responses techniques may be energetically favorable, especially the CRISPR-Cas system, because the cost of a new spacer acquisition is speculated to be low, although it is possible that additional types of resistance costs may

exist (Thingstad et al. 2014) (Fig. 1.4.1). Indeed, recent studies focusing on the differences in biotic complexity between *in vitro* and environments have revealed that coexistence among human pathogens amplified the fitness trade-offs associated with viral receptor mutations in *Pseudomonas aeruginosa* and therefore enhanced the evolution of CRISPR-based resistance (Alseth et al. 2019). Higher viral genetic diversity can also influence CRISPR-based evolution, with an example being that the majority of a *Pseudomonas* population more favorably evolved based on the mutation of viral receptors to resist a broader range of viral genotypes than CRISPR-based specific resistance (Broniewski et al. 2020). Thus, fitness trade-offs in bacterial host species and diversity of both bacterial host and viruses in the environment could be another important factor that affects host–virus coevolution. From the perspective of fitness trade-off, most recently, the “pan-immune system” concept has been proposed. This states that a single strain can access immune defense mechanisms in closely related strain via HGT, although it cannot possess all possible defense systems (Bernheim and Sorek 2020).

Footnotes

Red Queen like dynamics: A hypothesis proposing co-evolutionary process between

competing species (Valen 1973); in the case of virus-host interactions, this hypothesis explains continuous dynamics of resistance acquisition in microbial hosts and viral avoidance to the host resistance (Brockhurst et al. 2014).

King of the Mountain hypothesis: A hypothesis proposing that high recombination rate enables dominance of a competitive prokaryote in the ecosystem via horizontal transfer of genes involved in resistance to viral infection (Zhao et al. 2013).

1.5 Conclusion

Viruses, which are highly abundant biological entities lacking their own metabolism, can reprogram host cells toward the production of virus progeny, after which the host cell is lysed, releasing new virions into the surrounding environment. This reprogramming viral strategy can change the content and composition of host metabolites and releases a large amount of organic matter, thus considerably affecting biogeochemical cycles. In addition, viral infection checkmates not only microbial species that become dominant but also abundant genotypes within a single microbial species, and thereby enables the coexistence of diverse microbial species and genotypes in the aquatic ecosystem. Thus, viral infection is one of the key factors shaping microbial community diversity and maintaining high diversity within a single microbial population. Likewise, because temperate viruses may be prevalent under specific

combinations of environment and conditions, they could affect microbial diversity via superinfection exclusion during either the lysogenic cycle or induction of lytic cycle. Meanwhile, microorganisms have evolved extracellular and intracellular antiviral mechanisms with different fitness costs and specificities. Therefore, viruses with diverse host range and genetic diversity interact with abundant hosts by a complex balancing between fitness trade-offs and the specificity of extracellular- and intracellular antiviral resistance in hosts. Together, this results in continuous virus–host coevolution, leading to diversification in aquatic ecosystems. Collectively, viruses are important biological entities that sustain and generate microbial diversity and control the biogeochemical cycle in aquatic environments.

Acknowledgements

This work was supported by the Grants-in Aids for Scientific Research (B) (grant number 17H03850) from the Japan Society for the Promotion of Science (JSPS); the Research Council of Lithuania (grant number S-MIP-17-28 to SŠ); the Canon Foundation (grant number 203143100025); the JSPS Scientific Research on Innovative Areas (grant number 16H06437); and the Bilateral Open Partnership Joint Research Project (Japan–Lithuania Research Cooperative Program) “Research on prediction of environmental change in Baltic Sea based on comprehensive metagenomic analysis of

microbial viruses”. DM, KT, HT, SŠ and TY all contributed to manuscript preparation,
discussion, and revision.

References

Ahlgren NA, Fuchsman CA, Rocap G, Fuhrman JA (2019) Discovery of several novel,
widespread, and ecologically distinct marine Thaumarchaeota viruses that encode
amoC nitrification genes. ISME J 13:618–631.

<https://doi.org/10.1038/s41396-018-0289-4>

Alseth EO, Pursey E, Luján AM, et al (2019) Bacterial biodiversity drives the evolution
of CRISPR-based phage resistance. Nature 574:549–552.

<https://doi.org/10.1038/s41586-019-1662-9>

Anantharaman K, Duhaime MB, Breier JA, et al (2014) Sulfur Oxidation Genes in
Diverse Deep-Sea Viruses. Science (80-) 344:757 LP – 760.

<https://doi.org/10.1126/science.1252229>

Ankrah NYD, May AL, Middleton JL, et al (2014) Phage infection of an
environmentally relevant marine bacterium alters host metabolism and lysate
composition. ISME J 8:1089–1100. <https://doi.org/10.1038/ismej.2013.216>

Argov T, Azulay G, Pasechnek A, et al (2017) Temperate bacteriophages as regulators

563 of host behavior. *Curr Opin Microbiol* 38:81–87.

564 <https://doi.org/10.1016/j.mib.2017.05.002>

565 Avrani S, Schwartz DA, Lindell D (2012) Virus-host swinging party in the oceans. *Mob*

566 *Genet Elements* 2:88–95. <https://doi.org/10.4161/mge.20031>

567 Avrani S, Wurtzel O, Sharon I, et al (2011) Genomic island variability facilitates

568 *Prochlorococcus*-virus coexistence. *Nature* 474:604–608.

569 <https://doi.org/10.1038/nature10172>

570 Azam F, Fenchel T, Field JG, et al (1983) The Ecological Role of Water-Column

571 Microbes in the Sea. *Mar Ecol Prog Ser* 10:257–263

572 Bachy C, Charlesworth CJ, Chan AM, et al (2018) Transcriptional responses of the

573 marine green alga *Micromonas pusilla* and an infecting prasinovirus under

574 different phosphate conditions. *Environ Microbiol* 20:2898–2912.

575 <https://doi.org/10.1111/1462-2920.14273>

576 Balcázar JL (2018) How do bacteriophages promote antibiotic resistance in the

577 environment? *Clin Microbiol Infect* 24:447–449.

578 <https://doi.org/https://doi.org/10.1016/j.cmi.2017.10.010>

579 Bar-On YM, Milo R (2019) The Biomass Composition of the Oceans: A Blueprint of

580 Our Blue Planet. *Cell* 179:1451–1454. <https://doi.org/10.1016/j.cell.2019.11.018>

581 Barnet YM, Daft MJ, Stewart WDP (1981) Cyanobacteria-cyanophage interactions in
 582 continuous culture. *J Appl Bacteriol* 51:541–552.
 583 <https://doi.org/10.1111/j.1365-2672.1981.tb01273.x>

584 Bernheim A, Sorek R (2020) The pan-immune system of bacteria: antiviral defence as a
 585 community resource. *Nat Rev Microbiol* 18:113–119.
 586 <https://doi.org/10.1038/s41579-019-0278-2>

587 Bettarel Y, Sime-Ngando T, Bouvy M, et al (2005) Low consumption of virus-sized
 588 particles by heterotrophic nanoflagellates in two lakes of the French Massif
 589 Central . *Aquat Microb Ecol* 39:205–209

590 Bouvier T, Del Giorgio PA (2007) Key role of selective viral-induced mortality in
 591 determining marine bacterial community composition. *Environ Microbiol*
 592 9:287–297. <https://doi.org/10.1111/j.1462-2920.2006.01137.x>

593 Breitbart M (2011) Marine Viruses: Truth or Dare. *Ann Rev Mar Sci* 4:425–448.
 594 <https://doi.org/10.1146/annurev-marine-120709-142805>

595 Breitbart M, Bonnain C, Malki K, Sawaya NA (2018) Phage puppet masters of the
 596 marine microbial realm. *Nat Microbiol* 3:754–766.
 597 <https://doi.org/10.1038/s41564-018-0166-y>

598 Breitbart M, Thompson L, Suttle C, Sullivan M (2007) Exploring the vast diversity of

599 marine viruses. *Oceanography* 20:135–139.

600 <https://doi.org/10.5670/oceanog.2007.58>

601 Brockhurst MA, Chapman T, King KC, et al (2014) Running with the Red Queen: The

602 role of biotic conflicts in evolution. *Proc R Soc B Biol Sci* 281:.

603 <https://doi.org/10.1098/rspb.2014.1382>

604 Brockhurst MA, Morgan AD, Fenton A, Buckling A (2007) Experimental coevolution

605 with bacteria and phage: The *Pseudomonas fluorescens*— Φ 2 model system. *Infect*

606 *Genet Evol* 7:547–552.

607 <https://doi.org/https://doi.org/10.1016/j.meegid.2007.01.005>

608 Broniewski JM, Meaden S, Paterson S, et al (2020) The effect of phage genetic

609 diversity on bacterial resistance evolution. *ISME J* 14:828–836.

610 <https://doi.org/10.1038/s41396-019-0577-7>

611 Brum JR, Hurwitz BL, Schofield O, et al (2016) Seasonal time bombs: Dominant

612 temperate viruses affect Southern Ocean microbial dynamics. *ISME J* 10:437–449.

613 <https://doi.org/10.1038/ismej.2015.125>

614 Buckling A, Rainey PB (2002a) Antagonistic coevolution between a bacterium and a

615 bacteriophage. *Proc R Soc B Biol Sci* 269:931–936.

616 <https://doi.org/10.1098/rspb.2001.1945>

617 Buckling A, Rainey PB (2002b) The role of parasites in sympatric and allopatric host
618 diversification. *Nature* 420:496–499. <https://doi.org/10.1038/nature01164>

619 Campos J, Martínez E, Marrero K, et al (2003) Novel Type of Specialized Transduction
620 for CTX ϕ or Its Satellite Phage RS1 Mediated by Filamentous Phage V_{GJ} ϕ in
621 *Vibrio cholerae*. *J Bacteriol* 185:7231 LP – 7240.
622 <https://doi.org/10.1128/JB.185.24.7231-7240.2003>

623 Cannon RE, Shane MS, Bush VN (1971) Lysogeny of a blue-green alga, *Plectonema*
624 *boryanum*. *Virology* 45:149–153. [https://doi.org/10.1016/0042-6822\(71\)90121-8](https://doi.org/10.1016/0042-6822(71)90121-8)

625 Casjens SR, Hendrix RW (2015) Bacteriophage lambda: Early pioneer and still relevant.
626 *Virology* 479–480:310–330.
627 <https://doi.org/https://doi.org/10.1016/j.virol.2015.02.010>

628 Choi C, Jimenez V, Needham D, et al (2020) Seasonal and geographical transitions in
629 eukaryotic phytoplankton community structure in the Atlantic and Pacific Oceans.
630 *Frontiers (Boulder)* 11:. <https://doi.org/10.3389/fmicb.2020.542372>

631 Clokie MRJ, Shan J, Bailey S, et al (2006) Transcription of a “photosynthetic” T4-type
632 phage during infection of a marine cyanobacterium. *Environ Microbiol* 8:827–835.
633 <https://doi.org/10.1111/j.1462-2920.2005.00969.x>

634 Cohan FM, Koeppel AF (2008) The Origins of Ecological Diversity in Prokaryotes.

635 Curr Biol 18:1024–1034. <https://doi.org/10.1016/j.cub.2008.09.014>

636 Coleman ML, Chisholm SW (2007) Code and context: Prochlorococcus as a model for
 637 cross-scale biology. Trends Microbiol 15:398–407.
 638 <https://doi.org/10.1016/j.tim.2007.07.001>

639 Coutinho FH, Silveira CB, Gregoracci GB, et al (2017) Marine viruses discovered via
 640 metagenomics shed light on viral strategies throughout the oceans. Nat Commun
 641 8:15955. <https://doi.org/10.1038/ncomms15955>

642 Cowlishaw J, Mersa M (1975) Co-Evolution of a Virus-Alga System. Appl Microbiol
 643 29:234 LP – 239

644 Cram JA, Chow CET, Sachdeva R, et al (2015) Seasonal and interannual variability of
 645 the marine bacterioplankton community throughout the water column over ten
 646 years. ISME J 9:563–580. <https://doi.org/10.1038/ismej.2014.153>

647 Cram JA, Parada AE, Fuhrman JA (2016) Dilution reveals how viral lysis and grazing
 648 shape microbial communities. Limnol Oceanogr 61:889–905.
 649 <https://doi.org/10.1002/lno.10259>

650 Crummett LT, Puxty RJ, Weihe C, et al (2016) The genomic content and context of
 651 auxiliary metabolic genes in marine cyanomyoviruses. Virology 499:219–229.
 652 <https://doi.org/10.1016/j.virol.2016.09.016>

653 Cuadros-Orellana S, Martin-Cuadrado AB, Legault B, et al (2007) Genomic plasticity
654 in prokaryotes: The case of the square haloarchaeon. *ISME J* 1:235–245.
655 <https://doi.org/10.1038/ismej.2007.35>

656 De Smet J, Hendrix H, Blasdel BG, et al (2017) *Pseudomonas* predators: understanding
657 and exploiting phage–host interactions. *Nat Rev Microbiol* 15:517–530.
658 <https://doi.org/10.1038/nrmicro.2017.61>

659 De Smet J, Zimmermann M, Kogadeeva M, et al (2016) High coverage metabolomics
660 analysis reveals phage-specific alterations to *Pseudomonas aeruginosa* physiology
661 during infection. *ISME J* 10:1823–1835. <https://doi.org/10.1038/ismej.2016.3>

662 Dennehy JJ (2012) What can phages tell us about host-pathogen coevolution? *Int J Evol*
663 *Biol* 2012:396165. <https://doi.org/10.1155/2012/396165>

664 Doron S, Fedida A, Hernández-Prieto MA, et al (2016) Transcriptome dynamics of a
665 broad host-range cyanophage and its hosts. *ISME J* 10:1437–1455.
666 <https://doi.org/10.1007/s13398-014-0173-7.2>

667 Enav H, Mandel-Gutfreund Y, Béjà O (2014) Comparative metagenomic analyses
668 reveal viral-induced shifts of host metabolism towards nucleotide biosynthesis.
669 *Microbiome* 2:1–11. <https://doi.org/10.1186/2049-2618-2-9>

670 Feiner R, Argov T, Rabinovich L, et al (2015) A new perspective on lysogeny:

671 prophages as active regulatory switches of bacteria. *Nat Rev Microbiol*
 672 13:641–650. <https://doi.org/10.1038/nrmicro3527>
 673 Fernandes RM, de Lencastre H, Archer LJ (1989) Specialized transduction in *Bacillus*
 674 subtilis by the phages IG1, IG3, and IG4. *Arch Virol* 105:137–40
 675 Field CB, Behrenfeld MJ, Randerson JT, Falkowski P (1998) Primary production of the
 676 biosphere: Integrating terrestrial and oceanic components. *Science* (80-)
 677 281:237–240. <https://doi.org/10.1126/science.281.5374.237>
 678 Findlay SEG, Parr TB (2017) Chapter 24 - Dissolved Organic Matter. In: Lamberti GA,
 679 Hauer FRBT-M in SE (Third E (eds). Academic Press, pp 21–36
 680 Frias-Lopez J, Shi Y, Tyson GW, et al (2008) Microbial community gene expression in
 681 ocean surface waters. *Proc Natl Acad Sci* 105:3805–3810.
 682 <https://doi.org/10.1073/pnas.0708897105>
 683 Fuhrman J, Suttle C (1993) Viruses in Marine Planktonic Systems. *Oceanography*
 684 6:51–63. <https://doi.org/10.5670/oceanog.1993.14>
 685 Fuhrman JA (1999) Marine viruses and their biogeochemical and ecological effects.
 686 *Nature* 399:541–548. <https://doi.org/10.1038/21119>
 687 Gonzalez JM, Suttle CA (1993) Grazing by marine nanoflagellates on viruses and
 688 virus-sized particles: ingestion and digestion. *Mar Ecol Prog Ser* 94:1–10.

689 <https://doi.org/10.3354/meps094001>

690 Gottesman ME, Yarmolinsky MB (1968) Integration-negative mutants of bacteriophage

691 lambda. *J Mol Biol* 31:487–505. [https://doi.org/10.1016/0022-2836\(68\)90423-3](https://doi.org/10.1016/0022-2836(68)90423-3)

692 Guidi L, Chaffron S, Bittner L, et al (2016) Plankton networks driving carbon export in

693 the oligotrophic ocean. *Nature* 532:465–470. <https://doi.org/10.1038/nature16942>

694 Guttman B, Raya R, Kutter E (2004) *Basic Phage Biology*

695 Hall AR, Scanlan PD, Buckling A (2011a) Bacteria-phage coevolution and the

696 emergence of generalist pathogens. *Am Nat* 177:44–53.

697 <https://doi.org/10.1086/657441>

698 Hall AR, Scanlan PD, Morgan AD, Buckling A (2011b) Host–parasite coevolutionary

699 arms races give way to fluctuating selection. *Ecol Lett* 14:635–642.

700 <https://doi.org/10.1111/j.1461-0248.2011.01624.x>

701 Hendrix RW, Lawrence JG, Hatfull GF, Casjens S (2000) The origins and ongoing

702 evolution of viruses. *Trends Microbiol* 8:504–508.

703 [https://doi.org/https://doi.org/10.1016/S0966-842X\(00\)01863-1](https://doi.org/https://doi.org/10.1016/S0966-842X(00)01863-1)

704 Hobbs Z, Abedon ST (2016) Diversity of phage infection types and associated

705 terminology: the problem with ‘Lytic or lysogenic.’ *FEMS Microbiol Lett* 363:.

706 <https://doi.org/10.1093/femsle/fnw047>

707 Howard-Varona C, Hargreaves KR, Abedon ST, Sullivan MB (2017a) Lysogeny in
 708 nature: Mechanisms, impact and ecology of temperate phages. *ISME J*
 709 11:1511–1520. <https://doi.org/10.1038/ismej.2017.16>
 710 Howard-Varona C, Lindback MM, Bastien GE, et al (2020) Phage-specific metabolic
 711 reprogramming of virocells. *ISME J* 14:881–895.
 712 <https://doi.org/10.1038/s41396-019-0580-z>
 713 Howard-Varona C, Roux S, Dore H, et al (2017b) Regulation of infection efficiency in
 714 a globally abundant marine Bacterioidetes virus. *ISME J* 11:284–295.
 715 <https://doi.org/10.1038/ismej.2016.81>
 716 Hurwitz BL, Brum JR, Sullivan MB (2015) Depth-stratified functional and taxonomic
 717 niche specialization in the “core” and “flexible” Pacific Ocean Virome. *ISME J*
 718 9:472–484. <https://doi.org/10.1038/ismej.2014.143>
 719 Hurwitz BL, U’Ren JM (2016) Viral metabolic reprogramming in marine ecosystems.
 720 *Curr Opin Microbiol* 31:161–168. <https://doi.org/10.1016/j.mib.2016.04.002>
 721 Hutchinson GE (1961) The Paradox of the Plankton. *Am Nat* 95:137–145
 722 Ignacio-Espinoza JC, Ahlgren NA, Fuhrman JA (2020) Long-term stability and Red
 723 Queen-like strain dynamics in marine viruses. *Nat Microbiol* 5:265–271.
 724 <https://doi.org/10.1038/s41564-019-0628-x>

725 Jover LF, Effler TC, Buchan A, et al (2014) The elemental composition of virus
 726 particles: Implications for marine biogeochemical cycles. *Nat Rev Microbiol*
 727 12:519–528. <https://doi.org/10.1038/nrmicro3289>
 728 Kelly L, Ding H, Huang KH, et al (2013) Genetic diversity in cultured and wild marine
 729 cyanomyoviruses reveals phosphorus stress as a strong selective agent. *ISME J*
 730 7:1827–1841. <https://doi.org/10.1038/ismej.2013.58>
 731 Kettler GC, Martiny AC, Huang K, et al (2007) Patterns and implications of gene gain
 732 and loss in the evolution of *Prochlorococcus*. *PLoS Genet* 3:2515–2528.
 733 <https://doi.org/10.1371/journal.pgen.0030231>
 734 Kieft K, Zhou Z, Anderson RE, et al (2020) Ecology of inorganic sulfur auxiliary
 735 metabolism in widespread bacteriophages. *bioRxiv* 2020.08.24.253096.
 736 <https://doi.org/10.1101/2020.08.24.253096>
 737 Kimura S, Sako Y, Yoshida T (2013) Rapid *Microcystis* cyanophage gene
 738 diversification revealed by long and short-term genetic analyses of the tail sheath
 739 gene in a natural pond. *Appl Environ Microbiol* 79:2789–2795.
 740 <https://doi.org/10.1128/AEM.03751-12>
 741 Kimura S, Uehara M, Morimoto D, et al (2018) Incomplete selective sweeps of
 742 *Microcystis* population detected by the leader-end CRISPR fragment analysis in a

743 natural pond. *Front Microbiol* 9:article 425.
 744 <https://doi.org/10.3389/fmicb.2018.00425>

745 Kimura S, Yoshida T, Hosoda N, et al (2012) Diurnal infection patterns and impact of
 746 *Microcystis cyanophages* in a Japanese pond. *Appl Environ Microbiol*
 747 78:5805–5811. <https://doi.org/10.1128/AEM.00571-12>

748 Knowles B, Bailey B, Boling L, et al (2017) Variability and host density independence
 749 in inductions-based estimates of environmental lysogeny. *Nat Microbiol* 2:17064.
 750 <https://doi.org/10.1038/nmicrobiol.2017.64>

751 Knowles B, Silveira CB, Bailey BA, et al (2016) Lytic to temperate switching of viral
 752 communities. *Nature* 531:466–470. <https://doi.org/10.1038/nature17193>

753 Koskella B, Brockhurst MA (2014) Bacteria-phage coevolution as a driver of ecological
 754 and evolutionary processes in microbial communities. *FEMS Microbiol Rev*
 755 38:916–931. <https://doi.org/10.1111/1574-6976.12072>

756 Ku C, Ku C, Sheyn U, et al (2020) A single-cell view on alga-virus interactions reveals
 757 sequential transcriptional programs and infection states. *Sci Adv* 6:.
 758 <https://doi.org/10.1126/sciadv.aba4137>

759 Kuno S, Sako Y, Yoshida T (2014) Diversification of CRISPR within coexisting
 760 genotypes in a natural population of the bloom-forming cyanobacterium

761 *Microcystis aeruginosa*. Microbiology 160:903–916.

762 <https://doi.org/10.1099/mic.0.073494-0>

763 Kuno S, Yoshida T, Kaneko T, Sako Y (2012) Intricate interactions between the

764 bloom-forming cyanobacterium *Microcystis aeruginosa* and foreign genetic

765 elements, revealed by diversified clustered regularly interspaced short palindromic

766 repeat (CRISPR) signatures. Appl Environ Microbiol 78:5353–5360.

767 <https://doi.org/10.1128/AEM.00626-12>

768 Kutter E, Bryan D, Ray G, et al (2018) From host to phage metabolism: Hot tales of

769 phage T4’s takeover of *E. coli*. Viruses 10:. <https://doi.org/10.3390/v10070387>

770 Laber CP, Hunter JE, Carvalho F, et al (2018) Coccolithovirus facilitation of carbon

771 export in the North Atlantic. Nat Microbiol 3:537–547.

772 <https://doi.org/10.1038/s41564-018-0128-4>

773 Lawrence J, Töpper J, Petelenz-Kurdziel E, et al (2018) Viruses on the menu: The

774 appendicularian *Oikopleura dioica* efficiently removes viruses from seawater.

775 Limnol Oceanogr 63:S244–S253. <https://doi.org/10.1002/lno.10734>

776 Leenheer JA, Croué J-P (2003) Peer Reviewed: Characterizing Aquatic Dissolved

777 Organic Matter. Environ Sci Technol 37:18A-26A.

778 <https://doi.org/10.1021/es032333c>

779 Lenski RE, Levin BR (1985) Constraints on the coevolution of bacteria and virulent
 780 phage: A model, some experiments, and predictions for natural communities. *Am*
 781 *Nat* 125:585–602. <https://doi.org/10.1086/284364>
 782 Lin X, Ding H, Zeng Q (2016) Transcriptomic response during phage infection of a
 783 marine cyanobacterium under phosphorus-limited conditions. *Environ Microbiol*
 784 18:450–460. <https://doi.org/10.1111/1462-2920.13104>
 785 Lindell D, Jaffe JD, Coleman ML, et al (2007) Genome-wide expression dynamics of a
 786 marine virus and host reveal features of co-evolution. *Nature* 449:83–86.
 787 <https://doi.org/10.1038/nature06130>
 788 Lindell D, Jaffe JD, Johnson ZI, et al (2005) Photosynthesis genes in marine viruses
 789 yield proteins during host infection. *Nature* 438:86–89.
 790 <https://doi.org/10.1038/nature04111>
 791 Lindell D, Sullivan MB, Johnson ZI, et al (2004) Transfer of photosynthesis genes to
 792 and from *Prochlorococcus* viruses. *Proc Natl Acad Sci U S A* 101:11013–11018.
 793 <https://doi.org/10.1073/pnas.0401526101>
 794 Ma X, Coleman ML, Waldbauer JR (2018) Distinct molecular signatures in dissolved
 795 organic matter produced by viral lysis of marine cyanobacteria. *Environ Microbiol*
 796 20:3001–3011. <https://doi.org/10.1111/1462-2920.14338>

797 Maharjan R, Seeto S, Notley-McRobb L, Ferenci T (2006) Clonal adaptive radiation in
 798 a constant environment. *Science* (80-) 313:514–517.
 799 <https://doi.org/10.1126/science.1129865>
 800 Makarova KS, Wolf YI, Snir S, Koonin E V. (2011) Defense islands in bacterial and
 801 archaeal genomes and prediction of novel defense systems. *J Bacteriol*
 802 193:6039–6056. <https://doi.org/10.1128/JB.05535-11>
 803 Marston MF, Pierciey FJ, Shepard A, et al (2012) Rapid diversification of coevolving
 804 marine *Synechococcus* and a virus. *Proc Natl Acad Sci U S A* 109:4544–4549.
 805 <https://doi.org/10.1073/pnas.1120310109>
 806 Middelboe M, Hagström A, Blackburn N, et al (2001) Effects of bacteriophages on the
 807 population dynamics of four strains of pelagic marine bacteria. *Microb Ecol*
 808 42:395–406. <https://doi.org/10.1007/s00248-001-0012-1>
 809 Middelboe M, Holmfeldt K, Riemann L, et al (2009) Bacteriophages drive strain
 810 diversification in a marine *Flavobacterium*: implications for phage resistance and
 811 physiological properties. *Environ Microbiol* 11:1971–1982.
 812 <https://doi.org/10.1111/j.1462-2920.2009.01920.x>
 813 Mizoguchi K, Morita M, Fischer CR, et al (2003) Coevolution of bacteriophage PP01
 814 and *Escherichia coli* O157:H7 in continuous culture. *Appl Environ Microbiol*

815 69:170 LP – 176. <https://doi.org/10.1128/AEM.69.1.170-176.2003>

816 Mizuno CM, Ghai R, Rodriguez-Valera F (2014) Evidence for metaviromic islands in
817 marine phages. *Front Microbiol* 5:1–10. <https://doi.org/10.3389/fmicb.2014.00027>

818 Monier A, Chambouvet A, Milner DS, et al (2017) Host-derived viral transporter
819 protein for nitrogen uptake in infected marine phytoplankton. *Proc Natl Acad Sci*
820 114:E7489 LP-E7498. <https://doi.org/10.1073/pnas.1708097114>

821 Moniruzzaman M, Gann ER, Wilhelm SW (2018) Infection by a Giant Virus (AaV)
822 Induces Widespread Physiological Reprogramming in *Aureococcus*
823 *anophagefferens* CCMP1984 – A Harmful Bloom Algae . *Front. Microbiol.*
824 9:752

825 Moniruzzaman M, Martinez-Gutierrez CA, Weinheimer AR, Aylward FO (2020)
826 Dynamic genome evolution and complex virocell metabolism of
827 globally-distributed giant viruses. *Nat Commun* 11:1710.
828 <https://doi.org/10.1038/s41467-020-15507-2>

829 Morimoto D, Kimura S, Sako Y, Yoshida T (2018) Transcriptome analysis of a
830 bloom-forming cyanobacterium *Microcystis aeruginosa* during Ma-LMM01 phage
831 infection. *Front Microbiol* 9:article 2. <https://doi.org/10.3389/fmicb.2018.00002>

832 Morimoto D, Šulčius S, Tominaga K, Yoshida T (2020) Predetermined clockwork

833 microbial worlds: Current understanding of aquatic microbial diel response from
 834 model systems to complex environments. *Adv Appl Microbiol* 1–29.
 835 <https://doi.org/10.1016/bs.aambs.2020.06.001>

836 Morimoto D, Tominaga K, Nishimura Y, et al (2019) Cooccurrence of broad- and
 837 narrow-host-range viruses infecting the bloom-forming toxic cyanobacterium
 838 *Microcystis aeruginosa*. *Appl Environ Microbiol* 85:e01170-19

839 Morris RM, Cain KR, Hvorecny KL, Kollman JM (2020) Lysogenic host–virus
 840 interactions in SAR11 marine bacteria. *Nat Microbiol* 5:1011–1015.
 841 <https://doi.org/10.1038/s41564-020-0725-x>

842 Needham DM, Chow CET, Cram JA, et al (2013) Short-term observations of marine
 843 bacterial and viral communities: Patterns, connections and resilience. *ISME J*
 844 7:1274–1285. <https://doi.org/10.1038/ismej.2013.19>

845 Needham DM, Fichot EB, Wang E, et al (2018) Dynamics and interactions of highly
 846 resolved marine plankton via automated high-frequency sampling. *ISME J*
 847 12:2417–2432. <https://doi.org/10.1038/s41396-018-0169-y>

848 Needham DM, Sachdeva R, Fuhrman JA (2017) Ecological dynamics and
 849 co-occurrence among marine phytoplankton, bacteria and myoviruses shows
 850 microdiversity matters. *ISME J* 11:1614–1629.

851 <https://doi.org/10.1038/ismej.2017.29>

852 Needham DM, Yoshizawa S, Hosaka T, et al (2019) A distinct lineage of giant viruses
853 brings a rhodopsin photosystem to unicellular marine predators. *Proc Natl Acad*
854 *Sci* 116:20574 LP – 20583. <https://doi.org/10.1073/pnas.1907517116>

855 Nissimov JI, Vandzura R, Johns CT, et al (2018) Dynamics of transparent exopolymer
856 particle production and aggregation during viral infection of the coccolithophore,
857 *Emiliania huxleyi*. *Environ Microbiol* 20:2880–2897.
858 <https://doi.org/10.1111/1462-2920.14261>

859 Parada AE, Fuhrman JA (2017) Marine archaeal dynamics and interactions with the
860 microbial community over 5 years from surface to seafloor. *ISME J* 11:2510–2525.
861 <https://doi.org/10.1038/ismej.2017.104>

862 Parsons RJ, Breitbart M, Lomas MW, Carlson CA (2012) Ocean time-series reveals
863 recurring seasonal patterns of virioplankton dynamics in the northwestern Sargasso
864 Sea. *ISME J* 6:273–284. <https://doi.org/10.1038/ismej.2011.101>

865 Paul JH (2008) Prophages in marine bacteria: dangerous molecular time bombs or the
866 key to survival in the seas? *ISME J* 2:579–589.
867 <https://doi.org/10.1038/ismej.2008.35>

868 Pourtois J, Tarnita CE, Bonachela JA (2020) Impact of Lytic Phages on Phosphorus- vs.

869 Nitrogen-Limited Marine Microbes. *Front Microbiol* 11:.
870 <https://doi.org/10.3389/fmicb.2020.00221>

871 Reva O, Tümmler B (2008) Think big - Giant genes in bacteria. *Environ Microbiol*
872 10:768–777. <https://doi.org/10.1111/j.1462-2920.2007.01500.x>

873 Rodriguez-Brito B, Li L, Wegley L, et al (2010) Viral and microbial community
874 dynamics in four aquatic environments. *ISME J* 4:739–751.
875 <https://doi.org/10.1038/ismej.2010.1>

876 Rodriguez-Valera F, Martin-Cuadrado AB, Rodriguez-Brito B, et al (2009) Explaining
877 microbial population genomics through phage predation. *Nat Rev Microbiol*
878 7:828–836. <https://doi.org/10.1038/nrmicro2235>

879 Rodriguez-Valera F, Ussery DW (2012) Is the pan-genome also a pan-selectome?
880 *F1000Research* 1:1–7. <https://doi.org/10.3410/f1000research.1-16.v1>

881 Rosenwasser S, Mausz MA, Schatz D, et al (2014) Rewiring host lipid metabolism by
882 large viruses determines the fate of *Emiliana huxleyi*, a bloom-forming alga in the
883 ocean. *Plant Cell* 26:2689–2707. <https://doi.org/10.1105/tpc.114.125641>

884 Roucourt B, Lavigne R (2009) The role of interactions between phage and bacterial
885 proteins within the infected cell: a diverse and puzzling interactome. *Environ*
886 *Microbiol* 11:2789–2805. <https://doi.org/10.1111/j.1462-2920.2009.02029.x>

887 Schulz F, Roux S, Paez-Espino D, et al (2020) Giant virus diversity and host
 888 interactions through global metagenomics. *Nature* 578:432–436.
 889 <https://doi.org/10.1038/s41586-020-1957-x>

890 Schvarcz CR, Steward GF (2018) A giant virus infecting green algae encodes key
 891 fermentation genes. *Virology* 518:423–433.
 892 <https://doi.org/https://doi.org/10.1016/j.virol.2018.03.010>

893 Schwalbach MS, Hewson I, Fuhrman JA (2004) Viral effects on bacterial community
 894 composition in marine plankton microcosms. *Aquat Microb Ecol* 34:117–127.
 895 <https://doi.org/10.3354/ame034117>

896 Sharma RS, Mishra V, Mohmmmed A, Babu CR (2008) Phage specificity and
 897 lipopolysaccharides of stem- and root-nodulating bacteria (*Azorhizobium*
 898 *caulinodans*, *Sinorhizobium* spp., and *Rhizobium* spp.) of *Sesbania* spp. *Arch*
 899 *Microbiol* 189:411–418. <https://doi.org/10.1007/s00203-007-0322-x>

900 Sheyn U, Rosenwasser S, Lehahn Y, et al (2018) Expression profiling of host and virus
 901 during a coccolithophore bloom provides insights into the role of viral infection in
 902 promoting carbon export. *ISME J* 12:704–713.
 903 <https://doi.org/10.1038/s41396-017-0004-x>

904 Sieradzki ET, Ignacio-Espinoza JC, Needham DM, et al (2019) Dynamic marine viral

905 infections and major contribution to photosynthetic processes shown by
 906 spatiotemporal picoplankton metatranscriptomes. Nat Commun 10:.
 907 <https://doi.org/10.1038/s41467-019-09106-z>
 908 Silkin V, Pautova L, Kravchishina M, et al (2020) Dataset of the *Emiliania huxleyi*
 909 abundance and phytoplankton composition in the Barents Sea in summer
 910 2014–2018. Data Br 32:106251.
 911 <https://doi.org/https://doi.org/10.1016/j.dib.2020.106251>
 912 Sime-Ngando T (2014) Environmental bacteriophages: Viruses of microbes in aquatic
 913 ecosystems. Front Microbiol 5:1–14. <https://doi.org/10.3389/fmicb.2014.00355>
 914 Sorek R, Kunin V, Hugenholtz P (2008) CRISPR - A widespread system that provides
 915 acquired resistance against phages in bacteria and archaea. Nat Rev Microbiol
 916 6:181–186. <https://doi.org/10.1038/nrmicro1793>
 917 Stewart FM, Levin BR (1984) The population biology of bacterial viruses: Why be
 918 temperate. Theor Popul Biol 26:93–117.
 919 [https://doi.org/https://doi.org/10.1016/0040-5809\(84\)90026-1](https://doi.org/https://doi.org/10.1016/0040-5809(84)90026-1)
 920 Sullivan MB, Weitz JS, Wilhelm S (2017) Viral ecology comes of age. Environ
 921 Microbiol Rep 9:33–35. <https://doi.org/10.1111/1758-2229.12504>
 922 Suttle CA (2005) Viruses in the sea. Nature 437:356–361.

923 <https://doi.org/10.1038/nature04160>

924 Suttle CA (2007) Marine viruses - major players in the global ecosystem. *Nat Rev*

925 *Microbiol* 5:801–812. <https://doi.org/10.1038/nrmicro1750>

926 Suttle CA, Chen F (1992) Mechanisms and rates of decay of marine viruses in seawater.

927 *Appl Environ Microbiol* 58:3721–3729.

928 <https://doi.org/10.1128/aem.58.11.3721-3729.1992>

929 Tarutani K, Nagasaki K, Yamaguchi M (2000) Viral Impacts on Total Abundance and

930 Clonal Composition of the Harmful Bloom-Forming

931 PhytoplanktonHeterosigma akashiwo *Appl Environ*

932 *Microbiol* 66:4916 LP – 4920.

933 <https://doi.org/10.1128/AEM.66.11.4916-4920.2000>

934 Thierauf A, Perez G, Maloy S (2009) Generalized transduction. *Methods Mol Biol*

935 501:267–286. <https://doi.org/10.1007/978-1-60327-565-1>

936 Thingstad TF (2000) Elements of a theory for the mechanisms controlling abundance,

937 diversity, and biogeochemical role of lytic bacterial viruses in aquatic systems.

938 *Limnol Oceanogr* 45:1320–1328. <https://doi.org/10.4319/lo.2000.45.6.1320>

939 Thingstad TF, Vage S, Storesund JE, et al (2014) A theoretical analysis of how

940 strain-specific viruses can control microbial species diversity. *Proc Natl Acad Sci*

941 U S A 111:7813–7818. <https://doi.org/10.1073/pnas.1400909111>
 942 Thompson LR, Zeng Q, Kelly L, et al (2011) Phage auxiliary metabolic genes and the
 943 redirection of cyanobacterial host carbon metabolism. *Proc Natl Acad Sci*
 944 108:E757–E764. <https://doi.org/10.1073/pnas.1102164108>
 945 Touchon M, Bernheim A, Rocha EPC (2016) Genetic and life-history traits associated
 946 with the distribution of prophages in bacteria. *ISME J* 10:2744–2754.
 947 <https://doi.org/10.1038/ismej.2016.47>
 948 Touchon M, Moura de Sousa JA, Rocha EP (2017) Embracing the enemy: The
 949 diversification of microbial gene repertoires by phage-mediated horizontal gene
 950 transfer. *Curr Opin Microbiol* 38:66–73. <https://doi.org/10.1016/j.mib.2017.04.010>
 951 Valen VL (1973) A new evolutionary law. *Evol Theory* 1:1–30
 952 Warwick-Dugdale J, Buchholz HH, Allen MJ, Temperton B (2019) Host-hijacking and
 953 planktonic piracy: how phages command the microbial high seas. *Virol J* 16:15.
 954 <https://doi.org/10.1186/s12985-019-1120-1>
 955 Waterbury JB, Valois FW (1993) Resistance to co-occurring phages enables marine
 956 *Synechococcus* communities to coexist with cyanophages abundant in seawater.
 957 *Appl Environ Microbiol* 59:3393–3399.
 958 <https://doi.org/10.1128/aem.59.10.3393-3399.1993>

959 Wei Y, Kirby A, Levin BR, et al (2011) The population and evolutionary dynamics of
 960 *Vibrio cholerae* and its bacteriophage: Conditions for maintaining phage-limited
 961 communities. *Am Nat* 178:715–725. <https://doi.org/10.1086/662677>
 962 Wei Y, Ocampo P, Levin BR (2010) An experimental study of the population and
 963 evolutionary dynamics of *Vibrio cholerae* O1 and the bacteriophage JSF4. *Proc R*
 964 *Soc B Biol Sci* 277:3247–3254. <https://doi.org/10.1098/rspb.2010.0651>
 965 Weinbauer MG (2004) Ecology of prokaryotic viruses. *FEMS Microbiol Rev*
 966 28:127–181. <https://doi.org/10.1016/j.femsre.2003.08.001>
 967 Weitz JS, Stock CA, Wilhelm SW, et al (2015) A multitrophic model to quantify the
 968 effects of marine viruses on microbial food webs and ecosystem processes. *ISME J*
 969 9:1352–1364. <https://doi.org/10.1038/ismej.2014.220>
 970 Welsh JE, Steenhuis P, de Moraes KR, et al (2020) Marine virus predation by non-host
 971 organisms. *Sci Rep* 10:1–9. <https://doi.org/10.1038/s41598-020-61691-y>
 972 Whitman WB, Coleman DC, Wiebe WJ (1998) Prokaryotes: The unseen majority. *Proc*
 973 *Natl Acad Sci U S A* 95:6578–6583. <https://doi.org/10.1073/pnas.95.12.6578>
 974 Wigington CH, Sonderegger DL, Brussaard CPD, et al (2016) Re-examining of the
 975 relationship between marine virus and microbial cell abundances. *Nat Microbiol*
 976 1:10.1038/nmicrobiol.2015.24. <https://doi.org/10.1101/025544>

977 Wikner J, Vallino JJ, Steward GF, et al (1993) Nucleic acids from the host bacterium as
 978 a major source of nucleotides for three marine bacteriophages. *FEMS Microbiol*
 979 *Ecol* 12:237–248. <https://doi.org/10.1111/j.1574-6941.1993.tb00036.x>
 980 Wilhelm LJ, Tripp HJ, Givan SA, et al (2007) Natural variation in SAR11 marine
 981 bacterioplankton genomes inferred from metagenomic data. *Biol Direct* 2:1–19.
 982 <https://doi.org/10.1186/1745-6150-2-27>
 983 Wilhelm SW, Suttle CA (1999) Viruses and Nutrient Cycles in the Sea: Viruses play
 984 critical roles in the structure and function of aquatic food webs. *Bioscience*
 985 49:781–788. <https://doi.org/10.2307/1313569>
 986 Williamson SJ, Rusch DB, Yooseph S, et al (2008) The sorcerer II global ocean
 987 sampling expedition: Metagenomic characterization of viruses within aquatic
 988 microbial samples. *PLoS One* 3:. <https://doi.org/10.1371/journal.pone.0001456>
 989 Wilmes P, Simmons SL, Denef VJ, Banfield JF (2009) The dynamic genetic repertoire
 990 of microbial communities. *FEMS Microbiol Rev* 33:109–132.
 991 <https://doi.org/10.1111/j.1574-6976.2008.00144.x>
 992 Winter C, Bouvier T, Weinbauer MG, Thingstad TF (2010) Trade-Offs between
 993 Competition and Defense Specialists among Unicellular Planktonic Organisms: the
 994 “Killing the Winner” Hypothesis Revisited. *Microbiol Mol Biol Rev* 74:42–57.

995 <https://doi.org/10.1128/mmbr.00034-09>

996 Wommack KE, Colwell RR (2000) Virioplankton: Viruses in Aquatic Ecosystems.

997 Microbiol Mol Biol Rev 64:69–114.

998 <https://doi.org/10.1128/mmbr.64.1.69-114.2000>

999 Yamada Y, Guillemette R, Baudoux AC, et al (2020) Viral attachment to biotic and

1000 abiotic surfaces in seawater. Appl Environ Microbiol 86:1–18.

1001 <https://doi.org/10.1128/AEM.01687-19>

1002 Yamada Y, Tomaru Y, Fukuda H, Nagata T (2018) Aggregate formation during the

1003 viral lysis of a marine diatom. Front Mar Sci 5:1–7.

1004 <https://doi.org/10.3389/fmars.2018.00167>

1005 Yoshida-Takashima Y, Yoshida M, Ogata H, et al (2012) Cyanophage infection in the

1006 bloom-forming cyanobacteria *Microcystis aeruginosa* in surface freshwater.

1007 Microbes Environ 27:350–355. <https://doi.org/10.1264/jsme2.ME12037>

1008 Yoshida M, Yoshida T, Kashima A, et al (2008a) Ecological dynamics of the toxic

1009 bloom-forming cyanobacterium *Microcystis aeruginosa* and its cyanophages in

1010 freshwater. Appl Environ Microbiol 74:3269–3273.

1011 <https://doi.org/10.1128/AEM.02240-07>

1012 Yoshida M, Yoshida T, Satomi M, et al (2008b) Intra-specific phenotypic and genotypic

1013 variation in toxic cyanobacterial *Microcystis* strains. *J Appl Microbiol*
 1014 105:407–415. <https://doi.org/10.1111/j.1365-2672.2008.03754.x>
 1015 Yoshida M, Yoshida T, Takashima Y, et al (2005) Genetic diversity of the toxic
 1016 cyanobacterium *Microcystis* in Lake Mikata. *Environ Toxicol* 20:229–234.
 1017 <https://doi.org/10.1002/tox.20102>
 1018 Yoshida T, Morimoto D, Kimura S (2019) Bacteria-virus interactions. In: Nishida H,
 1019 Oshima T (eds) *DNA Traffic in the Environment*. Springer Singapore, Singapore,
 1020 pp 95–108
 1021 Zborowsky S, Lindell D (2019) Resistance in marine cyanobacteria differs against
 1022 specialist and generalist cyanophages. *Proc Natl Acad Sci* 116:16899–16908.
 1023 <https://doi.org/10.1073/pnas.1906897116>
 1024 Zeng Q, Chisholm SW (2012) Marine viruses exploit their host’s two-component
 1025 regulatory system in response to resource limitation. *Curr Biol* 22:124–128.
 1026 <https://doi.org/10.1016/j.cub.2011.11.055>
 1027 Zhao Y, Temperton B, Thrash JC, et al (2013) Abundant SAR11 viruses in the ocean.
 1028 *Nature* 494:357–360. <https://doi.org/10.1038/nature11921>
 1029 Zsolnay Á (2003) Dissolved organic matter: artefacts, definitions, and functions.
 1030 *Geoderma* 113:187–209.

1031 [https://doi.org/https://doi.org/10.1016/S0016-7061\(02\)00361-0](https://doi.org/https://doi.org/10.1016/S0016-7061(02)00361-0)

1032