

NON-REGULAR SEMIGROUPS WHICH ARE AMALGAMATION BASES *

KUNITAKA SHOJI DEPARTMENT OF MATHEMATICS
SHIMANE UNIVERSITY

In this paper, we study non-regular semigroups which are amalgamation basess for finite semigroups or for all semigroups.

1 Semigroup amalgamation bases

Definition. Let $\mathcal{A}$ be the class of finite semigroups or the class of all semigroups. Let S, T, U be semigroups in $\mathcal{A}$ such that U is a subsemigroup of S and T in common. Then a triple $[S, T; U]$ is called an amalgam of semigroups S, T with U as a core in $\mathcal{A}$. An amalgam $[S, T; U]$ of $\mathcal{A}$ is called to be *weakly embeddable* in $\mathcal{A}$ if there exist a semigroup K belonging to $\mathcal{A}$ and monomorphisms $\xi_1 : S \rightarrow K$, $\xi_2 : T \rightarrow K$ such that the restrictions to U of ξ_1 and ξ_2 are equal to each other (that is, $\xi_1(S) \cap \xi_2(T) \supseteq \xi_1(U)$). An amalgam $[S, T; U]$ of $\mathcal{A}$ is called to be *strongly embeddable* in $\mathcal{A}$ if $\xi_1(S) \cap \xi_2(T) = \xi_1(U)$. A semigroup U in $\mathcal{A}$ is *amalgamation base* [resp. *weak amalgamation base*] if any amalgam with a core U in $\mathcal{A}$ is strongly embeddable [resp. weakly embeddable] in $\mathcal{A}$.

We have the following results which will be used later.

Result 1[[4], Theorem 12]. *Any finite semigroup U is an amalgamation base for finite semigroups if and only if U is a weak amalgamation base for finite semigroups [for all semigroup] .*

Result 2[[6], Theorem 1]. *If a finite semigroup U is an amalgamation base for finite semigroups, then all $\mathcal{J}$ -classes of U form a chain.*

*This is an absrtact and the paper will appear elsewhere.

2 Non-regular amalgamation bases for all semigroups

In the several papers [1], [4], [12] and etc., the study of regular amalgamation bases for all semigroups has been made.

From now on we discuss non-regular amalgamation bases for all semigroups.

The Kimura's counter-example was stated in the second volume of the book [2].

Example. Let $U = \{u, v, w, 0\}$ be a null semigroup in which all products are equal to 0. Let $S_1 = U \cup \{a\}$ where $a \notin U$, $au = ua = v$, and all the other products in S_1 are set equal to z . Let $S_2 = U \cup \{b\}$ where $b \notin U$, $bv = vb = w$, and all other products in S_2 are set equal to 0. Then the amalgam $[S_1, S_2; U]$ is not embeddable in the class of all semigroups.

Actually, $w = bv = b(ua) = (bu)a = 0a = 0$ in any oversemigroup, a contradiction.

Thus we have

Theorem 1. *Any null semigroups with at least 3 elements are not amalgamation bases for all semigroups*

On the other hand, it was known that

Result 3[4, Corollary 26]. *2-element semigroups are amalgamation bases for all semigroups.*

Result 4[Theorem 4.1, [4]]. *Let S be a semigroup with 0 consisting of a group G of units and a nilpotent ideal N . Then S is an amalgamation base for all semigroups if and only if there exists $a \in N$ such that $N = Ga \cup Ga^2 \cup \dots \cup Ga^n \cup \{0\}$ and $Ga = aG$*

Definition. Let S be a commutative semigroup with only finitely many $\mathcal{J}$ -classes, where $\mathcal{J}$ denotes the Green's J -relation on S .

Define a quasi-order $\geq_{\mathcal{J}}$ on S by $s \geq_{\mathcal{J}} t$ if and only if $Ss \supseteq St$.

$s >_{\mathcal{J}} t$ if $S^1s \supset S^1t$.

Let a, b be elements of S . Then we say that the pair a and b are $\mathcal{J}$ -comparable if $a \geq_{\mathcal{J}} b$ or $b \geq_{\mathcal{J}} a$. Otherwise, a, b are $\mathcal{J}$ -incomparable. Also, $\mathcal{J}$ -incomparable elements a, b of S are called E -distinct if there exists an idempotent $e \in S$ such that (i) either $a\mathcal{J}ea, b >_{\mathcal{J}} eb$ or (ii) $a >_{\mathcal{J}} ea, b\mathcal{J}eb$.

A subset A of S is called E -distinct if any pair of $\mathcal{J}$ -incomparable elements of A are E -distinct.

Result 5[The main theorem, [10]]. *Let S be a commutative semigroup with only finitely many $\mathcal{J}$ -classes. Then S is E -distinct if and only if S is an amalgamation base for all semigroups.*

Result 6[Corollary, [10]]. *Let S be a commutative semigroup with only finitely many $\mathcal{J}$ -classes. Then S is an amalgamation base for all semigroups if and only if all factor semigroups of S are an amalgamation base for all semigroups.*

3 Non-regular amalgamation bases for finite semigroups

Result 7[Theorem 20, [4]]. *Any finite cyclic semigroup is an amalgamation base for semigroups and for finite semigroups*

Theorem 2 *Let S be a semigroup with 0 consisting of a group G of units and a nilpotent ideal N . Then S is an amalgamation base for finite semigroups if and only if there exists $a \in N$ such that $N = Ga \cup Ga^2 \cup \dots \cup Ga^n \cup \{0\}$ and $Ga = aG$.*

References

- [1] S. Bulman-Fleming and K. McDowell. *Absolutely flat semigroups*. Pacific J. Math. **107**(1983), 319-333.
- [2] A. H. Clifford and G. B. Preston, *Algebraic theory of semigroups*, Amer. Math. Soc., Math. Survey, No.7, Providence, R.I., Vol.I(1961); Vol.II(1967).
- [3] L. M. Gluskin, *Elementary Generalized Groups* Mat. Sb. **41**(83)(1957), 23-36.
- [4] T. E. Hall, *Representation extension and amalgamation for semigroups*. Quart. J. Math. Oxford (2) **29**(1978), 309-334.
- [5] T. E. Hall. *Finite inverse semigroups and amalgamations*. Semigroups and their applications, Reidel, 1987, pp. 51-56.
- [6] T. E. Hall and M.S. Putcha, *The potential $\mathcal{J}$ -relation and amalgamation bases for finite semigroups*, Proc. Amer. Math. Soc. **95**(1985), 309-334.
- [7] T. E. Hall and K. Shoji, *Finite bands and amalgamation bases for finite semigroups*, Communications in Algebra **30**(2) (2002), 911-933.

- [8] J. Okniński and M.S. Putcha, *Embedding finite semigroup amalgams*, J. Austral. Math. Soc. (Series A) **51**(1991), 489-496.
- [9] *Amalgamation bases for semigroups*, Math. Japonicae **35**(1990), 473-483.
- [10] *Commutative semigroups which are semigroup amalgamation bases*, J. Algebra, **238**(2001), 1-50.
- [11] K. Shoji, *A proof of Okniński and Putcha's theorem*, Proc. of the Third Inter. Colloq. on Words, Languages and Combinatorics. Ed. by M. Ito & T. Imaoka World Scientific, 2003, 420-427.
- [12] K. Shoji, *Regular semigroups which are amalgamation bases for finite semigroups*, Algebra Colloquium **14**(2007) 245-254.
- [13] B. M. Schein, *A symmetric semigroup of transformations is covered by its inverse subsemigroups*, Acta Math. Acad. Sc. Hungar. **22**(1971), 163-171.