

On Behaviors of Hypergeometric Series
of One, Two, Three and n Variables
Near Boundaries of Their Convergence Regions
of Logarithmic Case

Megumi Saigo

Department of Applied Mathematics, Fukuoka University

Fukuoka 814-01, Japan

西郷 恵 (福岡大・理)

1. Introduction

A lot of properties are already known for various hypergeometric series in the literature. Especially, representation formulas exhibiting behaviors of the series near boundaries of their convergence regions are interesting and applicable to some calculations in [12], [13] and [27].

The present report is intended to summarize the results of the behaviors to distinguishing that for which series the behavior is known and to serve further investigation of such a calculation by finding some new ideas.

The generalized hypergeometric series of one variable is defined by

$$(1.1) \quad {}_pF_q(\alpha_1, \dots, \alpha_p; \beta_1, \dots, \beta_q; x) = \sum_{m=0}^{\infty} \frac{\prod_{j=1}^p (\alpha_j)_m}{\prod_{j=1}^q (\beta_j)_m} \frac{x^m}{m!},$$

where $(\alpha)_m$ denotes the Pochhammer symbol defined by $\Gamma(\alpha+m)/\Gamma(\alpha)$. The series (1.1) is called Gaussian series if $p = 2$, $q = 1$, and Clausenian series if $p = 3$, $q = 2$. It is well known (see e.g. [8], [26]) that the series (1.1) with $q = p-1$ converges in the region $|x| < 1$ for any complex parameters α_j and $\beta_j \notin$ nonpositive integers, as well as at $x = 1$ for $\operatorname{Re}(m) > 0$, where

$$(1.2) \quad m = \sum_{j=1}^{p-1} \beta_j - \sum_{j=1}^p \alpha_j.$$

We are interesting how the series ${}_pF_{p-1}(x)$ behaves near the point $x = 1$ when $m = 0$, for which the series is called zero-balanced. In fact, Srinivasa Ramanujan already discovered the property for the case $p = 3$ in his Notebook [9, Chapt. 11, Entry 24, Cor. 2] without proof, which may be read in the form:

$$(1.3) \quad {}_3F_2(a, b, c; d, a+b+c-d; 1-p) \\ = \frac{(a+b-d)(d-c)\Gamma(d)\Gamma(a+b+c-d)}{\Gamma(a+1)\Gamma(b+1)\Gamma(c)} {}_4F_3(1, 1, a+b-d+1, d-c+1; 2, a+1, b+1; 1) \\ - \frac{\Gamma(d)\Gamma(a+b+c-d)}{\Gamma(a)\Gamma(b)\Gamma(c)} [2\gamma + \psi(a) + \psi(b) + \log p] + o(1), \quad (p \rightarrow +0),$$

for $\operatorname{Re}(c) > 0$, where $\psi(a)$ means the psi function defined by the logarithmic derivative of the gamma function $\Gamma'(a)/\Gamma(a)$, and $\gamma = -\psi(1)$ is the Euler-Macheroni constant. Berndt express (1.3) to be "a very beautiful and significant formula" in his book [2].

The formula (1.3) implies, if we set $d = c$, the formula for the Gauss series:

$$(1.4) \quad {}_2F_1(a, b; a+b; 1-p) = -\frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} [2\gamma + \psi(a) + \psi(b) + \log p] + o(1),$$

$$(p \rightarrow +0),$$

which is a simple result from the continuation formula for the Gauss function (see [14]).

The formula (1.3) was left long mysterious fact, because Ramanujan did not refer to any proof in his Notebook, and no one could derive from the theorem (Entry 24), though it was displayed as a corollary. About the formula (1.3), Berndt mentioned in [2] as "Corollary 2 in Section 24 offers a certain asymptotic formula for zero-balanced ${}_3F_2$ series. Such formulas in the literature have previously been established only for zero-balanced ${}_2F_1$ series. It is interesting that this elegant formula had been overlooked for 60 years after Ramanujan's death." Recently, Evans and Stanton [5] and Bühring [3] proved (1.3) by their own methods. Independently of their proofs, the author established the formula (1.3) in [14], that was published earlier (1983) than the papers [5] (1984) and [3] (1987). The author did not know until 1988 that the formula (1.3) was the result by Ramanujan and left long as a mystery. The author's interest was in the direction of some calculations of the multiplication of the fractional integral and derivative which were defined by him as generalizations of that of Riemann-Liouville and Erdélyi-Kober (cf. [10], [12], [13], [27]). During the stay at the University of Victoria in Canada (March, 1988 - February, 1989), the author came to know the formula (1.3) occupied the attention and its proofs appeared in [5] and [3]. Concerning the proofs in [5] and [3], Berndt also wrote in [2] that "It would be interesting to have a more direct proof that might shed some light on Ramanujan's approach." The author is certain that his proof is just the direct one.

Moreover, the author knew at the University of Victoria that a generalization of the formula (1.3) to ${}_pF_{p-1}$ with $p \geq 4$ was posed as an open problem by Evans [4, p.553]. But the problem was solved by us, the author and H.M. Srivastava, in [24] by a similar method with that for $p = 3$ in [14]. The result for $p = 4$ is:

$$\begin{aligned}
 (1.5) \quad & {}_4F_3(a_1, a_2, a_3, a_4; b_1, b_2, b_3; 1-p) \\
 &= \frac{\Gamma(b_1)\Gamma(b_2)\Gamma(b_3)}{\Gamma(a_1+1)\Gamma(a_2+1)\Gamma(a_3)\Gamma(a_4)} (b_1-a_3)(a_1+a_2-b_1) \\
 &\quad \cdot {}_4F_3(1, 1, b_1-a_3+1, a_1+a_2-b_1+1; 2, a_1+1, a_2+1; 1) \\
 &\quad + \frac{\Gamma(b_2)\Gamma(b_3)}{\Gamma(a_4)\Gamma(b_2+b_3-a_4+1)} (b_2-a_4)(b_3-a_4) \\
 &\quad \cdot F_{1:1;1}^{0:3;3} \left[\begin{array}{c} \text{---} : a_1, a_2, a_3; b_2-a_4+1, b_3-a_4+1, 1; \\ b_2+b_3-a_4+1; \quad b_1; \quad 2; \end{array} \begin{array}{c} 1, 1 \\ 1, 1 \end{array} \right] \\
 &\quad - \frac{\Gamma(b_1)\Gamma(b_2)\Gamma(b_3)}{\Gamma(a_1)\Gamma(a_2)\Gamma(a_3)\Gamma(a_4)} [2\gamma + \psi(a_1) + \psi(a_2) + \log p] + o(1), \\
 &\hspace{25em} (p \rightarrow +0),
 \end{aligned}$$

where $a_1+a_2+a_3+a_4 = b_1+b_2+b_3$, $\operatorname{Re}(a_3) > 0$, $\operatorname{Re}(a_4) > 0$, and $F_{1:1;1}^{0:3;3}$ means the Kampé de Fériet function (see e.g. [1], [26]). The formula (1.5) implies (1.3) if $a_4 = b_3$.

For general, if $p \geq 5$ we have

$$(1.6) \quad {}_pF_{p-1}(a_1, \dots, a_p; b_1, \dots, b_{p-1}; 1-\rho)$$

$$= \frac{\Gamma(b_1) \cdots \Gamma(b_{p-1})}{\Gamma(a_1) \cdots \Gamma(a_p)} \frac{b_1 - a_3}{a_1 a_2} \left\{ \sum_{j=2}^{p-1} b_j - \sum_{j=3}^p a_j \right\} \\
\cdot {}_4F_3 \left[1, 1, b_1 - a_3 + 1, \sum_{j=2}^{p-1} b_j - \sum_{j=3}^p a_j + 1; 2, a_1 + 1, a_2 + 1; 1 \right] \\
+ \sum_{k=3}^{p-1} \frac{\Gamma(b_{k-1}) \cdots \Gamma(b_{p-1})}{\Gamma(a_{k+1}) \cdots \Gamma(a_p)} \frac{(b_{k-1} - a_{k+1}) \left\{ \sum_{j=k}^{p-1} b_j - \sum_{j=k+1}^p a_j \right\}}{\Gamma \left(\sum_{j=k-1}^{p-1} b_j - \sum_{j=k+1}^p a_j + 1 \right)} \\
\cdot {}_{F_{1:k-2;1}}^{0: k;3} \left[\begin{array}{c} \text{---} : a_1, \dots, a_k, b_{k-1} - a_{k+1} + 1, \sum_{j=k}^{p-1} b_j - \sum_{j=k+1}^p a_j + 1, 1; \\ \sum_{j=k-1}^{p-1} b_j - \sum_{j=k+1}^p a_j + 1 : b_1, \dots, b_{k-2}; \end{array} \right] \\
- \frac{\Gamma(b_1) \cdots \Gamma(b_{p-1})}{\Gamma(a_1) \cdots \Gamma(a_p)} [2\gamma + \psi(a_1) + \psi(a_2) + \log \rho] + o(1),$$

$$(\rho \rightarrow +0),$$

where $a_1 + \dots + a_p = b_1 + \dots + b_{p-1}$, $\operatorname{Re}(a_j) > 0$, ($j = 3, \dots, p$).

We introduce in the next section various hypergeometric series which are interesting to discuss through our methods, and their convergence regions of three variables are figured. For their precise definitions and the demonstration of their convergence regions, we refer to [1], [6], [7], [8] and [26]. Section 3 is devoted to mention the results obtained up to the present.

2. Hypergeometric Series and Their Convergence Regions

Single variable: As indicated in the previous section, the series ${}_pF_{p-1}(x)$ has the region of convergence $|x| < 1$ for any integer $p \geq 2$.

Two variables: The Appell series of two variables are as follows:

$$F_1(a, b_1, b_2; c; x, y) \quad \text{in} \quad |x| < 1, |y| < 1$$

$$F_2(a, b_1, b_2; c_1, c_2; x, y) \quad \text{in} \quad |x| + |y| < 1$$

$$F_3(a_1, a_2, b_1, b_2; c; x, y) \quad \text{in} \quad |x| < 1, |y| < 1$$

$$F_4(a, b; c_1, c_2; x, y) \quad \text{in} \quad \sqrt{|x|} + \sqrt{|y|} < 1$$

Three variables: Let $|x| = r$, $|y| = s$, and $|z| = t$, the convergence regions of triple hypergeometric series of Lauricella and Srivastava are shown as follows with figures:

$$F_A(a, a, a, b_1, b_2, b_3; c_1, c_2, c_3; x, y, z) \quad r+s+t < 1$$

$$F_B(a_1, a_2, a_3, b_1, b_2, b_3; c, c, c; x, y, z) \quad r < 1, s < 1, t < 1$$

$$F_C(a, a, a, b, b, b; c_1, c_2, c_3; x, y, z) \quad \sqrt{r} + \sqrt{s} + \sqrt{t} < 1$$

$$F_D(a, a, a, b_1, b_2, b_3; c, c, c; x, y, z) \quad r < 1, s < 1, t < 1$$

$$F_E(a, a, a, b_1, b_2, b_2; c_1, c_2, c_3; x, y, z)$$

$$r < 1 - (\sqrt{s} + \sqrt{t})^2$$

$$F_F(a, a, a, b_1, b_2, b_1; c_1, c_2, c_2; x, y, z)$$

$$\sqrt{r} + \sqrt{t} < 1, \quad s < \frac{1}{2}(1+t-r+\sqrt{(1+t-r)^2-4t})$$

$$F_G(a, a, a, b_1, b_2, b_3; c_1, c_2, c_2; x, y, z)$$

$$r+s < 1, \quad r+t < 1$$

$$F_K(a_1, a_2, a_2, b_1, b_2, b_1; c_1, c_2, c_3; x, y, z)$$

$$r < 1, \quad s < 1, \quad t < (1-s)(1-r)$$

$$F_M(a_1, a_2, a_2, b_1, b_2, b_1; c_1, c_2, c_2; x, y, z)$$

$$r+t < 1, \quad s < 1$$

$$F_N(a_1, a_2, a_3, b_1, b_2, b_1; c_1, c_2, c_2; x, y, z)$$

$$r+t < 1, \quad s < 1$$

$$F_P(a_1, a_2, a_1, b_1, b_1, b_2; c_1, c_2, c_2; x, y, z)$$

$$r+s < 1, \quad r+t < 1$$

$$F_R(a_1, a_2, a_1, b_1, b_2, b_1; c_1, c_2, c_2; x, y, z)$$

$$\sqrt{r} + \sqrt{t} < 1, \quad s < 1$$

$$F_S(a_1, a_2, a_2, b_1, b_2, b_3; c, c, c; x, y, z)$$

$$r < 1, \quad s < 1, \quad t < 1$$

$$F_T(a_1, a_2, a_2, b_1, b_2, b_1; c, c, c; x, y, z)$$

$$r < 1, \quad s < 1, \quad t < 1$$

$$H_A(a_1, a_2, a_3; c_1, c_2; x, y, z)$$


$$r < (1-s)(1-t), \quad s < 1, \quad t < 1$$

$$H_B(a_1, a_2, a_3; c_1, c_2, c_3; x, y, z)$$


$$r+s+t+2\sqrt{rst} < 1$$

$$H_C(a_1, a_2, a_3; c; x, y, z)$$


$$r < 1, \quad s < 1, \quad t < 1, \quad r+s+t-2\sqrt{(1-r)(1-s)(1-t)} < 2$$




$$F_A: r+s+t < 1$$




$$F_B: r < 1, s < 1, t < 1$$




$$F_C: \sqrt{r} + \sqrt{s} + \sqrt{t} < 1$$









$$F_D: r < 1, s < 1, t < 1$$




$$F_E: r + (\sqrt{s} + \sqrt{t})^2 < 1$$




$$F_F: \sqrt{r} + \sqrt{t} < 1, s < \frac{1}{2} \{ 1 + t - r + \sqrt{(1+t-r)^2 - 4t} \}$$



 $F_G: r+s < 1, r+t < 1$

 $F_K: r < 1, s < 1, t < (1-r)(1-s)$

 $F_M: r+t < 1, s < 1$

 $F_N: r+t < 1, s < 1$

 $F_P: r+s < 1, r+t < 1$

 $F_R: \sqrt{r} + \sqrt{t} < 1, s < 1$




$$F_S: r < 1, s < 1, t < 1$$




$$F_T: r < 1, s < 1, t < 1$$



$$H_A: r < (1-s)(1-t), s < 1, t < 1$$



$$H_B: r + s + t + 2\sqrt{rst} < 1$$



$$H_C: r < 1, s < 1, t < 1, r + s + t - 2\sqrt{(1-r)(1-s)(1-t)} < 2$$

Multivariables: Let $|x_i| = r_i$, ($i = 1, \dots, n$), then the convergence regions of Lauricella hypergeometric series of n variables are as follows:

$$F_A^{(n)}(a, b_1, \dots, b_n; c_1, \dots, c_n; x_1, \dots, x_n) \quad r_1 + \dots + r_n < 1$$

$$F_B^{(n)}(a_1, \dots, a_n, b_1, \dots, b_n; c; x_1, \dots, x_n) \quad r_1 < 1, \dots, r_n < 1$$

$$F_C^{(n)}(a, b; c_1, \dots, c_n; x_1, \dots, x_n) \quad \sqrt{r_1} + \dots + \sqrt{r_n} < 1$$

$$F_D^{(n)}(a, b_1, \dots, b_n; c; x_1, \dots, x_n) \quad r_1 < 1, \dots, r_n < 1$$

3. Properties of Hypergeometric Series

We shall state here whole possibility of the boundary points of convergence regions of hypergeometric series introduced in the previous section. The reference numbers indicate that their behaviors were already calculated. The term "complete" means the formula is known with no restriction on parameters without some zero-balancedness, and "2 cases" means that two formulas are evaluated with some other restrictions. The calculations are tried only for the listed Gaussian complete series. The cases which are not indicated the reference numbers are open problems for the reader.

Single variable:

$${}_2F_1[1-p] \text{ complete: } [14]$$

$${}_3F_2[1-p] \text{ complete: } [14]$$

$p^F_{p-1}[1-p]$ complete: [24]

Two variables:

$F_1[1-p, y]$ complete: included in [15]

$F_1[1-px, 1-py]$ complete: [11]

$F_2[x, 1-x-p]$ 2 cases: [14] complete: [20]

$F_3[1-p, y]$ complete: [14]

$F_3[1-px, 1-py]$ complete: included in [23]

$F_4[x, (1-\sqrt{y}-p)^2]$ 3 cases: [21]

Three variables:

$F_A[x, y, 1-x-y-p]$ complete: included in [18]

$F_B[x, y, 1-p]$ complete: included in [23]

$F_B[x, 1-py, 1-pz]$ complete: included in [23]

$F_B[1-px, 1-py, 1-pz]$ complete: included in [23]

$$F_C[x, y, (1 - \sqrt{x} - \sqrt{y} - \rho)^2] \quad 1 \text{ case:} \quad [21]$$

$$F_D[x, y, 1 - \rho] \quad \text{complete:} \quad [15]$$

$$F_D[x, 1 - \rho y, 1 - \rho z] \quad \text{complete:} \quad [15]$$

$$F_D[1 - \rho x, 1 - \rho y, 1 - \rho z] \quad \text{complete:} \quad [15]$$

$$F_E[1 - (\sqrt{y} + \sqrt{z})^2 - \rho, y, z]$$

$$F_F[x, y, (1 - \sqrt{x} - \rho)^2]$$

$$F_F[x, \frac{1}{2}(1 - x + z + \sqrt{(1 - x + z)^2 - 4z} - \rho), z]$$

$$F_F[x, 1 - \sqrt{x} - \rho y, (1 - \sqrt{x} - \rho z)^2]$$

$$F_G[x, y, 1 - x - \rho] \quad 2 \text{ cases:} \quad [16] \quad \text{complete:} \quad [17]$$

$$F_G[x, 1 - x - \rho, 1 - x - \rho] \quad \text{complete:} \quad [17]$$

$$F_K[x, y, (1 - x)(1 - y) - \rho] \quad 1 \text{ case:} \quad [17]$$

$$F_M[x, 1 - \rho, z] \quad \text{complete:} \quad [16]$$

$$F_M[x, y, 1 - x - \rho] \quad 1 \text{ case:} \quad [16]$$

$$F_M[x, 1-\rho y, 1-x-\rho z]$$

$$F_N[x, 1-\rho, z] \quad \text{complete:} \quad [17]$$

$$F_N[x, y, 1-x-\rho] \quad 1 \text{ case:} \quad [21] \quad \text{complete:} \quad [17]$$

$$F_N[x, 1-\rho y, 1-x-\rho z]$$

$$F_P[x, y, 1-x-\rho]$$

$$F_P[x, 1-x-\rho y, 1-x-\rho z]$$

$$F_R[x, y, (1-\sqrt{x}-\rho)^2] \quad 1 \text{ case:} \quad [21]$$

$$F_R[x, 1-\rho, z] \quad \text{complete:} \quad [17]$$

$$F_R[x, 1-\rho y, (1-\sqrt{x}-\rho z)^2]$$

$$F_S[x, y, 1-\rho] \quad \text{complete:} \quad [25]$$

$$F_S[x, 1-\rho y, 1-\rho z] \quad \text{complete:} \quad [25]$$

$$F_S[1-\rho x, 1-\rho y, 1-\rho z] \quad \text{complete:} \quad [25]$$

$$F_S[1-\rho, y, z] \quad \text{complete:} \quad [25]$$

$$F_S[1-\rho x, 1-\rho y, z] \quad \text{complete:} \quad [25]$$

$F_T[x, y, 1-p]$ complete: [25]

$F_T[x, 1-py, 1-pz]$ complete: [25]

$F_T[1-px, 1-py, 1-pz]$

$F_T[1-p, y, z]$ complete: [25]

$F_T[1-px, 1-py, z]$ complete: [25]

$H_A[(1-y)(1-z)-p, y, z]$

$H_B[x, y, z], \quad x+y+z+2\sqrt{xyz} = 1-p$

$H_C[1-p, y, z]$ complete: [19]

$H_C[1-px, y, 1-y-p]$

$H_C[x, y, z], \quad x+y+z-2\sqrt{(1-x)(1-y)(1-z)} = 2-p$

Multivariables:

$F_A^{(n)}[x_1, \dots, x_{n-1}, 1-x_1-\dots-x_{n-1}-p]$ complete: [18]

$F_B^{(n)}[x_1, \dots, x_k, 1-px_{k+1}, \dots, 1-px_n]$ complete: [23]

$$F_C^{(n)} [x_1, \dots, x_{n-1}, (\sqrt{|x_1|} + \dots + \sqrt{|x_{n-1}|})^2 - p]$$

$$F_D^{(n)} [x_1, \dots, x_k, 1 - px_{k+1}, \dots, 1 - px_n] \text{ complete: } [22]$$

References

- [1] P. Appell et J. Kampé de Fériet: Fonctions Hypergéométriques et Hypersphériques, Polynômes d'Hermite, Gauthier-Villars, Paris, 1926.
- [2] B.C. Berndt: Ramanujan's Notebooks, Part II, Springer-Verlag, New York, Berlin, Heidelberg, London, Paris and Tokyo, 1989.
- [3] W. Bühring: The behavior at unit argument of the hypergeometric function ${}_3F_2$, SIAM J. Math. Anal., **18**(1987), 1227-1234.
- [4] R.J. Evans: Ramanujan's second notebook; Asymptotic expansions for hypergeometric series and related functions, Ramanujan Revisited (G.E. Andrews et al. edited: Proc. of the Ramanujan Centenary Conference), 537-560, Academic Press, New York, 1988.
- [5] R.J. Evans and D. Stanton: Asymptotic formulas for zero-balanced hypergeometric series, SIAM J. Math. Anal., **15**(1984), 1010-1020.
- [6] H. Exton: Multiple Hypergeometric Functions and Applications, Halsted Press (Ellis Horwood, Chichester), John Wiley and Sons, New York, London, Sidney and Toronto, 1976.
- [7] G. Lauricella: Sulle funzioni ipergeometriche a più variabili, Rend. Circ. Mat. Palermo, **7**(1893), 111-158.

- [8] W. Magnus, F. Oberhettinger and R.P. Soni: Formulas and Theorems for the Special Functions of Mathematical Physics, Third enlarged ed., Springer-Verlag, New York, 1966.
- [9] S. Ramanujan: Notebooks of Srinivasa Ramanujan, Vol.2, Tata Inst. of Fund. Res., Bombay, 1957.
- [10] M. Saigo: A remark on integral operators involving the Gauss hypergeometric functions, Math. Rep. Kyushu Univ., **11**(1978), 135-143.
- [11] M. Saigo: On a property of the Appell hypergeometric function F_1 , Math. Rep. Kyushu Univ., **12**(1980), 63-67.
- [12] M. Saigo: A certain boundary value problem for the Euler-Darboux equation, II, Math. Japon., **25**(1980), 211-220.
- [13] M. Saigo: A certain boundary value problem for the Euler-Darboux equation, III, Math. Japon., **26**(1981), 103-119.
- [14] M. Saigo: On properties of the Appell hypergeometric functions F_2 and F_3 and the generalized Gauss function ${}_3F_2$, Bull. Central Res. Inst. Fukuoka Univ., **66**(1983), 27-32.
- [15] M. Saigo: On properties of the Lauricella hypergeometric functions F_D , Bull. Central Res. Inst. Fukuoka Univ., **104**(1988), 13-31.
- [16] M. Saigo: On properties of hypergeometric functions of three variables, F_M and F_G , Rend. Circ. Mat. Palermo (2), **37**(1988), 449-468.
- [17] M. Saigo: The asymptotic behaviors of triple hypergeometric series, F_G , F_K , F_N and F_R near boundaries of their convergence regions, (Kyoto, 1989), Sûrikaiseikikenkyûsho Kôkyûroku, this volume (1990).
- [18] M. Saigo: The behaviors of the Lauricella hypergeometric series $F_A^{(n)}$ in n variables near the boundaries of its convergence region, (in preparation).

- [19] M. Saigo: The asymptotic behavior of the Srivastava hypergeometric series of three variables H_C near the boundary of its convergence region, (in preparation).
- [20] M. Saigo, O.I. Marichev and Nguyen Thanh Hai: Asymptotic representations of Gaussian series ${}_2F_1$, Clausenian series ${}_3F_2$ and Appell series F_2 and F_3 near boundaries of their convergence regions, Fukuoka Univ. Sci. Rep., **19**(1989), 83-90.
- [21] M. Saigo and H.M. Srivastava: The behaviors of the Appell double hypergeometric series F_4 and certain Lauricella triple hypergeometric series near the boundaries of their convergence regions, Fukuoka Univ. Sci. Rep., **19**(1989), 1-10.
- [22] M. Saigo and H.M. Srivastava: The behaviors of the Lauricella hypergeometric series $F_D^{(n)}$ in n variables near the boundary of its convergence region, Univ. Victoria Rep., No. DM-480-IR, 1-23, (1988).
- [23] M. Saigo and H.M. Srivastava: The asymptotic behaviour of Lauricella's second hypergeometric series $F_B^{(n)}$ in n variables near the boundaries of its convergence region, Fukuoka Univ. Sci. Rep., **19**(1989), 71-82.
- [24] M. Saigo and H.M. Srivastava: The behaviors of the zero-balanced hypergeometric series ${}_pF_{p-1}$ near the boundary of its convergence region, Proc. Amer. Math. Soc., (to appear).
- [25] M. Saigo and H.M. Srivastava: Some asymptotic formulas exhibiting the behaviors of the triple hypergeometric series F_S and F_T near the boundaries of their convergence regions, Univ. Victoria Rep., No. DMS-515-IR, 1-38, (1989).
- [26] H.M. Srivastava and P.W. Karlsson: Multiple Gaussian Hypergeometric Series, Halsted Press (Ellis Horwood Ltd., Chichester), John Wiley and Sons, New York, Chichester, Brisbane and Toronto, 1985.

- [27] H.M. Srivastava and M. Saigo: Multiplication of fractional calculus operators and boundary value problems involving the Euler-Darboux equation, J. Math. Anal. Appl., **121**(1987), 325-369.