

Homocoupling of aryl halides in flow: Space integration of lithiation and FeCl₃ promoted homocoupling

Aiichiro Nagaki, Yuki Uesugi, Yutaka Tomida and Jun-ichi Yoshida*

Full Research Paper

Open Access

Address:

Department of Synthetic Chemistry and Biological Chemistry,
Graduate School of Engineering, Kyoto University,
Kyotodaigakukatsura, Nishikyo-ku, Kyoto, 615-8510, Japan

Email:

Aiichiro Nagaki - anagaki@sbchem.kyoto-u.ac.jp; Jun-ichi Yoshida* -
yoshida@sbchem.kyoto-u.ac.jp

* Corresponding author

Keywords:

homocoupling; iron salts; microreactor; organolithiums

Beilstein J. Org. Chem. **2011**, *7*, 1064–1069.

doi:10.3762/bjoc.7.122

Received: 15 April 2011

Accepted: 29 June 2011

Published: 02 August 2011

This article is part of the Thematic Series "Chemistry in flow systems II".

Guest Editor: A. Kirschning

© 2011 Nagaki et al; licensee Beilstein-Institut.

License and terms: see end of document.

Abstract

The use of FeCl₃ resulted in a fast homocoupling of aryllithiums, and this enabled its integration with the halogen–lithium exchange reaction of aryl halides in a flow microreactor. This system allows the homocoupling of two aryl halides bearing electrophilic functional groups, such as CN and NO₂, in under a minute.

Introduction

Biaryl structures often occur in various organic compounds including natural products, bioactive compounds, functional polymers, ligands in catalysts and theoretically interesting molecules, and the oxidative homocoupling of arylmetals is one of the most useful methods for the construction of biaryl frameworks [1]. Stoichiometric amounts of transition metal salts such as TiCl₄ [2], TiCl₃ [3], VO(OEt)Cl₂ [4], CoCl₂ [5], CuCl₂ [6] and Pd(OAc)₂ [7] have been used for homocoupling of arylmetals. In some cases catalytic processes in the presence of a reoxidant, such as oxygen or other organic oxidants, are effective. Recently, iron salts have been also used because of their low costs and lack of toxicity [8–18]. For example, Hayashi et al. reported the iron-catalyzed oxidative homocoupling of Grignard reagents, using 1,2-dihalogenoethanes as an oxidant [19]. Cahiez et al. have also reported the FeCl₃-catalyzed homocou-

pling reaction of Grignard reagents bearing functional groups, using atmospheric oxygen [20]. The use of aryllithium compounds instead of Grignard reagents is very interesting, because they are easily generated by halogen–lithium exchange under homogeneous conditions, thus enabling the generation in a flow. However, to the best of our knowledge, oxidative homocoupling of aryllithiums using iron salts has not been reported so far. One of the major reasons for this seems to be the instability of aryllithiums, especially of those bearing electrophilic functional groups such as cyano and nitro groups [21], making the subsequent homocoupling difficult or impossible.

Recently, we have reported that flow microreactor systems [22–85] are quite effective for the generation and reaction of highly reactive organolithiums such as functionalized aryllithiums,

oxiranyllithiums, aziridinyllithiums, and allenyllithiums [86–98]. Herein we report that integration [99,100] of the generation of aryllithiums, especially those bearing electrophilic functional groups, by halogen–lithium exchange and FeCl_3 promoted homocoupling has been effectively accomplished in an integrated flow microreactor system.

Results and Discussion

First, we focused on the generation of *p*-methoxyphenyllithium from *p*-bromoanisole (Scheme 1). A flow microreactor system, consisting of two T-shaped micromixers (**M1** and **M2**) and two microtube reactors (**R1** and **R2**) shown in Figure 1, was used. A solution of *p*-bromoanisole (Ar-X) (0.10 M in THF, flow rate: 6.0 mL/min) and a solution of *n*-butyllithium (0.40 M in hexane, flow rate: 1.5 mL/min) were introduced to **M1** ($\varnothing = 250 \mu\text{m}$) by syringe pumps. The resulting mixture was passed through **R1** to conduct the bromine–lithium exchange reaction. Methanol (neat, flow rate: 3.0 mL/min) was added in **M2** ($\varnothing = 500 \mu\text{m}$) and the mixture was passed through **R2** ($\varnothing = 1000 \mu\text{m}$, $L = 50 \text{ cm}$) to protonate *p*-methoxyphenyllithium. The reactions were carried out with varying residence time in **R1** (t^{R1} : 0.2–6.3 s) and varying temperature (T : -78 to 24°C). The temperature (T) was controlled by adjusting the bath temperature. The residence time (t^{R1}) was adjusted by changing the inner diameter and the length in the microtube reactor **R1** with a fixed flow rate. After a steady state was reached, the product solution was collected for 30 s. As shown in Figure 2, the yield of the protonated product, anisole, depends on both T and t^{R1} . The reaction at low temperatures ($T < -48^\circ\text{C}$) with short residence times ($t^{\text{R1}} < 0.79 \text{ s}$) resulted in very low yields, because the Br–Li exchange reaction was not complete. The increase in T and t^{R1} caused an increase in the yield, and high yields ($>85\%$) were obtained through the appropriate choice of T and t^{R1} .

Next, we examined the integration of the halogen–lithium exchange reaction with FeCl_3 promoted homocoupling (Scheme 2). Integrated flow microreactor systems consisting of three micromixers (**M1**, **M2**, and **M3**) and three microtube reac-




Figure 1: Flow microreactor system for halogen–lithium exchange of aryl halide followed by reaction with methanol. T-shaped micromixer: **M1** (inner diameter: $250 \mu\text{m}$), and **M2** (inner diameter: $500 \mu\text{m}$), microtube reactor: **R1** and **R2** ($\varnothing = 1000 \mu\text{m}$, length = 50 cm), a solution of aryl halides: 0.10 M in THF (6.0 mL/min), a solution of lithium reagent: 0.40 M or 0.42 M in hexane (*n*-BuLi) or Et_2O (PhLi) (1.5 mL/min), a solution of methanol: Neat (3.0 mL/min).






Figure 2: Effects of the temperature (T) and the residence time in **R1** (t^{R1}) on the yield of anisole in the Br–Li exchange reaction of *p*-bromoanisole followed by reaction with methanol in the flow microreactor system. The contour plot with scatter overlay shows the yields of anisole (%), which are indicated by small circles.





Scheme 1: Halogen–lithium exchange of *p*-bromoanisole followed by reaction with methanol.



Scheme 2: Halogen–lithium exchange of *p*-bromoanisole followed by oxidative homocoupling with FeCl_3 .

tors (**R1**, **R2**, and **R3**) were used, as shown in Figure 3. A solution of *p*-bromoanisole (Ar-X) (0.10 M in THF, flow rate: 6.0 mL/min) and a solution of *n*-butyllithium (0.42 M in hexane, flow rate: 1.5 mL/min) were introduced to **M1** ($\varnothing = 250\ \mu\text{m}$) by syringe pumps. The resulting mixture was passed through **R1** ($t^{\text{R1}} = 13\ \text{s}$ ($-78\ ^\circ\text{C}$), $t^{\text{R1}} = 13\ \text{s}$ ($-48\ ^\circ\text{C}$), $t^{\text{R1}} = 3.1\ \text{s}$ ($-28\ ^\circ\text{C}$), $t^{\text{R1}} = 3.1\ \text{s}$ ($0\ ^\circ\text{C}$), $t^{\text{R1}} = 3.1\ \text{s}$ ($24\ ^\circ\text{C}$)) at the corresponding temperatures and was mixed with a solution of FeCl_3 (0.10 M in THF, flow rate: 6.0 mL/min) in **M2** ($\varnothing = 500\ \mu\text{m}$). The resulting mixture was passed through **R2** and was then mixed with methanol (neat, flow rate: 1.5 mL/min) in **M3** ($\varnothing = 500\ \mu\text{m}$) to protonate the unchanged *p*-methoxyphenyllithium. The resulting solution was passed through **R3** ($\varnothing = 1000\ \mu\text{m}$, $L = 50\ \text{cm}$). The temperature (T) was controlled by adjusting the bath temperature, and the residence time in **R2** (t^{R2}) by changing the inner diameter and the length in **R2** with the fixed flow rate. After a steady state was reached, the product solution was collected for 30 s. The results obtained from varying t^{R2} and T are summarized in Figure 4, in which the yield of 4,4'-dimethoxybiphenyl is plotted against T and t^{R2} as a contour map with scattered overlay (see Supporting Information File 1 for details). The yield depends on both T and t^{R2} . At $-78\ ^\circ\text{C}$, the yield increased with t^{R2} because of the progress of the homocoupling. At $0\ ^\circ\text{C}$, the homocoupling product was obtained in reasonable yields for a wide range of t^{R2} . The productivity of the present system is acceptable for large scale laboratory synthesis (6.2 g/h). It is noteworthy that the integrated reactions were complete within the overall residence time of 14.7 s, even at low temperatures such as $-48\ ^\circ\text{C}$. Thus, we envisaged that the reaction could also be applied to less stable aryllithium compounds that decompose very quickly.

One of the major benefits of flow microreactor synthesis is the ability to use highly unstable reactive intermediates. Such inter-



















mediates can be rapidly generated and transferred to another location to be used in a subsequent reaction before they decompose. We have already reported the generation and reactions of unstable aryllithium species such as *o*-bromophenyllithiums, and aryllithiums bearing alkoxycarbonyl, cyano, nitro, and ketone carbonyl groups [86,87,93,95,96,98], which are difficult to use in conventional macro batch reactors. As shown in Table 1, reactions of aryllithiums bearing cyano and nitro groups proceeded successfully to give the corresponding homocoupling products, where in contrast it is very difficult to achieve such reactions using conventional batch reactors. A mechanism involving transmetalation of the aryl group from lithium to iron followed by reductive elimination of the homocoupling product seems to be plausible, while a similar mechanism is proposed for homo-coupling of organomagnesium compounds with FeCl_3 [19,20]. The regioselectivity of the coupling is consistent with this mechanism. Radical coupling seems to be less likely.

Conclusion

In conclusion, we found that the use of FeCl_3 results in fast oxidative homocoupling of aryllithiums, which enables its integration with the halogen–lithium exchange of aryl halides. Various aryl halides, including those bearing electrophilic functional groups, can be used for this transformation in the integrated flow microreactor system. Hence, the method greatly enhances the synthetic utility of aryllithium compounds and adds a new dimension to the chemistry of coupling reactions.

Table 1: Homocoupling of aryl halides using the integrated flow microreactor system.

Ar–X	T (°C)	t ^{R1} (s)	Ar–Ar	Yield (%)
	24	3.100		72
	24	3.100		69
	0	3.100		76
	–28	0.055		75
	0	0.055		66
	24	0.055		76
	–48	0.014		53 ^a
	–48	0.014		63 ^a

^aPhLi instead of *n*-BuLi was used as lithiating reagent.

Supporting Information

Supporting Information features experimental procedures and full spectroscopic data for all new compounds.

Supporting Information File 1

Experimental details.

[<http://www.beilstein-journals.org/bjoc/content/supplementary/1860-5397-7-122-S1.pdf>]

Acknowledgements

This work was financially supported in part by a Grant-in-Aid for Scientific Research from the Japan Society for the Promotion of Science and NEDO projects.

References

- Cepanec, I. *Synthesis of Biaryls*; Elsevier: Amsterdam, Boston, 2004.
- Inoue, A.; Kitagawa, K.; Shinokubo, H.; Oshima, K. *Tetrahedron* **2000**, *56*, 9601–9605. doi:10.1016/S0040-4020(00)00929-7
- McKillop, A.; Elsom, L. F.; Taylor, E. C. *J. Am. Chem. Soc.* **1968**, *90*, 2423–2424. doi:10.1021/ja01011a041
- Ishikawa, T.; Ogawa, A.; Hirao, T. *Organometallics* **1998**, *17*, 5713–5716. doi:10.1021/om980607c
- Kharasch, M. S.; Fields, E. K. *J. Am. Chem. Soc.* **1941**, *63*, 2316–2320. doi:10.1021/ja01854a006
- Sakellarios, E.; Kyrimis, T. *Ber. Dtsch. Chem. Ges.* **1924**, *57*, 322–326. doi:10.1002/cber.19240570233
- Lei, A.; Srivastava, M.; Zhang, X. *J. Org. Chem.* **2002**, *67*, 1969–1971. doi:10.1021/jo011098i
- Cahiez, G.; Marquais, S. *Pure Appl. Chem.* **1996**, *68*, 53–60. doi:10.1351/pac199668010053
- Cahiez, G.; Marquais, S. *Tetrahedron Lett.* **1996**, *37*, 1773–1776. doi:10.1016/0040-4039(96)00116-5

10. Cahiez, G.; Avedissian, H. *Synthesis* **1998**, 1199–1205. doi:10.1055/s-1998-2135
11. Dohle, W.; Kopp, F.; Cahiez, G.; Knochel, P. *Synlett* **2001**, 1901–1904. doi:10.1055/s-2001-18748
12. Fürstner, A.; Leitner, A.; Méndez, M.; Krause, H. *J. Am. Chem. Soc.* **2002**, *124*, 13856–13863. doi:10.1021/ja027190t
13. Fürstner, A.; Leitner, A. *Angew. Chem., Int. Ed.* **2002**, *41*, 609–612. doi:10.1002/1521-3773(20020215)41:4<609::AID-ANIE609>3.0.CO;2-M
14. Quintin, J.; Franck, X.; Hocquemiller, R.; Figadère, B. *Tetrahedron Lett.* **2002**, *43*, 3547–3549. doi:10.1016/S0040-4039(02)00568-3
15. Martin, R.; Fürstner, A. *Angew. Chem., Int. Ed.* **2004**, *43*, 3955–3957. doi:10.1002/anie.200460504
16. Scheiper, B.; Bonnekessel, M.; Krause, H.; Fürstner, A. *J. Org. Chem.* **2004**, *69*, 3943–3949. doi:10.1021/jo0498866
17. Nakamura, M.; Matsuo, K.; Ito, S.; Nakamura, E. *J. Am. Chem. Soc.* **2004**, *126*, 3686–3687. doi:10.1021/ja049744t
18. Nagano, T.; Hayashi, T. *Org. Lett.* **2004**, *6*, 1297–1299. doi:10.1021/ol049779y
19. Nagano, T.; Hayashi, T. *Org. Lett.* **2005**, *7*, 491–493. doi:10.1021/ol047509+
20. Cahiez, G.; Chaboche, C.; Mahuteau-Betzer, F.; Ahr, M. *Org. Lett.* **2005**, *7*, 1943–1946. doi:10.1021/ol050340v
21. Knochel, P., Ed. *Handbook of Functionalized Organometallics*; Wiley-VCH: Weinheim, Germany, 2005. doi:10.1002/9783527619467
22. Yoshida, J. *Flash Chemistry: Fast Organic Synthesis in Microsystems*; Wiley: Hoboken, N.J., 2008.
23. Fletcher, P. D. I.; Haswell, S. J.; Pombo-Villar, E.; Warrington, B. H.; Watts, P.; Wong, S. Y. F.; Zhang, X. *Tetrahedron* **2002**, *58*, 4735–4757. doi:10.1016/S0040-4020(02)00432-5
24. Jähnisch, K.; Hessel, V.; Löwe, H.; Baerns, M. *Angew. Chem., Int. Ed.* **2004**, *43*, 406–446. doi:10.1002/anie.200300577
25. Yoshida, J. *Chem. Commun.* **2005**, 4509–4516. doi:10.1039/b508341a
26. Doku, G. N.; Verboom, W.; Reinhoudt, D. N.; van den Berg, A. *Tetrahedron* **2005**, *61*, 2733–2742. doi:10.1016/j.tet.2005.01.028
27. Yoshida, J.; Nagaki, A.; Iwasaki, T.; Suga, S. *Chem. Eng. Technol.* **2005**, *28*, 259–266. doi:10.1002/ceat.200407127
28. Geyer, K.; Codee, J. D. C.; Seeberger, P. H. *Chem.–Eur. J.* **2006**, *12*, 8434–8442. doi:10.1002/chem.200600596
29. Whitesides, G. *Nature* **2006**, *442*, 368–373. doi:10.1038/nature05058
30. deMello, A. J. *Nature* **2006**, *442*, 394–402. doi:10.1038/nature05062
31. Song, H.; Chen, D. L.; Ismagilov, R. F. *Angew. Chem., Int. Ed.* **2006**, *45*, 7336–7356. doi:10.1002/anie.200601554
32. Kobayashi, J.; Mori, Y.; Kobayashi, S. *Chem.–Asian J.* **2006**, *1*, 22–35. doi:10.1002/asia.200600058
33. Mason, B. P.; Price, K. E.; Steinbacher, J. L.; Bogdan, A. R.; McQuade, D. T. *Chem. Rev.* **2007**, *107*, 2300–2318. doi:10.1021/cr050944c
34. Ahmed-Omer, B.; Brandtand, J. C.; Wirth, T. *Org. Biomol. Chem.* **2007**, *5*, 733–740. doi:10.1039/b615072a
35. Watts, P.; Wiles, C. *Chem. Commun.* **2007**, 443–467. doi:10.1039/b609428g
36. Yoshida, J.; Nagaki, A.; Yamada, T. *Chem.–Eur. J.* **2008**, *14*, 7450–7459. doi:10.1002/chem.200800582
37. Fukuyama, T.; Rahman, M. T.; Sato, M.; Ryu, I. *Synlett* **2008**, 151–163. doi:10.1055/s-2007-1000884
38. McMullen, J. P.; Jensen, K. F. *Annu. Rev. Anal. Chem.* **2010**, *3*, 19–42. doi:10.1146/annurev.anchem.111808.073718
39. Yoshida, J. *Chem. Rec.* **2010**, *10*, 332–341. doi:10.1002/tcr.201000020
40. Yoshida, J.; Kim, H.; Nagaki, A. *ChemSusChem* **2011**, *4*, 331–340. doi:10.1002/cssc.201000271
41. Watts, P.; Wiles, C.; Haswell, S. J.; Pombo-Villar, E.; Styring, P. *Chem. Commun.* **2001**, 990–991. doi:10.1039/b102125g
42. Suga, S.; Okajima, M.; Fujiwara, K.; Yoshida, J. *J. Am. Chem. Soc.* **2001**, *123*, 7941–7942. doi:10.1021/ja015823i
43. Fukuyama, T.; Shinmen, M.; Nishitani, S.; Sato, M.; Ryu, I. *Org. Lett.* **2002**, *4*, 1691–1694. doi:10.1021/ol0257732
44. Ueno, M.; Hisamoto, H.; Kitamori, T.; Kobayashi, S. *Chem. Commun.* **2003**, 936–937. doi:10.1039/b301638b
45. Garcia-Egido, E.; Spikmans, V.; Wong, S. Y. F.; Warrington, B. H. *Lab Chip* **2003**, *3*, 73–76. doi:10.1039/b302381h
46. Lai, S. M.; Martin-Aranda, R.; Yeung, K. L. *Chem. Commun.* **2003**, 218–219. doi:10.1039/b209297b
47. Mikami, K.; Yamanaka, M.; Islam, M. N.; Kudo, K.; Seino, N.; Shinoda, M. *Tetrahedron Lett.* **2003**, *44*, 7545–7548. doi:10.1016/S0040-4039(03)01835-5
48. Kobayashi, J.; Mori, Y.; Okamoto, K.; Akiyama, R.; Ueno, M.; Kitamori, T.; Kobayashi, S. *Science* **2004**, *304*, 1305–1308. doi:10.1126/science.1096956
49. Nagaki, A.; Kawamura, K.; Suga, S.; Ando, T.; Sawamoto, M.; Yoshida, J. *J. Am. Chem. Soc.* **2004**, *126*, 14702–14703. doi:10.1021/ja044879k
50. Lee, C.-C.; Sui, G.; Elizarov, A.; Shu, C. J.; Shin, Y.-S.; Dooley, A. N.; Huang, J.; Daridon, A.; Wyatt, P.; Stout, D.; Kolb, H. C.; Witte, O. N.; Satyamurthy, N.; Heath, J. R.; Phelps, M. E.; Quake, S. R.; Tseng, H.-R. *Science* **2005**, *310*, 1793–1796. doi:10.1126/science.1118919
51. Horcajada, R.; Okajima, M.; Suga, S.; Yoshida, J. *Chem. Commun.* **2005**, 1303–1305. doi:10.1039/b417388k
52. Kawaguchi, T.; Miyata, H.; Ataka, K.; Mae, K.; Yoshida, J. *Angew. Chem., Int. Ed.* **2005**, *44*, 2413–2416. doi:10.1002/anie.200462466
53. Ducry, L.; Roberge, D. M. *Angew. Chem., Int. Ed.* **2005**, *44*, 7972–7975. doi:10.1002/anie.200502387
54. Nagaki, A.; Togai, M.; Suga, S.; Aoki, N.; Mae, K.; Yoshida, J. *J. Am. Chem. Soc.* **2005**, *127*, 11666–11675. doi:10.1021/ja0527424
55. He, P.; Watts, P.; Marken, F.; Haswell, S. J. *Angew. Chem., Int. Ed.* **2006**, *45*, 4146–4149. doi:10.1002/anie.200600951
56. Uozumi, Y.; Yamada, Y.; Beppu, T.; Fukuyama, N.; Ueno, M.; Kitamori, T. *J. Am. Chem. Soc.* **2006**, *128*, 15994–15995. doi:10.1021/ja066697r
57. Tanaka, K.; Motomatsu, S.; Koyama, K.; Tanaka, S.; Fukase, K. *Org. Lett.* **2007**, *9*, 299–302. doi:10.1021/ol062777o
58. Sahoo, H. R.; Kralj, J. G.; Jensen, K. F. *Angew. Chem., Int. Ed.* **2007**, *46*, 5704–5708. doi:10.1002/anie.200701434
59. Hornung, C. H.; Mackley, M. R.; Baxendale, I. R.; Ley, S. V. *Org. Process Res. Dev.* **2007**, *11*, 399–405. doi:10.1021/op700015f
60. Iwasaki, T.; Nagaki, A.; Yoshida, J. *Chem. Commun.* **2007**, 1263–1265. doi:10.1039/b615159k
61. Nagaki, A.; Iwasaki, T.; Kawamura, K.; Yamada, D.; Suga, S.; Ando, T.; Sawamoto, M.; Yoshida, J. *Chem.–Asian J.* **2008**, *3*, 1558–1567. doi:10.1002/asia.200800081
62. Nagaki, A.; Tomida, Y.; Yoshida, J. *Macromolecules* **2008**, *41*, 6322–6330. doi:10.1021/ma800769n
63. Fukuyama, T.; Kobayashi, M.; Rahman, M. T.; Kamata, N.; Ryu, I. *Org. Lett.* **2008**, *10*, 533–536. doi:10.1021/ol702718z

64. Nagaki, A.; Tomida, Y.; Miyazaki, A.; Yoshida, J. *Macromolecules* **2009**, *42*, 4384–4387. doi:10.1021/ma900551a
65. Tricotet, T.; O'Shea, D. F. *Chem.–Eur. J.* **2010**, *16*, 6678–6686. doi:10.1002/chem.200903284
66. Greenway, G. M. S.; Haswell, J.; Morgan, D. O.; Skelton, V.; Styring, P. *Sens. Actuators, B* **2000**, *63*, 153–158. doi:10.1016/S0925-4005(00)00352-X
67. Haswell, S. J.; O'Sullivan, B.; Styring, P. *Lab Chip* **2001**, *1*, 164–166. doi:10.1039/b104035a
68. Niwa, S.; Eswaramoorthy, M.; Nair, J.; Raj, A.; Itoh, N.; Shoji, H.; Namba, T.; Mizukami, F. *Science* **2002**, *295*, 105–107. doi:10.1126/science.1066527
69. Jas, G.; Kirschning, A. *Chem.–Eur. J.* **2003**, *9*, 5708–5723. doi:10.1002/chem.200305212
70. Basheer, C.; Hussain, F. S. J.; Lee, H. K.; Valiyaveetil, S. *Tetrahedron Lett.* **2004**, *45*, 7297–7300. doi:10.1016/j.tetlet.2004.08.017
71. Liu, S.; Fukuyama, T.; Sato, M.; Ryu, I. *Org. Process Res. Dev.* **2004**, *8*, 477–481. doi:10.1021/op034200h
72. Kunz, U.; Schönfeld, H.; Solodenko, W.; Jas, G.; Kirschning, A. *Ind. Eng. Chem. Res.* **2005**, *44*, 8458–8467. doi:10.1021/ie048891x
73. Comer, E.; Organ, M. G. *J. Am. Chem. Soc.* **2005**, *127*, 8160–8167. doi:10.1021/ja0512069
74. Lee, C. K. Y.; Holmes, A. B.; Ley, S. V.; McConvey, I. F.; Al-Duri, B.; Leeke, G. A.; Santos, R. C. D.; Sevilged, J. P. K. *Chem. Commun.* **2005**, 2175–2177. doi:10.1039/b418669a
75. Mauger, C.; Buisine, O.; Caravieilhies, S.; Mignani, G. *J. Organomet. Chem.* **2005**, *690*, 3627–3629. doi:10.1016/j.jorganchem.2005.03.071
76. Kirschning, A.; Solodenko, W.; Mennecke, K. *Chem.–Eur. J.* **2006**, *12*, 5972–5990. doi:10.1002/chem.200600236
77. Shore, G.; Morin, S.; Organ, M. G. *Angew. Chem., Int. Ed.* **2006**, *45*, 2761–2766. doi:10.1002/anie.200503600
78. Shi, G.; Hong, F.; Liang, Q.; Fang, H.; Nelson, S.; Weber, S. G. *Anal. Chem.* **2006**, *78*, 1972–1979. doi:10.1021/ac051844+
79. Rahman, M. T.; Fukuyama, T.; Kamata, N.; Sato, M.; Ryu, I. *Chem. Commun.* **2006**, 2236–2238. doi:10.1039/b600970k
80. Murphy, E. R.; Martinelli, J. R.; Zaborenko, N.; Buchwald, S. L.; Jensen, K. F. *Angew. Chem., Int. Ed.* **2007**, *46*, 1734–1737. doi:10.1002/anie.200604175
81. Kawanami, H.; Matsushima, K.; Sato, M.; Ikushima, Y. *Angew. Chem., Int. Ed.* **2007**, *46*, 5129–5132. doi:10.1002/anie.200700611
82. Yamada, Y. M. A.; Watanabe, T.; Torii, K.; Uozumi, Y. *Chem. Commun.* **2009**, 5594–5596. doi:10.1039/b912696a
83. Jin, J.; Cai, M.-M.; Li, J. *Synlett* **2009**, 2534–2538. doi:10.1055/s-0029-1217730
84. Ahmed-Omer, B.; Barrow, D. A.; Wirth, T. *Tetrahedron Lett.* **2009**, *50*, 3352–3355. doi:10.1016/j.tetlet.2009.02.133
85. McMullen, J. P.; Stone, M. T.; Buchwald, S. L.; Jensen, K. F. *Angew. Chem., Int. Ed.* **2010**, *49*, 7076–7080. doi:10.1002/anie.201002590
86. Usutani, H.; Tomida, Y.; Nagaki, A.; Okamoto, H.; Nokami, T.; Yoshida, J. *J. Am. Chem. Soc.* **2007**, *129*, 3046–3047. doi:10.1021/ja068330s
87. Nagaki, A.; Tomida, Y.; Usutani, H.; Kim, H.; Takabayashi, N.; Nokami, T.; Okamoto, H.; Yoshida, J. *Chem.–Asian J.* **2007**, *2*, 1513–1523. doi:10.1002/asia.200700231
88. Nagaki, A.; Takabayashi, N.; Tomida, Y.; Yoshida, J. *Org. Lett.* **2008**, *10*, 3937–3940. doi:10.1021/ol8015572
89. Nagaki, A.; Kim, H.; Yoshida, J. *Angew. Chem., Int. Ed.* **2008**, *47*, 7833–7836. doi:10.1002/anie.200803205
90. Nagaki, A.; Takizawa, E.; Yoshida, J. *J. Am. Chem. Soc.* **2009**, *131*, 1654–1655. doi:10.1021/ja809325a
91. Nagaki, A.; Takabayashi, N.; Tomida, Y.; Yoshida, J. *Beilstein J. Org. Chem.* **2009**, *5*, No. 16. doi:10.3762/bjoc.5.16
92. Tomida, Y.; Nagaki, A.; Yoshida, J. *Org. Lett.* **2009**, *11*, 3614–3617. doi:10.1021/ol901352t
93. Nagaki, A.; Kim, H.; Yoshida, J. *Angew. Chem., Int. Ed.* **2009**, *48*, 8063–8065. doi:10.1002/anie.200904316
94. Nagaki, A.; Takizawa, E.; Yoshida, J. *Chem.–Eur. J.* **2010**, *16*, 14149–14158. doi:10.1002/chem.201000815
95. Nagaki, A.; Kim, H.; Moriwaki, Y.; Matsuo, C.; Yoshida, J. *Chem.–Eur. J.* **2010**, *16*, 11167–11177. doi:10.1002/chem.201000876
96. Nagaki, A.; Kim, H.; Matsuo, C.; Yoshida, J. *Org. Biomol. Chem.* **2010**, *8*, 1212–1217. doi:10.1039/b919325c
97. Tomida, Y.; Nagaki, A.; Yoshida, J. *J. Am. Chem. Soc.* **2011**, *133*, 3744–3747. doi:10.1021/ja110898s
98. Kim, H.; Nagaki, A.; Yoshida, J. *Nat. Commun.* **2011**, *2*, No. 264. doi:10.1038/ncomms1264
99. Suga, S.; Yamada, D.; Yoshida, J. *Chem. Lett.* **2010**, *39*, 404–406. doi:10.1246/cl.2010.404
100. Nagaki, A.; Kenmoku, A.; Moriwaki, Y.; Hayashi, A.; Yoshida, J. *Angew. Chem., Int. Ed.* **2010**, *49*, 7543–7547. doi:10.1002/anie.201002763

License and Terms

This is an Open Access article under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The license is subject to the *Beilstein Journal of Organic Chemistry* terms and conditions: (<http://www.beilstein-journals.org/bjoc>)

The definitive version of this article is the electronic one which can be found at:
[doi:10.3762/bjoc.7.122](https://doi.org/10.3762/bjoc.7.122)