

# Semi-classical Asymptotics for the Partition Function of an Abstract Bose Field Model

Yuta Aihara

Department of Mathematics, Hokkaido University, Sapporo, 060-0810, Japan

Semi-classical asymptotics for the partition function of an abstract Bose field model is considered.

Keywords: semi-classical asymptotics, Bose field, partition function, second quantization, Fock space.

## I. INTRODUCTION

In quantum mechanics, in which a physical constant  $\hbar := h/2\pi$  ( $h$  : the Planck constant) plays an important role, the limit  $\hbar \rightarrow 0$  for various quantities (if it exists) is called the classical limit. Trace formulas in the abstract boson Fock space and the classical limit for the trace  $Z(\beta\hbar)$  (the partition function) of the heat semigroup of a perturbed second quantization operator were derived by Arai [2], where  $\beta > 0$  denotes the inverse temperature. Generally speaking, the classical limit is regarded as the zero-th order approximation in  $\hbar$ . From this point of view, it is interesting to derive higher order asymptotics of various quantities in  $\hbar$ . Such asymptotics are called semi-classical asymptotics. In this paper the asymptotic formula for  $Z(\beta\hbar)$  is stated, which is derived in [1].

## II. A CLASSICAL LIMIT IN THE ABSTRACT BOSON FOCK SPACE

In this section we review a classical limit for the trace of a perturbed second quantization operator and some fundamental facts related to it.

Let  $\mathcal{H}$  be a real separable Hilbert space, and  $A$  be a strictly positive self-adjoint operator acting in  $\mathcal{H}$ . We denote by  $\{\mathcal{H}_s(A)\}_{s \in \mathbb{R}}$  the Hilbert scale associated with  $A$  [3]. For all  $s \in \mathbb{R}$ , the dual space of  $\mathcal{H}_s(A)$  can be naturally identified with  $\mathcal{H}_{-s}(A)$ .

We denote by  $\mathcal{I}_1(\mathcal{H})$  the ideal of the trace class operators on  $\mathcal{H}$ . Let  $\gamma > 0$  be fixed. Throughout this paper, we assume the following.

*Assumption I.*  $A^{9-\gamma} \in \mathcal{I}_1(\mathcal{H})$ .

Under Assumption I, the embedding mapping of  $\mathcal{H}$  into

$$E := \mathcal{H}_{-\gamma}(A)$$

is Hilbert-Schmidt. Hence, by Minlos' theorem, there exists a unique probability measure  $\mu$  on  $(E, \mathcal{B})$  such that the Borel field  $\mathcal{B}$  is generated by  $\{\phi(f) | f \in \mathcal{H}_\gamma(A)\}$  and

$$\int_E e^{i\phi(f)} d\mu(\phi) = e^{-\|f\|_{\mathcal{H}}^2/2}, \quad f \in \mathcal{H}_\gamma(A),$$

where  $\|\cdot\|_{\mathcal{H}}$  denotes the norm of  $\mathcal{H}$ .

The complex Hilbert space  $L^2(E, d\mu)$  is canonically isomorphic to the boson Fock space over  $\mathcal{H}$ , which is called the  $Q$ -space representation of it [3]. We denote by  $d\Gamma(A)$  the second quantization of  $A$  and set

$$H_0 = d\Gamma(A).$$

Then for all  $\beta > 0$ ,  $e^{-\beta H_0} \in \mathcal{I}_1(L^2(E, d\mu))$ .

**DEFINITION 2.1.** *A mapping  $V$  of a Banach space  $X$  into a Banach space  $Y$  is said to be polynomially continuous if there exists a polynomial  $P$  of two real variables with positive coefficients such that*

$$\|V(\phi) - V(\psi)\| \leq P(\|\phi\|, \|\psi\|) \|\phi - \psi\|, \quad \phi, \psi \in X.$$

Let  $V$  be a real valued function on  $E$ . Throughout this paper, we assume the following.

*Assumption II.* *The function  $V$  is bounded from below, 3-times Fréchet differentiable, and  $V, V', V'', V'''$  are polynomially continuous.*

For  $\hbar > 0$ , we define  $V_\hbar$  by

$$V_\hbar(\phi) := V(\sqrt{\hbar} \phi), \quad \phi \in E.$$

and set

$$H_\hbar := H_0 \dot{+} \frac{1}{\hbar} V_\hbar,$$

where  $\dot{+}$  denotes the quadratic form sum.

Under Assumption I, II, for all  $\beta > 0$ ,  $e^{-\beta H_\hbar} \in \mathcal{I}_1(L^2(E, d\mu))$  [ 2 ].

**THEOREM 2.2.** [ 2 ]. *Let  $\beta > 0$ . Then*

$$\lim_{\hbar \rightarrow 0} \frac{\text{Tr } e^{-\beta \hbar H_\hbar}}{\text{Tr } e^{-\beta \hbar H_0}} = \int_E \exp \left( -\beta V \left( \sqrt{\frac{2}{\beta}} A^{-1/2} \phi \right) \right) d\mu(\phi).$$

### III. A CLASS OF LOCALLY CONVEX SPACES

In this section we introduce a class of locally convex spaces, which gives a general framework for the asymptotic analysis discussed in this paper.

We denote by  $\mathbb{R}_+$  the set of the nonnegative real numbers.

DEFINITION 3.1. *A mapping  $f$  from  $\mathbb{R}_+$  to a locally convex space  $X$  is said to be locally bounded if for all  $\delta > 0$  and every continuous seminorm  $p$  on  $X$ ,*

$$p_\delta(f) := \sup_{0 \leq \varepsilon \leq \delta} p(f(\varepsilon)) < \infty.$$

We denote by  $(X^{\mathbb{R}_+})_{\text{l.b.}}$  the linear space of the locally bounded mappings from  $\mathbb{R}_+$  to  $X$ . The topology defined by the seminorms  $\{p_\delta\}_{p,\delta}$  turns  $(X^{\mathbb{R}_+})_{\text{l.b.}}$  into a locally convex space. If  $X$  is a Fréchet space,  $(X^{\mathbb{R}_+})_{\text{l.b.}}$  is a Fréchet space.

Let  $\{E_n\}_{n \in \mathbb{N}}$  be a family of Banach spaces with the property that

$$E_{n+1} \subset E_n, \quad \|\phi\|_n \leq \|\phi\|_{n+1}, \quad \phi \in E_{n+1},$$

for all  $n \in \mathbb{N}$ , where  $\|\cdot\|_n$  denotes the norm of  $E_n$ . Then, the topology defined by the norms  $\{\|\cdot\|_n\}_{n \in \mathbb{N}}$  turns  $\bigcap_{n \in \mathbb{N}} E_n$  into a Fréchet space.

Let  $(X, P)$  be a probability space and  $Y$  be a Banach space. We denote by  $L^p(X, dP; Y)$  the Banach space of the  $Y$ -valued  $L^p$ -functions on  $(X, P)$ . Then  $\bigcap_{p \in \mathbb{N}} L^p(X, dP; Y)$  can be provided with the structure of Fréchet space.

DEFINITION 3.2. *Let  $f$  be a mapping from  $\mathbb{R}_+$  to  $\bigcap_{p \in \mathbb{N}} L^p(X, dP; Y)$ . We say that  $f$  is in  $(\bigcap_{p \in \mathbb{N}} L^p(X, dP; Y))_{\text{u.i.}}^{\mathbb{R}_+}$  if and only if for each  $\delta > 0$ , there exists a nonnegative function  $g \in \bigcap_{p \in \mathbb{N}} L^p(X, dP)$  such that*

$$\sup_{0 \leq \varepsilon \leq \delta} \|f(\varepsilon)(x)\|_Y \leq g(x),$$

*$P$ -a.e.x.*

The set  $(\bigcap_{p \in \mathbb{N}} L^p(X, dP; Y))_{\text{u.i.}}^{\mathbb{R}_+}$  is a linear subspace of  $(\bigcap_{p \in \mathbb{N}} L^p(X, dP; Y))_{\text{l.b.}}^{\mathbb{R}_+}$ . In what follows, we omit  $x$  in  $f(\varepsilon)(x)$ .

Let  $X_1, \dots, X_n$  and  $Z$  be non-empty sets and  $G$  be a real-valued function on  $X_1 \times \dots \times X_n$  and  $F_j$  be a mapping from  $Z$  to  $X_j$ ,  $j = 1, \dots, n$ . We define  $G(F_1, \dots, F_n)$ , the real-valued function on  $Z$ , by

$$G(F_1, \dots, F_n)(z) = G(F_1(z), \dots, F_n(z)), \quad z \in Z.$$

Then we can prove the following propositions.

PROPOSITION 3.3. Let  $Q$  be a polynomial of  $n$  real valuables. Then the mapping  $(F_1, \dots, F_n) \mapsto Q(\|F_1\|, \dots, \|F_n\|)$  from  $\left(\bigcap_{p \in \mathbb{N}} L^p(X, dP; Y)\right)_{\text{u.i.}}^{\mathbb{R}_+}$  to  $\left(\bigcap_{p \in \mathbb{N}} L^p(X, dP)\right)_{\text{u.i.}}^{\mathbb{R}_+}$  is continuous.

PROPOSITION 3.4. Let  $Z_j$  be a Banach space ( $j = 1, \dots, n$ ),  $L$  be a continuous multilinear form on  $Z_1 \times \dots \times Z_n$ , and  $V_j$  be a polynomially continuous mapping from  $Y$  to  $Z_j$  ( $j = 1, \dots, n$ ). Then the mapping  $(F_1, \dots, F_n) \mapsto L(V_1 \circ F_1, \dots, V_n \circ F_n)$  from  $\left(\bigcap_{p \in \mathbb{N}} L^p(X, dP; Y)\right)_{\text{u.i.}}^{\mathbb{R}_+}$  to  $\left(\bigcap_{p \in \mathbb{N}} L^p(X, dP)\right)_{\text{u.i.}}^{\mathbb{R}_+}$  is continuous.

#### IV. AN ASYMPTOTIC FORMULA

Let  $\{\lambda_n\}_{n=1}^\infty$  be the eigenvalues of  $A$ , and  $\{e_n\}_{n=1}^\infty$  be the complete orthonormal system (CONS) of  $\mathcal{H}$  with  $Ae_n = \lambda_n e_n$ , and

$$\sum_{n=1}^\infty \frac{1}{\lambda_n^{\gamma-9}} < \infty \quad (4.1)$$

Let  $\varphi$  be a bijection from  $\mathbb{N} \times \mathbb{N}$  to  $\mathbb{N}$ . For all  $n, m \in \mathbb{N}$ , we set  $f_{n,m} = e_{\varphi(n,m)}$ . Then  $\{f_{n,m}\}_{n,m=1}^\infty$  is a CONS of  $\mathcal{H}$ . For all  $\phi \in E$ , we define

$$\phi_n := \phi(e_n), \quad \phi_{n,m} := \phi(f_{n,m}).$$

Then  $\{\phi_n\}_n$  and  $\{\phi_{n,m}\}_{n,m}$  are families of independent Gaussian random variables such that for all  $n, m, n', m' \in \mathbb{N}$ ,

$$\int_E \phi_n d\mu(\phi) = 0, \quad \int_E \phi_n \phi_m d\mu(\phi) = \delta_{nm} \quad (4.2)$$

$$\int_E \phi_{n,m} \phi_{n',m'} d\mu(\phi) = \delta_{nn'} \delta_{mm'}. \quad (4.3)$$

For all  $m_1, \dots, m_p \in \mathbb{N}$ , we have

$$\sup_{n_1, \dots, n_p \in \mathbb{N}} \int_E |\phi_{n_1}|^{m_1} \dots |\phi_{n_p}|^{m_p} d\mu(\phi) < \infty. \quad (4.4)$$

For all  $N, M \in \mathbb{N}$ , we set

$$\begin{aligned} F_{N,M}(\varepsilon, \omega, s) &= \sqrt{\frac{2}{\beta}} \sum_{n=1}^N \frac{\phi_n}{\sqrt{\lambda_n}} e_n + \sum_{n=1}^N \sum_{m=1}^M \sqrt{\frac{4\varepsilon^2 \lambda_n}{\beta(\varepsilon^2 \lambda_n^2 + (2\pi m)^2)}} (\psi_{n,m} \cos(2\pi m s) \\ &+ \theta_{n,m} \sin(2\pi m s)) e_n, \quad \varepsilon \geq 0, \quad \omega = (\phi, \psi, \theta) \in \Omega, \quad 0 \leq s \leq 1. \end{aligned} \quad (4.5)$$

Then we have

$$\frac{\text{Tre}^{-\beta \hbar H_h}}{\text{Tre}^{-\beta \hbar H_0}} = \lim_{N, M \rightarrow \infty} \int_\Omega \exp \left( -\beta \int_0^1 V(F_{N,M}(\varepsilon, \omega, s)) ds \right) d\nu(\omega), \quad (4.6)$$

where  $\varepsilon = \beta \hbar$  (See [ 2 ], Lemma 5.2, Lemma 5.3. ).

We set

$$Z(\varepsilon) = \lim_{N,M \rightarrow \infty} \int_{\Omega} \exp \left( -\beta \int_0^1 F_{N,M}(\varepsilon, \omega, s) ds \right) d\nu(\omega), \quad \varepsilon \geq 0, \quad (4.7)$$

For all  $n, m \in \mathbb{N}$ , we set

$$\alpha_{n,m}(\varepsilon) = \sqrt{\frac{4\varepsilon^2 \lambda_n}{\beta(\varepsilon^2 \lambda_n^2 + (2\pi m)^2)}}, \quad \varepsilon \geq 0.$$

Then, for all  $\delta > 0$ , there exists a constant  $C > 0$  such that

$$|\alpha_{n,m}(\varepsilon)| \leq \frac{C\sqrt{\lambda_n}}{m}, \quad n, m \in \mathbb{N}, \quad 0 \leq \varepsilon \leq \delta. \quad (4.8)$$

$$|\alpha'_{n,m}(\varepsilon)| \leq \frac{C\sqrt{\lambda_n}}{m}, \quad n, m \in \mathbb{N}, \quad 0 \leq \varepsilon \leq \delta. \quad (4.9)$$

$$|\alpha''_{n,m}(\varepsilon)| \leq \frac{C\lambda_n^{5/2}}{m}, \quad n, m \in \mathbb{N}, \quad 0 \leq \varepsilon \leq \delta. \quad (4.10)$$

$$|\alpha'''_{n,m}(\varepsilon)| \leq \frac{C(\lambda_n^{5/2} + \lambda_n^{9/2})}{m}, \quad n, m \in \mathbb{N}, \quad 0 \leq \varepsilon \leq \delta. \quad (4.11)$$

We denote by  $\mu_{[0,1]}^{(L)}$  the Lebesgue measure on  $[0, 1]$ . Then by (4.8),(4.9),(4.10) and (4.11), we can prove the following lemma.

LEMMA 4.1.  $\{F_{N,M}\}_{N,M \in \mathbb{N}}, \{F'_{N,M}\}_{N,M \in \mathbb{N}}, \{F''_{N,M}\}_{N,M \in \mathbb{N}}, \{F'''_{N,M}\}_{N,M \in \mathbb{N}}$  are Cauchy nets in  $(\bigcap_{p \in \mathbb{N}} L^p(\Omega \times [0, 1], d(\nu \otimes \mu_{[0,1]}^{(L)}; E))_{\text{u.i.}}^{\mathbb{R}^+}$ .

For all  $N, M \in \mathbb{N}$ , we set

$$G_{N,M}(\varepsilon, \omega) = \exp \left( -\beta \int_0^1 V(F_{N,M}(\varepsilon, \omega, s)) ds \right) \quad \varepsilon \geq 0, \quad \omega \in \Omega.$$

Then by Proposition 3.4 and Lemma 4.1, we can prove the following lemma.

LEMMA 4.2.  $\{G_{N,M}\}_{N,M \in \mathbb{N}}, \{G'_{N,M}\}_{N,M \in \mathbb{N}}, \{G''_{N,M}\}_{N,M \in \mathbb{N}}, \{G'''_{N,M}\}_{N,M \in \mathbb{N}}$  are Cauchy nets in  $(\bigcap_{p \in \mathbb{N}} L^p(\Omega, d\nu))_{\text{u.i.}}^{\mathbb{R}^+}$ .

By Lemma 4.2 and the fact that  $\alpha_{n,m}$  is infinitely differentiable for all  $n, m \in \mathbb{N}$ ,  $\int_{\Omega} H_{N,M}(\varepsilon, \omega) d\nu(\omega)$  with  $H_{N,M} = G_{N,M}, G'_{N,M}, G''_{N,M}, G'''_{N,M}$  uniformly converges in  $\varepsilon$ . Hence one can interchange the limit  $\lim_{N,M \rightarrow \infty}$  with differentiations in  $\varepsilon$  and see that  $Z$  is 3-times continuously differentiable in  $\mathbb{R}_+$ .

We can prove the following theorem.

THEOREM 4.3. For all  $\beta > 0$ ,

$$\begin{aligned}
& \frac{\text{Tre}^{-\beta\hbar H_\hbar}}{\text{Tre}^{-\beta\hbar H_0}} \\
&= \int_E \exp\left(-\beta V\left(\sqrt{\frac{2}{\beta}}A^{-1/2}\phi\right)\right) d\mu(\phi) \\
&\quad - \frac{\beta^3\hbar^2}{2} \sum_{m=1}^{\infty} \int_{E^2} d\mu(\phi)d\mu(\psi) \exp\left(-\beta V\left(\sqrt{\frac{2}{\beta}}A^{-1/2}\phi\right)\right) \\
&\times V''\left(\sqrt{\frac{2}{\beta}}A^{-1/2}\phi\right) \left(A^{1/2}\left(\frac{1}{\sqrt{\beta\pi m}} \sum_{n=1}^{\infty} \psi_{n,m}e_n\right), A^{1/2}\left(\frac{1}{\sqrt{\beta\pi m}} \sum_{n=1}^{\infty} \psi_{n,m}e_n\right)\right) \\
&\quad + o(\hbar^2)
\end{aligned}$$

as  $\hbar \rightarrow 0$ .

#### REFERENCES

- [1] Y. Aihara, Semi-classical asymptotics in an abstract Bose field model, *IJPAM* **85** (2013), 265-284.
- [2] A. Arai, Trace formulas, a Golden-Thompson inequality and classical limit in Boson Fock space, *J. Funct. Anal.* **136** (1996), 510-546.
- [3] A. Arai, Functional Integral Methods in Quantum Mathematical Physics, Kyoritsu-shuppan (2010) (in Japanese).