

Radial symmetry of self-similar solutions for semilinear heat equations

神戸大学工学部 内藤 雄基 (Yūki Naito)
大阪大学理学研究科 鈴木 貴 (Takashi Suzuki)

We consider the symmetry properties of positive solutions of the equation

$$\Delta u + \frac{1}{2}x \cdot \nabla u + \frac{1}{p-1}u + u^p = 0 \quad \text{in } \mathbf{R}^n, \quad (1.1)$$

where $n \geq 2$ and $p > 1$. This equation arises in the study of (forward) self-similar solutions of the semilinear heat equation

$$w_t = \Delta w + w^p \quad \text{in } \mathbf{R}^n \times (0, \infty). \quad (1.2)$$

It is well known that if $w(x, t)$ satisfies (1.2), then, for $\mu > 0$ the rescaled functions

$$w_\mu(x, t) = \mu^{2/(p-1)}w(\mu x, \mu^2 t)$$

define a one parameter family of solutions to (1.2). A solution w is said to be self-similar, when $w_\mu(x, t) = w(x, t)$ for all $\mu > 0$. It can be easily checked that w is a self-similar solution to (1.2) if and only if w has the form

$$w(x, t) = t^{-1/(p-1)}u(x/\sqrt{t}), \quad (1.3)$$

where u satisfies the elliptic Eq. (1.1). Moreover, if u has spherical symmetry, that is if $u = u(r)$, $r = |x|$, then u satisfies the ordinary differential equation

$$u'' + \left(\frac{n-1}{r} + \frac{r}{2}\right)u' + \frac{1}{p-1}u + u^p = 0, \quad r > 0. \quad (1.4)$$

Such self-similar solutions are often used to describe the large time behavior of global solutions to the Cauchy problem, see, e.g., [11, 13, 3, 14, 5, and 15], and to show nonuniqueness of solution to (1.2) with zero initial data in a certain functional space, see [12].

First we state the result concerning the symmetry properties of the solution of (1.1).

THEOREM 1.1. *Let $u \in C^2(\mathbf{R}^n)$ be a positive solution of (1.1) such that*

$$u(x) = o(|x|^{-2/(p-1)}) \quad \text{as } |x| \rightarrow \infty. \quad (1.5)$$

Then u must be radially symmetric about the origin.

The proof of Theorem 1.1 is based on the moving planes argument. This technique was developed by Serrin [18] in PDE theory, and extended and generalized by Gidas, Ni, and Nirenberg [9, 10]. We remark that with a change of variables we are still able to prove a radial symmetry result for Eq. (1.1).

Let us consider the problem

$$\begin{cases} u'' + \left(\frac{n-1}{r} + \frac{r}{2} \right) u' + \frac{1}{p-1} u + |u|^{p-1} u = 0, & r > 0, \\ u'(0) = 0 \quad \text{and} \quad u(0) = \alpha \in \mathbf{R}. \end{cases} \quad (1.6)$$

The problem (1.6) has been investigated extensively in [12, 16, 20, and 2]. We denote by $u(r; \alpha)$ the unique solution of (1.6). We recall that $u(r; \alpha)$ has the following properties:

- (i) $\lim_{r \rightarrow \infty} r^{2/(p-1)} u(r; \alpha) = L(\alpha)$ exists and is finite for every $\alpha \in \mathbf{R}$ (see [12, Theorem 5]);
- (ii) if $L(\alpha) = 0$, then there exists a constant $A \neq 0$ such that

$$u(r; \alpha) = A e^{-r^2/4} r^{2/(p-1)-n} \{1 + O(r^{-2})\} \quad \text{as } r \rightarrow \infty$$

(see [16, Theorem 1]);

- (iii) if $p \geq (n+2)/(n-2)$, then $u(r; \alpha)$ is positive on $[0, \infty)$ and $L(\alpha) > 0$ for every $\alpha > 0$ (see [12, Theorem 5]);

- (iv) if $(n+2)/n < p < (n+2)/(n-2)$, then there exists a unique $\alpha > 0$ such that $u(r; \alpha)$ is positive on $[0, \infty)$ and $L(\alpha) = 0$ (see [20, Theorem 1] and [2, Theorem 1.2 and Corollary 1.3]).

By virtue of Theorem 1.1 we obtain the following:

COROLLARY 1.1. (i) *Assume that $p \geq (n+2)/(n-2)$. Then there exists no positive solution u of (1.1) satisfying (1.5).*

(ii) *Assume that $(n+2)/n < p < (n+2)/(n-2)$. Then there exists a unique positive solution $u(x)$ satisfying (1.5). Moreover, the solution u is radially symmetric about the origin.*

Remark. The result (i) is differently proven by [3, Proposition 4.3] based on the Pohozaev identity.

Following the notations in [3] and [14], we define

$$L^2(K) = \{u : \mathbf{R}^n \rightarrow \mathbf{R}; \int_{\mathbf{R}^n} |u|^2 K(x) dx < \infty\} \quad \text{and}$$

$$H^1(K) = \{u : \mathbf{R}^n \rightarrow \mathbf{R}; \int_{\mathbf{R}^n} (|u|^2 + |\nabla u|^2) K(x) dx < \infty\},$$

where $K(x) = \exp(-|x|^2/4)$. Escobedo and Kavian have shown in [3, Proposition 3.5] that if $1 < p < (n+2)/(n-2)$ and if $u \in H^1(K)$ is a solution of (1.1), then $u \in C^2(\mathbf{R}^n)$ and satisfies $u(x) = O(\exp(-|x|^2/8))$ as $|x| \rightarrow \infty$. As a consequence of Corollary 1.1, we obtain the following:

COROLLARY 1.2. *Assume that $(n+2)/n < p < (n+2)/(n-2)$. Then the problem*

$$\begin{cases} \Delta u + \frac{1}{2}x \cdot \nabla u + \frac{1}{p-1}u + u^p = 0 & \text{in } \mathbf{R}^n, \\ u \in H^1(K) \quad \text{and} \quad u > 0 & \text{in } \mathbf{R}^n, \end{cases} \quad (1.7)$$

has a unique solution.

Let us consider the Cauchy problem

$$\begin{cases} w_t = \Delta w + w^p & \text{in } \mathbf{R}^n \times (0, \infty), \\ w(x, 0) = \tau w_0 & \text{in } \mathbf{R}^n, \end{cases} \quad (1.8)$$

where $w_0 \in L^2(K) \cap L^\infty(\mathbf{R}^n)$, $w_0 \geq 0$, and $\tau > 0$ is a parameter. We denote by $w(x, t; \tau)$ the unique solutions of (1.8) (see [15]). Combining the result by Kawanago [15, Theorem 1] and Corollary 1.2, we obtain the following, where the asymptotic behavior of $w(\cdot, t; \tau)$ as $t \rightarrow \infty$ becomes clearer.

COROLLARY 1.3. *Assume that $(n+2)/n < p < (n+2)/(n-2)$. Then there exists a unique $\tau_0 > 0$ such that the solution $w(x, t; \tau)$ is a global solution if $\tau \in (0, \tau_0]$, and $w(x, t; \tau)$ blows up in finite time if $\tau \in (\tau_0, \infty)$. Moreover, $w(x, t; \tau_0)$ satisfies*

$$\lim_{t \rightarrow \infty} \left\| t^{1/(p-1)} w(\cdot, t; \tau_0) - u_0(\cdot/\sqrt{t}) \right\|_{L^\infty(\mathbf{R}^n)} = 0,$$

where u_0 is a unique solution of the problem (1.7).

Next we consider the existence of nonradial solutions of (1.1). Let $p > (n+2)/n$ and let $U(r)$ be a positive solution of (1.4) satisfying

$$U'(0) = 0 \quad \text{and} \quad \lim_{r \rightarrow \infty} r^{2/(p-1)} U(r) > 0. \quad (1.9)$$

The existence of such U is obtained by [12, Theorem 5]. Define $\ell = \ell(U) > 0$ as

$$\ell = \lim_{r \rightarrow \infty} r^{2/(p-1)} U(r). \quad (1.10)$$

We investigate the Cauchy problem for Eq. (1.2) with

$$w(x, 0) = w_0 \in L^1_{\text{loc}}(\mathbf{R}^n), \quad (1.11)$$

where

$$0 \leq w_0(x) \leq \ell |x|^{-2/(p-1)}, \quad w_0 \not\equiv 0, \quad x \in \mathbf{R}^n \setminus \{0\}. \quad (1.12)$$

Relation (1.11) is taken in the sense of $L^1_{\text{loc}}(\mathbf{R}^n)$, that is,

$$\int_K |w(x, t) - w_0(x)| dx \rightarrow 0 \quad \text{as } t \rightarrow 0$$

for any compact subset K of $\mathbf{R}^n$. We note that $w_0 \in L^1_{\text{loc}}(\mathbf{R}^n)$ if (1.12) holds with $p > (n+2)/n$.

THEOREM 1.2. *Let $p > (n+2)/n$. Assume that (1.12) holds, where ℓ is the constant in (1.10). Then there exists a positive solution $w \in C^{2,1}(\mathbf{R}^n \times (0, \infty))$ of (1.2) and (1.11). Assume, furthermore, that $w_0 \in C(\mathbf{R}^n \setminus \{0\})$, then w satisfies*

$$w(x, t) \rightarrow w_0(x) \quad \text{as } t \rightarrow 0 \quad \text{uniformly in } |x| \geq r \text{ for every } r > 0. \quad (1.13)$$

Moreover, w is self-similar if $\mu^{2/(p-1)} w_0(\mu x) = w_0(x)$ for every $\mu > 0$.

COROLLARY 1.4. *Let $p > (n+2)/n$. Assume that $A : S^{n-1} \rightarrow \mathbf{R}$ is continuous and satisfies*

$$0 \leq A(\sigma) \leq \ell, \quad A \not\equiv 0, \quad \sigma \in S^{n-1}. \quad (1.14)$$

Then there exists a positive self-similar solution $w \in C^{2,1}(\mathbf{R}^n \times (0, \infty))$ of (1.2) satisfying (1.11) and (1.13) with $w_0(x) = A(x/|x|)|x|^{-2/(p-1)}$.

Recall that self-similar solutions w to (1.2) have the form (1.3) with u satisfying (1.1). Therefore, $w(\sigma, t) = r^{2/(p-1)} u(r\sigma)$ for $\sigma \in S^{n-1}$, where $r = 1/\sqrt{t}$. Then we obtain the following corollary, which shows that condition (1.5) in Theorem 1.1 is crucial.

COROLLARY 1.5. Let $p > (n + 2)/n$. Assume that $A : S^{n-1} \rightarrow \mathbf{R}$ is continuous and satisfies (1.14). Then there exists a positive non-radial solution u of (1.1) satisfying

$$r^{2/(p-1)}u(r\sigma) \rightarrow A(\sigma) \quad \text{as } r \rightarrow \infty \quad \text{uniformly in } \sigma \in S^{n-1}.$$

Remark. (i) If $1 < p \leq (n + 2)/n$, no time global, non-negative, and nontrivial solution exists in (1.2) (see, e.g., [7], [19]). Therefore, (1.1) admits a positive solution only if $p > (n + 2)/n$.

(ii) We find that the solution w of (1.2) and (1.11) obtained in Theorem 1.2 is a minimal solution of the integral equation

$$w(x, t) = \int_{R^n} \Gamma(x - y : t) w_0(y) dy + \int_0^t \int_{R^n} \Gamma(x - y : t - s) [w(y, s)]^p dy ds,$$

where $\Gamma(x : t) = (4\pi t)^{-n/2} e^{-|x|^2/4t}$. See the proof of Theorem 1.2 below.

(iii) Galaktionov and Vazquez [8] studied the Cauchy problem (1.2) and (1.11) with singular initial values for the case $p > n/(n - 2)$.

REFERENCES

- [1] E. DiBenedetto, "An Introduction to Partial Differential Equations", Birkhäuser, Boston, 1995.
- [2] C. Dohmen and M. Hirose, Structure of positive radial solutions to the Haraux-Weissler equation, *Nonlinear Anal. TMA* **33** (1998), 51-69.
- [3] M. Escobedo and O. Kavian, Variational problems related to self-similar solutions for the heat equation, *Nonlinear Anal. TMA* **11** (1987), 1103-1133.
- [4] M. Escobedo and O. Kavian, Asymptotic behavior of positive solutions of a nonlinear heat equation, *Houston J. Math.* **13** (1987), 39-50.
- [5] M. Escobedo, O. Kavian, and H. Matano, Large time behavior of solutions of a dissipative semilinear heat equation, *Comm. Partial Differential Equations* **27** (1995), 1427-1452.
- [6] A. Friedman, "Partial Differential Equations of Parabolic Type", Prentice-Hall, New Jersey, 1964.
- [7] H. Fujita, On the blowing up of solutions of the Cauchy problem for $u_t = \Delta u + u^{1+\alpha}$, *J. Fac. Sci. Univ. Tokyo, Sect. I* **13** (1966), 109-124.
- [8] V. A. Galaktionov and J. L. Vazquez, Continuation of blowup solutions of nonlinear heat equations in several space dimensions, *Comm. Pure Appl. Math.* **50** (1997), 1-67.
- [9] B. Gidas, W.-M. Ni, and L. Nirenberg, Symmetry and related properties via the maximum principle, *Comm. Math. Phys.* **68** (1979), 209-243.

- [10] B. Gidas, W.-M. Ni, and L. Nirenberg, Symmetry of positive solutions of nonlinear elliptic equations in $\mathbf{R}^n$ in "Mathematical Analysis and Applications, Part A", ed. by L. Nachbin, Adv. Math. Suppl. Stud. **7**, Academic Press, New York, 1981, 369-402.
- [11] A. Gmira and L. Veron, Large time behaviour of the solutions of a semilinear parabolic equation in $\mathbf{R}^N$, *J. Differential Equations* **53** (1984), 258-276.
- [12] A. Haraux and F. B. Weissler, Non-uniqueness for a semilinear initial value problem, *Indiana Univ. Math. J.* **31** (1982), 167-189.
- [13] S. Kamin and L. A. Peletier, Large time behaviour of solutions of the heat equation with absorption, *Ann. Scuola Norm. Sup. Pisa* **12** (1985), 393-408.
- [14] O. Kavian, Remarks on the large time behavior of a nonlinear diffusion equation, *Annal. Institut Henri Poincaré-Analyse nonlinéaire* **4** (1987), 423-452.
- [15] T. Kawanago, Asymptotic behavior of solutions of a semilinear heat equation with subcritical nonlinearity, *Annal. Institut Henri Poincaré-Analyse nonlinéaire* **13** (1996), 1-15.
- [16] L. A. Peletier, D. Terman, and F. B. Weissler, On the equation $\Delta u + \frac{1}{2}x \cdot \nabla u + f(u) = 0$, *Arch. Rational Mech. Anal.* **94** (1986), 83-99.
- [17] M. Protter and H. Weinberger, "Maximal Principles in Differential Equations", Prentice-Hall, Englewood Cliffs, N.J. 1967.
- [18] J. Serrin, A symmetry problem in potential theory, *Arch. Rational Mech. Anal.* **43** (1971), 304-318.
- [19] F. B. Weissler, Existence and non-existence of global solutions for a semilinear heat equation, *Israel J. Math.* **38** (1981), 29-40.
- [20] E. Yanagida, Uniqueness of rapidly decaying solutions to the Haraux-Weissler equation, *J. Differential Equations* **127** (1996), 561-570.