

SYSTEMS OF NONLINEAR VARIATIONAL INEQUALITIES ARISING FROM PHASE TRANSITION PHENOMENA

N. KENMOCHI and M. NIEZGODKA

1. Introduction

We consider an evolution system, consisting of a nonlinear second-order parabolic PDE and a nonlinear fourth-order parabolic PDE with constraint, which is described as follows:

$$\rho(u)_t + \lambda(w)_t - \Delta u = h(t, x) \quad \text{in } Q := (0, T) \times \Omega, \quad (1.1-1)$$

$$\frac{\partial u}{\partial n} + n_o u = h_o(t, x) \quad \text{on } \Sigma := (0, T) \times \Gamma, \quad (1.1-2)$$

$$u(0, \cdot) = u_o \quad \text{in } \Omega, \quad (1.1-3)$$

$$w_t - \Delta(-\nu \Delta w + \xi + g(w) - \lambda_o(w)u) = 0 \quad \text{in } Q, \quad (1.2-1)$$

$$\frac{\partial w}{\partial n} = 0, \quad \frac{\partial}{\partial n}(-\nu \Delta w + \xi + g(w) - \lambda_o(w)u) = 0 \quad \text{on } \Sigma, \quad (1.2-2)$$

$$\xi \in \beta(w) \quad \text{on } Q, \quad (1.2-3)$$

$$w(0, \cdot) = w_o \quad \text{in } \Omega. \quad (1.2-4)$$

Here Ω is a bounded domain in $\mathbf{R}^N$ ($1 \leq N \leq 3$) with smooth boundary $\Gamma = \partial\Omega$; $\rho : \mathbf{R} \rightarrow \mathbf{R}$, $g : \mathbf{R} \rightarrow \mathbf{R}$ and $\lambda : \mathbf{R} \rightarrow \mathbf{R}$ are given functions and $\lambda_o(r) = \lambda'(r)$ (= the derivative of λ) for $r \in \mathbf{R}$; $\nu > 0$ and $n_o \geq 0$ are given constants, and h and h_o are given functions on Q and Σ , respectively; u_o and w_o are initial data; β is a given maximal monotone graph in $\mathbf{R} \times \mathbf{R}$.

The system (1.1)-(1.2) is interpreted as a simplified model for thermodynamical phase separation in which w represents the order parameter, $\theta = -\frac{1}{u}$ the (Kelvin) temperature and the free energy functional $F(\theta, w)$ is supposed to be dependent upon the temperature θ and to be given by the formula

$$F(\theta, w) := \int_{\Omega} f(\theta, w, \nabla w) dx, \quad w \in H^1(\Omega), \quad (1.3)$$

$$f(\theta, w, \nabla w) = \left\{ \frac{1}{2}(\nu_o + \nu_1 \theta) |\nabla w|^2 + \tau(\theta) + \theta(\hat{\beta}(w) + \hat{g}(w)) + \lambda(w) \right\},$$

where $\hat{\beta}$ is a proper l.s.c. convex function such that $\partial \hat{\beta} = \beta$ in $\mathbf{R} \times \mathbf{R}$, $\hat{g}$ is a primitive of g on $\mathbf{R}$, λ is the same as above, $\nu_o \geq 0$, $\nu_1 > 0$ are constants and $\tau : \mathbf{R} \rightarrow \mathbf{R}$ is a smooth function.

In some general settings, various models for thermodynamical phase separation phenomena have been proposed and studied for instance by Luckhaus-Visintin [11] and Alt-Pawlow [1,2]. However, in their models the constraint (1.2-3) is not taken account of.

To illustrate our system (1.1)-(1.2), for instance, consider a binary system of alloys with components A and B occupying Ω ; let $w := w_A$ and w_B be the local concentrations of A and B , respectively, such that

$$w_A + w_B = \text{const.};$$

suppose that the free energy functional $F(\theta, w)$ of the Ginzburg-Landau type is of the form (1.3). Then, according to the thermodynamics approach of DeGroot-Mazur [5] and Alt-Pawlow [1,2], we can derive from (1.3) with transformation $u := -1/\theta$, the mass and energy balance equations:

$$\rho(u)_t + \lambda(w)_t + [\frac{1}{2}\nu_o|\nabla w|^2]_t + \nabla \cdot \mathbf{q} = h(t, x) \quad \text{in } Q, \quad (1.4)$$

$$w_t + \nabla \cdot \mathbf{j} = 0 \quad \text{in } Q, \quad (1.5)$$

where $\rho(u) = \tau(\theta) - \theta\tau'(\theta)$, $\mathbf{q}$ is the energy flux due to heat and mass transfer, $\mathbf{j}$ is the mass flux of the component A and h is a given heat source. Now suppose further that the fluxes $\mathbf{q}$ and $\mathbf{j}$ are described by the following constitutive relations:

$$\mathbf{q} = \nabla\left(\frac{1}{\theta}\right) (= -\nabla u) \quad \text{in } Q, \quad (1.6)$$

$$\mathbf{j} = -\nabla\left(\frac{\mu}{\theta}\right) (= \nabla(u\mu)) \quad \text{in } Q, \quad (1.7)$$

where

$$\frac{\mu}{\theta} = \frac{\delta}{\delta w} \left[\int_{\Omega} \frac{f(\theta, w, \nabla w)}{\theta} dx \right] \quad (1.8)$$

and $\frac{\delta}{\delta w}$ denotes the functional derivative with respect to w . Since $f(\theta, w, \nabla w)$ includes the non-smooth term $\hat{\beta}(w)$, the right hand side of (1.8) is here understood in the multivalued sense

$$\begin{aligned} & \frac{\delta}{\delta w} \left[\int_{\Omega} \frac{f(\theta, w, \nabla w)}{\theta} dx \right] \\ &= \left\{ -\nabla \cdot \left(\frac{\nu_o}{\theta} + \nu_1 \right) \nabla w + \xi + g(w) + \frac{\lambda'(w)}{\theta}; \xi \in L^2(\Omega), \xi \in \beta(w) \text{ a.e. on } \Omega \right\}, \end{aligned} \quad (1.9)$$

Now, combine (1.4)-(1.5) with (1.6)-(1.9). Then we obtain

$$\rho(u)_t + \lambda(w)_t + [\frac{\nu_o}{2}|\nabla w|^2]_t - \Delta u = h \quad \text{in } Q, \quad (1.10)$$

and

$$w_t - \Delta(-\nabla \cdot (\nu_1 - \nu_o u) \nabla w + \xi + g(w) - \lambda'(w)u) = 0 \quad \text{in } Q, \quad (1.11)$$

$$\xi \in \beta(w) \quad \text{in } Q. \quad (1.12)$$

Therefore, if $\nu_o = 0$ and $\nu_1 = \nu$, or if in (1.10) the term $[\frac{\nu_o}{2}|\nabla w|^2]_t$ is experimentally allowed to be neglected and in (1.11) the coefficient $(\nu_1 - \nu_o u)$ of ∇w replaced by a positive constant ν , then system (1.1-1)-(1.2- i), $i = 1, 3$, is regarded as a simplified form of (1.10)-(1.12). System (1.1)-(1.2) consists of these equations and initial-boundary conditions (1.1- i), $i = 2, 3$, and (1.2- i), $i=2,4$.

The aim of this paper is to study a weak formulation for system (1.1)-(1.2) in the variational sense, taking advantage of subdifferential techniques in Hilbert spaces.

2. Main results

Throughout this note, for a general (real) Banach space X we denote by $|\cdot|_X$ the norm in X and by X^* the dual space of X .

For simplicity we use the notations:

$$\begin{aligned} (v, w) &:= \int_{\Omega} v w dx && \text{for } v, w \in L^2(\Omega), \\ (v, w)_{\Gamma} &:= \int_{\Gamma} v w d\Gamma(x) && \text{for } v, w \in L^2(\Gamma), \\ a(v, w) &:= \int_{\Omega} \nabla v \cdot \nabla w dx && \text{for } v, w \in H^1(\Omega). \end{aligned}$$

Moreover we put

$$\begin{aligned} H &:= L^2(\Omega), && V := H^1(\Omega), \\ H_o &:= \{z \in H; \int_{\Omega} z dx = 0\}, && V_o := V \cap H_o, \end{aligned}$$

and denote by π the projection from H onto H_o , i.e.

$$\pi(z)(x) := z(x) - \frac{1}{|\Omega|} \int_{\Omega} z(y) dy, \quad z \in H.$$

Also, H_o is a Hilbert space with $|z|_{H_o} = |z|_H$ as well as V_o with $|z|_{V_o} = |\nabla z|_H$; we use sometimes symbol $(\cdot, \cdot)_o$ for the inner product in H_o and $\langle \cdot, \cdot \rangle_o$ for the duality pairing between V_o^* and V_o .

As usual, identifying H with its dual, we have

$$V \subset H \subset V^*$$

with dense and compact embeddings. Similarly, identifying H_o with its dual, we have

$$V_o \subset H_o \subset V_o^*$$

with dense and compact embeddings. Also, we denote by J_o the duality mapping from V_o onto V_o^* which is defined by the formula

$$\langle J_o z, \eta \rangle_o = a(z, \eta) \quad \text{for all } z, \eta \in V_o.$$

Therefore, in particular, if $z^* := J_o z \in H_o$, then $z \in H^2(\Omega)$ and z is the unique solution of the Neumann problem

$$-\Delta z = z^* \quad \text{in } \Omega, \quad \frac{\partial z}{\partial n} = 0 \quad \text{on } \Gamma, \quad \int_{\Omega} z dx = 0. \quad (2.1)$$

Accordingly, if $\eta \in H^2(\Omega)$ and $\frac{\partial \eta}{\partial n} = 0$ a.e. on Γ , then $J_o[\pi(\eta)] = -\Delta \eta$.

Now, we denote by (P) the system (1.1)-(1.2) mentioned in section 1 and discuss it under the following assumptions (A1)-(A6):

- (A1) $\rho : \mathbf{R} \rightarrow \mathbf{R}$ is an increasing Lipschitz continuous function with Lipschitz continuous inverse $\rho^{-1} : \mathbf{R} \rightarrow \mathbf{R}$; we denote by C_ρ a common Lipschitz constant of ρ and ρ^{-1} .
- (A2) $\lambda, \lambda_o : \mathbf{R} \rightarrow \mathbf{R}$ are Lipschitz continuous functions and $\lambda_o = \lambda'$; we denote by C_λ a common Lipschitz constant of λ and λ_o .
- (A3) $g : \mathbf{R} \rightarrow \mathbf{R}$ is a Lipschitz continuous function; we denote by C_g the Lipschitz constant of g .
- (A4) ν is a positive constant and n_o is a non-negative constant.
- (A5) β is a maximal monotone graph in $\mathbf{R} \times \mathbf{R}$ with bounded and non-empty interior $\text{int}.D(\beta)$ of the domain $D(\beta)$ in $\mathbf{R}$; we put $\text{int}.D(\beta) = (\sigma_*, \sigma^*)$ for $-\infty < \sigma_* < \sigma^* < \infty$ and hence $\overline{D(\beta)} = [\sigma_*, \sigma^*]$, and we may assume that β is the subdifferential of a non-negative, proper, l.s.c. and convex function $\hat{\beta}$ on $\mathbf{R}$, since the range $R(\beta)$ of β is the whole $\mathbf{R}$.
- (A6) $0 < T < \infty$, $h \in L^2(0, T; H)$, $h_o \in W^{1,2}(0, T; L^2(\Gamma))$ and $u_o \in H$, $w_o \in V$ with $\hat{\beta}(w_o) \in L^1(\Omega)$.

We introduce

$$K(\hat{\beta}) := \{z \in H; \hat{\beta}(z) \in L^1(\Omega)\}$$

and

$$K_m(\hat{\beta}) := \{z \in K(\hat{\beta}); \frac{1}{|\Omega|} \int_{\Omega} z dx = m\} \quad \text{for each } m \in \mathbf{R}.$$

We next give the weak formulation for (P).

Definition 2.1. A couple $\{u, w\}$ of functions $u : [0, T] \rightarrow V$ and $w : [0, T] \rightarrow H^2(\Omega)$ is called a (weak) solution of (P), if the following conditions (w1)-(w4) are satisfied:

- (w1) $u \in L^2(0, T; V) \cap L^\infty(0, T; H)$, $\rho(u) \in C_w([0, T]; H)$, $C_w([0, T]; H)$ being the space of all weakly continuous functions from $[0, T]$ into H , $\rho(u)' (= \frac{d}{dt}\rho(u)) \in L^1(0, T; V^*)$, $w \in L^2(0, T; H^2(\Omega)) \cap L^\infty(0, T; V)$, $w' \in L^2(0, T; V^*)$ and $\lambda(w)' \in L^1(0, T; V^*)$;
- (w2) $\rho(u)(0) = \rho(u_o)$ and $w(0) = w_o$;
- (w3) for a.e. $t \in [0, T]$ and all $z \in V$,

$$\frac{d}{dt}(\rho(u(t)) + \lambda(w(t)), z) + a(u(t), z) + (n_o u(t) - h_o(t), z)_\Gamma = (h(t), z); \quad (2.2)$$

- (w4) for a.e. $t \in [0, T]$,

$$\frac{\partial}{\partial n} w(t) = 0 \quad \text{a.e. on } \Gamma, \quad (2.3)$$

and there is a function $\xi \in L^2(0, T; H)$ such that

$$\xi \in \beta(w) \quad \text{a.e. in } Q \quad (2.4)$$

and

$$\frac{d}{dt}(w(t), \eta) + \nu(\Delta w(t), \Delta \eta) - (g(w(t)) + \xi(t) - \lambda'(w(t))u(t), \Delta \eta) = 0 \quad (2.5)$$

for all $\eta \in H^2(\Omega)$ with $\frac{\partial \eta}{\partial n}$ a.e. on Γ , and a.e. $t \in [0, T]$.

When it is necessary to indicate the data h, h_o, u_o, w_o , we denote problem (P) by $(P; h, h_o, u_o, w_o)$.

Remark 2.1. Let $\{u, w\}$ be any solution of (P). Then it follows from (2.5) in (w4) that

$$\frac{d}{dt}(w(t), 1) = 0 \quad \text{for a.e. } t \in [0, T],$$

whence

$$\int_{\Omega} w(t, x) dx = \int_{\Omega} w_o dx \quad \text{for all } t \in [0, T].$$

Therefore, putting

$$m := \frac{1}{|\Omega|} \int_{\Omega} w_o dx, \quad (2.6)$$

we observe that $w(t) - m \in V_o$ for all $t \in [0, T]$.

Our main results of this paper are stated as follows:

Theorem 2.1. Assume that $1 \leq N \leq 3$ and (A1)-(A6) hold, and assume with notation (2.6) that

$$m \in \text{int}.D(\beta), \quad \text{i.e. } \sigma_* < m < \sigma^*.$$

Then (P) has one and only one solution $\{u, w\}$. Moreover, the solution $\{u, w\}$ has the following bounds:

$$\begin{aligned} & |u|_{L^\infty(0, T; H)} + |u|_{L^2(0, T; V)} + |w|_{L^\infty(0, T; V)} + |\hat{\beta}(w)|_{L^\infty(0, T; L^1(\Omega))} + |w'|_{L^2(0, T; V^*)} \\ & \leq R_o(|u_o|_H, |w_o|_V, |\hat{\beta}(w_o)|_{L^1(\Omega)}, |h|_{L^2(0, T; H)}, |h_o|_{L^2(0, T; L^2(\Gamma))}), \end{aligned} \quad (2.7)$$

where $R_o : \mathbf{R}_+^5 \rightarrow \mathbf{R}$ is a function which is bounded on each bounded subset of $\mathbf{R}_+^5$;

$$\begin{aligned} & |w|_{L^2(0, T; H^2(\Omega))} + |\rho(u)'|_{L^1(0, T; V^*)} + |\lambda(w)'|_{L^1(0, T; V^*)} \\ & \leq R_1\left(\frac{1}{\delta}, r(\delta), |u_o|_H, |w_o|_V, |\hat{\beta}(w_o)|_{L^1(\Omega)}, |h|_{L^2(0, T; H)}, |h_o|_{L^2(0, T; L^2(\Gamma))}\right), \end{aligned} \quad (2.8)$$

where $R_1 : \mathbf{R}_+^7 \rightarrow \mathbf{R}_+$ is a function which is bounded on each bounded subset of $\mathbf{R}_+^7$, δ is an arbitrary number satisfying

$$0 < \delta < 1, \quad \sigma_* < m - \delta < m + \delta < \sigma^*, \quad (2.9)$$

and

$$r(\delta) = \sup\{|r'|; r' \in \beta(m - \delta) \cap \beta(m + \delta)\}. \quad (2.10)$$

Remark 2.2. In estimates (2.7) and (2.8), the dependence of the solution $\{u, w\}$ upon functions ρ , λ , g and β is not explicitly indicated. However, as will be able to be easily checked, the functions R_o and R_1 are chosen so as to be independent of them, as long as the Lipschitz constants C_ρ , C_λ , C_g and the length $\sigma^* - \sigma_*$ of $D(\beta)$ vary in a bounded subset of $\mathbf{R}_+$.

Theorem 2.2. Assume that $1 \leq N \leq 3$ and (A1)-(A5) hold. Let $\{h_n\}, \{h_{on}\}, \{u_{on}\}$ and $\{w_{on}\}$ be bounded sequences in $L^2(0, T; H)$, $W^{1,2}(0, T; L^2(\Gamma))$, H and V , respectively, and assume that $\{\hat{\beta}(w_{on})\}$ is bounded in $L^1(\Omega)$. Further suppose that as $n \rightarrow \infty$

$$h_n \rightarrow h \quad \text{in } L^2(0, T; H), \quad h_{on} \rightarrow h_o \quad \text{in } L^2(0, T; L^2(\Gamma))$$

and

$$u_{on} \rightarrow u_o \quad \text{in } H, \quad w_{on} \rightarrow w_o \quad \text{in } V.$$

Then we have the following statements (i) and (ii):

(i) Suppose that

$$\sigma_* < m_n := \frac{1}{|\Omega|} \int_{\Omega} w_{on} dx < \sigma^* \quad \text{for all } n, \quad (2.11)$$

and

$$\sigma_* < m := \frac{1}{|\Omega|} \int_{\Omega} w_o dx < \sigma^*.$$

Let $\{u_n, w_n\}$ be the solution of $(P_n) := (P; h_n, h_{on}, u_{on}, w_{on})$ for each n and $\{u, w\}$ be the solution of $(P) := (P; h, h_o, u_o, w_o)$. Then, as $n \rightarrow \infty$,

$$u_n \rightarrow u \quad \text{in } L^2(0, T; H),$$

$$\rho(u_n) \rightarrow \rho(u) \quad \text{weakly in } H \text{ and uniformly in } t \in [0, T],$$

$$w_n \rightarrow w \quad \text{in } L^2(0, T; V) \text{ and weakly}^* \text{ in } L^\infty(0, T; V)$$

and

$$w'_n \rightarrow w' \quad \text{weakly in } L^2(0, T; V^*).$$

(ii) Suppose that (2.11) holds and

$$m = \sigma_* \text{ or } \sigma^*,$$

Then, for the solution $\{u_n, w_n\}$ of (P_n) , we have as $n \rightarrow \infty$,

$$w_n \rightarrow m \quad \text{in } C([0, T]; H)$$

$$u_n \rightarrow u \quad \text{in } L^2(0, T; H) \text{ and weakly in } L^2(0, T; V)$$

and

$$\rho(u_n) \rightarrow \rho(u) \quad \text{weakly in } H \text{ and uniformly in } t \in [0, T],$$

where $u \in C([0, T]; H) \cap W_{loc}^{1,2}([0, T]; H) \cap L_{loc}^\infty([0, T]; V) \cap L^2(0, T; V)$ is the unique solution of

$$\frac{d}{dt}(\rho(u(t)), z) + a(u(t), z) + (n_o u(t) - h_o(t), z)_\Gamma = (h(t), z)$$

$$\text{for all } z \in V, \text{ a.e. } t \in [0, T], \quad (2.12)$$

$$u(0) = u_o.$$

Remark 2.3. In (ii) of Theorem 2.2, moreover if $h \in L^\infty(Q)$, $h_o \in L^\infty(\Sigma)$, $u_o \in L^\infty(\Omega)$ and $m \in D(\beta)$, then the pair $\{u, w\}$, with the solution u of (2.12) and $w = m$, is the solution of (P) in the sense of Definition 2.1. In fact, under such restrictions on the data we see that $u \in L^\infty(Q)$ and hence $\xi := k - g(m) + \lambda'(m)u \in \beta(m)$ on Q for a certain constant k . Thus condition (w4) of Definition 2.1 is satisfied.

3. Sketch of proofs

(1) (Uniqueness) The uniqueness of the solution of (P) can be proved by using Gronwall's inequality with the help of the following embedding inequalities:

$$|z|_{L^q(\Omega)} \leq C_o |\nabla z|_H, \quad |z|_{L^q(\Omega)} \leq \delta |\nabla z|_H + C_\delta |z|_{V^*}$$

for all $z \in V_o$ and $1 \leq q < 6$, where C_o is a positive constant, and δ is an arbitrary positive constant with a constant C_δ dependent only on δ .

(2) (Existence) For the construction of a solution of (P) we consider the approximate problem $(P)_\mu (= (P_\mu; h, h_o, u_o, w_o))$, with parameter $0 < \mu \leq 1$, to find a pair of functions $u_\mu : [0, T] \rightarrow V$ and $w_\mu : [0, T] \rightarrow H^2(\Omega)$ fulfilling the following conditions $(w1)_\mu$ -(w4) $_\mu$:

$$(w1)_\mu \quad u_\mu \in W^{1,2}(0, T; H) \cap L^\infty(0, T; V), w_\mu \in W^{1,2}(0, T; H) \cap L^\infty(0, T; V) \cap L^2(0, T; H^2(\Omega));$$

$$(w2)_\mu \quad u_\mu(0) = u_o \text{ and } w_\mu(0) = w_o;$$

$$(w3)_\mu \quad \text{for a.e. } t \in [0, T] \text{ and all } z \in V,$$

$$(\rho(u_\mu)'(t) + \lambda(w_\mu)'(t), z) + a(u_\mu(t), z) + (n_o u_\mu(t) - h_o(t), z)_\Gamma = (h(t), z); \quad (3.1)$$

$$(w4)_\mu \quad \text{for a.e. } t \in [0, T],$$

$$\frac{\partial w_\mu(t)}{\partial n} = 0 \quad \text{a.e. on } \Gamma, \quad (3.2)$$

and there is a function $\xi_\mu \in L^2(0, T; H)$ such that

$$\xi_\mu \in \beta(w_\mu) \quad \text{a.e. on } Q \quad (3.3)$$

and

$$\begin{aligned} (w'_\mu(t), \eta) - \mu(w'_\mu(t), \Delta \eta) + \nu(\Delta w_\mu(t), \Delta \eta) \\ - (g(w_\mu(t)) - \lambda'(w_\mu(t))u_\mu(t) + \xi_\mu(t), \Delta \eta) = 0 \end{aligned} \quad (3.4)$$

for all $\eta \in H^2(\Omega)$ with $\frac{\partial \eta}{\partial n} = 0$ a.e. on Γ and a.e. $t \in [0, T]$.

Besides we reformulate $(P)_\mu$ as a system of evolution equations including subdifferential operators. For this purpose, let us introduce convex functions φ on H_o and ψ^t , $t \leq t \leq T$, on H as follows:

$$\varphi(z) := \begin{cases} \frac{\nu}{2} |\nabla z|_H^2 + \int_\Omega \hat{\beta}(z+m) dx & \text{if } z \in V_o \text{ and } \hat{\beta}(z+m) \in L^1(\Omega), \\ \infty & \text{otherwise,} \end{cases} \quad (3.5)$$

where

$$m := \frac{1}{|\Omega|} \int_\Omega w_o dx,$$

and

$$\psi^t(z) := \begin{cases} \frac{1}{2} |\nabla z|_H^2 + \frac{n_o}{2} |z|_{L^2(\Gamma)}^2 - (h_o(t), z)_\Gamma & \text{if } z \in V, \\ \infty & \text{otherwise.} \end{cases} \quad (3.6)$$

We then consider the subdifferential $\partial\varphi$ of φ in H_o and the subdifferential $\partial\psi^t$ of ψ in H . It is easy to see that

(i) $z^* \in \partial\varphi(z)$ if and only if $z^* \in H_o$, $z \in V_o \cap (K_m(\hat{\beta}) - m)$ and

$$(z^*, v - z)_o \leq \nu a(z, v - z) + \int_\Omega \hat{\beta}(v + m) dx - \int_\Omega \hat{\beta}(z + m) dx$$

$$\text{for all } v \in V_o \cap (K_m(\hat{\beta}) - m);$$

(ii) $\partial\psi^t$ is singlevalued, and $z^* = \partial\psi^t(z)$ if and only if $z^* \in H$, $z \in V$ and

$$(z^*, v) = a(z, v) + (n_o z - h_o(t), v)_\Gamma \quad \text{for all } v \in V.$$

For each $\mu \in (0, 1]$, problem $(P)_\mu$ has at most one solution and we have:

Lemma 3.1. *Let $\sigma_* < m < \sigma^*$, and $\lambda_1(r) := \lambda(r + m)$ and $g_1(r) := g(r + m)$ for $r \in \mathbf{R}$. Then a pair $\{u_\mu, w_\mu\}$ of functions is a solution of $(P)_\mu$ if and only if the pair $\{u_\mu, v_\mu\}$ with $v_\mu := w_\mu - m$ is a solution of the problem $(P)_\mu'$ defined below:*

(P) $'_\mu$ Find a pair $\{u_\mu, v_\mu\}$ of functions satisfying the following conditions $(w1)_\mu' - (w4)_\mu'$:

(w1) $'_\mu$ $u_\mu \in W^{1,2}(0, T; H) \cap L^\infty(0, T; V)$ and $v_\mu \in W^{1,2}(0, T; H_o) \cap L^\infty(0, T; V_o)$;

(w2) $'_\mu$ $u_\mu(0) = u_o$ and $v_\mu(0) = v_o := w_o - m$;

(w3) $'_\mu$ for a.e. $t \in [0, T]$,

$$\rho(u_\mu)'(t) + \lambda_1(v_\mu)'(t) + \partial\psi^t(u_\mu(t)) = h(t); \quad (3.7)$$

(w4) $'_\mu$ for a.e. $t \in [0, T]$,

$$(J_o^* + \mu I)v_\mu'(t) + \partial\varphi(v_\mu(t)) + \pi[g_1(v_\mu(t)) - \lambda_1'(v_\mu(t))u_\mu(t)] \ni 0. \quad (3.8)$$

We can prove Lemma 3.1 by using the following lemma which is concerned with the Lagrange multipliers of elliptic variational inequalities.

Lemma 3.2. *Let $\sigma_* < m < \sigma^*$ and ℓ be any element of H . Consider the following two problems (M_m) and $(M_m)'$:*

(M_m) *Find a function $z_m \in K_m(\hat{\beta}) \cap V$ such that*

$$\nu a(z_m, z_m - \eta) + \int_{\Omega} \hat{\beta}(z_m) dx \leq (\ell, z_m - \eta) + \int_{\Omega} \hat{\beta}(\eta) dx \quad \text{for all } \eta \in K_m(\hat{\beta}) \cap V.$$

(M_m)' *Find a function $z_m \in K_m(\hat{\beta}) \cap H^2(\Omega)$, $\gamma_m \in \mathbf{R}$ and $\xi_m \in H$ such that*

$$-\nu \Delta z_m + \xi_m = \ell + \gamma_m \quad \text{in } \Omega$$

and

$$\xi_m \in \beta(z_m) \quad \text{a.e. on } \Omega, \quad \frac{\partial z_m}{\partial n} = 0 \quad \text{a.e. on } \Gamma.$$

Then $(M_m)'$ has a solution $\{z_m, \xi_m, \gamma_m\}$ and the function z_m is the unique solution of (M_m) . Moreover, γ_m can be chosen so that

$$|\gamma_m| \leq 4M^5(1 + |\ell|_H), \quad (3.9)$$

where $M = \max\{\frac{1}{\delta}, r(\delta), \sigma^* - \sigma_*, |\Omega|, \frac{1}{|\Omega|}\}$ for δ and $r(\delta)$ satisfying (2.9) and (2.10); z_m satisfies that

$$(-\Delta z_m, \xi_m) \geq 0 \quad (3.10)$$

and

$$\nu |\Delta z_m|_H \leq |\ell|_H + |\gamma_m| |\Omega|^{\frac{1}{2}}. \quad (3.11)$$

For the detail proof of Lemma 3.2 we refer to [9; Proposition 5.1]. Thanks to the additional term $\mu v'_\mu$ problem $(P_\mu)'$, hence (P_μ) , is uniquely solved in the Hilbert spaces H and H_o by applying time-dependent subdifferential techniques evolved in [4, 10]. In fact, we have the following result.

Proposition 3.1. *In addition to all the conditions of Theorem 2.1, assume that $u_o \in V$. Then, for each $\mu \in (0, 1]$, problem $(P)_\mu$ has one and only one solution $\{u_\mu, w_\mu\}$. Moreover, the solution $\{u_\mu, w_\mu\}$ satisfies the bounds of the following type:*

$$\begin{aligned} & |u_\mu|_{C([0,T];H)} + |\nabla u_\mu|_{L^2(0,T;H)} + |w'_\mu|_{L^2(0,T;V^*)} \\ & + \mu |w'_\mu|_{L^2(0,T;H)}^2 + |w_\mu|_{L^\infty(0,T;V)} + |\hat{\beta}(w_\mu)|_{L^\infty(0,T;L^1(\Omega))} \\ & \leq \tilde{R}_o(|u_o|_H, |w_o|_V, |\hat{\beta}(w_o)|_{L^1(\Omega)}, |h|_{L^2(0,T;H)}, |h_o|_{L^2(0,T;L^2(\Gamma))}), \end{aligned}$$

where $\tilde{R}_o : \mathbf{R}_+^5 \rightarrow \mathbf{R}_+$ is a function which is independent of μ and bounded on each bounded subset of $\mathbf{R}_+^5$;

$$|w_\mu|_{L^2(0,T;H^2(\Omega))} + |\rho(u_\mu)'|_{L^1(0,T;V^*)} + |\lambda(w_\mu)'|_{L^1(0,T;V^*)}$$

$$\leq \tilde{R}_1\left(\frac{1}{\delta}, r(\delta), |u_o|_H, |w_o|_V, |\hat{\beta}(w_o)|_{L^1(\Omega)}, |h|_{L^2(0,T;H)}, |h_o|_{L^2(0,T;L^2(\Gamma))}\right),$$

where $\tilde{R}_1 : \mathbf{R}_+^7 \rightarrow \mathbf{R}_+$ is a function which is independent of μ and bounded on each bounded subset of $\mathbf{R}_+^7$, δ is an arbitrary number satisfying (2.9) and $r(\delta)$ is a constant given by (2.10).

By the above proposition we obtain a solution $\{u, w\}$ of (P), passing to the limit in $\mu \rightarrow 0$, and see that the solution satisfies estimates (2.7) and (2.8).

(3) (Proof of Theorem 2.2) The assertions of Theorem 2.2 follow easily from estimates for the solution of (P) in Theorem 2.1.

Remark 3.1. In this paper the domain $D(\beta)$ of β is supposed to be bounded in $\mathbf{R}$. However this is not essential for the assertions of Theorems 2.1 and 2.2. For instance, our results can be extended to the case when $\text{int}.D(\beta) \neq \emptyset$ and there are constants $k_\beta > 0$ and $k'_\beta > 0$ such that

$$|\beta(r)| \geq k_\beta |r| - k'_\beta \quad \text{for all } r \in D(\beta);$$

note that under this condition we may assume that

$$\hat{\beta}(r) \geq \hat{k}_\beta |r|^2 \quad \text{for all } r \in D(\hat{\beta}),$$

where $\hat{k}_\beta > 0$ is a certain constant.

Application. As a typical example of maximal monotone graphs β in $\mathbf{R} \times \mathbf{R}$ arising in the context of phase separation (cf. [3]), we consider an increasing smooth function $\beta^c : (0, 1) \rightarrow \mathbf{R}$ defined by

$$\beta^c(w) := c \log \frac{w}{1-w}$$

with positive real parameter c . Also, as an example of non-smooth β , we consider the subdifferential β^0 of the indicator function of the interval $[0, 1]$ in $\mathbf{R}$, which is the limit of β^c as $c \rightarrow 0$ in the sense of maximal monotone graphs in $\mathbf{R} \times \mathbf{R}$.

By virtue of Theorem 2.1, problem (P) with $\beta = \beta^c$ ($c \geq 0$) has one and only one solution $\{u^c, w^c\}$, provided that $u_o \in H$, $w_o \in V$ with $0 < m < 1$ and $\log \frac{w_o}{1-w_o} \in L^1(\Omega)$, $h \in L^2(0, T; H)$ and $h_o \in W^{1,2}(0, T; L^2(\Gamma))$. Moreover, it easily follows from the estimates (2.7), (2.8) and the uniqueness of solutions to (P) that as $c \rightarrow 0$, the solution $\{u^c, w^c\}$ converges to the solution $\{u^0, w^0\}$ in the similar sense as in (i) of Theorem 2.2.

References

- [1] H. W. Alt and I. Pawlow, Dynamics of non-isothermal phase separation, in *Free Boundary Value Problems*, K.-H. Hoffmann and J. Sprekels ed., ISNM 95, Birkhäuser, Basel, 1990, pp. 1-26.
- [2] H. W. Alt and I. Pawlow, Existence of solutions for non-isothermal phase separation, preprint.

- [3] F. E. Blowey and C. M. Elliott, The Cahn-Hilliard gradient theory for phase separation with non-smooth free energy, Part I: Mathematical analysis, *European J. Appl. Math.* 2(1991), 233-280.
- [4] P. Colli and A. Visintin, On a class of doubly nonlinear evolution equations, *Comm. Partial Differential Equations* 15(1990), 737-756.
- [5] S. R. DeGroot and P. Mazur, *Non-Equilibrium Thermodynamics*, Dover Publ., New York, 1984.
- [6] E. DiBenedetto and R. E. Showalter, A pseudo-parabolic variational inequality and Stefan problem, *Nonlinear Anal. TMA* 6(1982), 279-291.
- [7] C. E. Elliott, The Cahn-Hilliard model for the kinetics of phase separation, in *Mathematical Models for Phase Change Problems*, J. F. Rodrigues ed., ISNM 88, Birkhäuser, Basel, 1989, pp. 35-73.
- [8] C. M. Elliott and S. Zheng, On the Cahn-Hilliard equation, *Arch. Rat. Mech. Anal.* 96(1986), 339-357.
- [9] N. Kenmochi, M. Niezgódka and I. Pawlow, Subdifferential operator approach to the Cahn-Hilliard equation with constraint, preprint.
- [10] N. Kenmochi and I. Pawlow, A class of nonlinear elliptic-parabolic equations with time-dependent constraints, *Nonlinear Anal. TMA* 10(1986), 1181-1202.
- [11] S. Luckhaus and A. Visintin, Phase transition in multicomponent systems, *Manuscripta Math.* 43(1983), 261-288.
- [12] Y. Oono and S. Puri, Study of the phase separation dynamics by use of cell dynamical systems, I. Modelling., *Phys. Rev. A* 38(1988), 434-453.
- [13] R. Temam, *Infinite Dimensional Dynamical Systems in Mechanics and Physics*, Springer Verlag, Berlin, 1988.
- [14] W. von Wahl, On the Cahn-Hilliard equation $u' + \Delta^2 u - \Delta f(u) = 0$, *Delft Progress Report* 10(1985), 291-310.
- [15] S. Zheng, Asymptotic behaviour of the solution to the Cahn-Hilliard equation, *Applicable Anal.* 23(1986), 165-184.

N. Kenmochi: Department of Mathematics, Faculty of Education, Chiba University
1-33 Yayoi-cho, Chiba, 260 Japan

M. Niezgódka: Institute of Applied Mathematics and Mechanics, Warsaw University
Banacha 2, 00-913 Warsaw, Poland