

# Weak and strong convergence theorems for monotone nonexpansive-type mappings in Banach spaces

Sachiko Atsushiba

Department of Mathematics, Tokyo Woman's Christian University,  
2-6-1 Zempukuji, Suginami-ku, Tokyo 167-8585, Japan

## Abstract

In this paper, we prove weak and strong convergence theorems for finite monotone nonexpansive mappings in uniformly convex Banach spaces endowed with a partial order.

## 1 Introduction

Let  $E$  be a real Banach space, let  $C$  be a nonempty subset of  $E$ . A mapping  $T : C \rightarrow C$  is called nonexpansive if

$$\|Tx - Ty\| \leq \|x - y\|$$

for all  $x, y \in C$ . For a mapping  $T : C \rightarrow C$ , we denote by  $F(T)$  the set of *fixed points* of  $T$ , i.e.,  $F(T) = \{z \in C : Tz = z\}$ .

Ran and Reurings [16] proved an analogue of the classical Banach contraction principle [5] in metric spaces endowed with a partial order. In particular, they show how this extension is useful when dealing with some special matrix equations (see also [11, 15, 22, 23]).

Mann [14] introduced an iteration process for approximation of fixed points of a mapping  $T$  in a Hilbert space as follows:

$$x_{n+1} = \alpha_n x_n + (1 - \alpha_n)Tx_n$$

---

E-mail address: s-atsushiba@lab.twcu.ac.jp

*Keywords. and phrases.* fixed point, monotone mapping, nonexpansive mapping, weak convergence, ordered Banach space

*2020 Mathematics Subject Classification* 47H05, 47H07, 47H09, 47H25

for all  $n \geq 1$ , where  $\{\alpha_n\}$  are sequences in  $[0, 1]$ . Later, Reich [13] discussed this iteration process in a uniformly convex Banach space whose norm is Frechet differentiable. Bin Dehaish and Khamsi [8] proved a weak convergence theorem of Mann's type [14] iteration for monotone nonexpansive mappings in Banach spaces endowed with a partial order (see also [1, 18]).

In this paper, we prove weak convergence theorems for finite monotone nonexpansive mappings in uniformly convex Banach spaces endowed with a partial order.

## 2 Preliminaries and notations

Throughout this paper, we assume that  $E$  is a real Banach space with norm  $\|\cdot\|$  and endowed with a *partial order*  $\preceq$  compatible with the linear structure of  $E$ , that is,

$$x \preceq y \text{ implies } x + z \preceq y + z,$$

$$x \preceq y \text{ implies } \lambda x \preceq \lambda y$$

for every  $x, y, z \in E$  and  $\lambda \geq 0$ . As usual we adopt the convention  $x \succeq y$  if and only if  $y \preceq x$ . It follows that all *order intervals*  $[x, \rightarrow) = \{z \in E : x \preceq z\}$  and  $(\leftarrow, y] = \{z \in E : z \in E : z \preceq y\}$  are convex. Moreover, we assume that each order intervals  $[x, \rightarrow)$  and  $(\leftarrow, y]$  are closed. Recall that an order interval is any of the subsets  $[a, \rightarrow) = \{x \in X; a \preceq x\}$  or  $(\leftarrow, a] = \{x \in X; x \preceq a\}$ . for any  $a \in E$ . As a direct consequence of this, the subset

$$[a, b] = \{x \in X; a \preceq x \preceq b\} = [a, \rightarrow) \cap (\leftarrow, b]$$

is also closed and convex for each  $a, b \in E$ .

Let  $E$  be a real Banach space with norm  $\|\cdot\|$  and endowed with a partial order  $\preceq$  compatible with the linear structure of  $E$ . We will say that this Banach space  $(E, \|\cdot\|, \preceq)$  is an *ordered Banach space*. Let  $C$  be a nonempty subset of  $E$ . A mapping  $T : C \rightarrow C$  is called *nonexpansive* if

$$\|Tx - Ty\| \leq \|x - y\|$$

for all  $x, y \in C$ . A mapping  $T : C \rightarrow C$  is called *monotone* if

$$Tx \preceq Ty$$

for each  $x, y \in C$  such that  $x \preceq y$  (see also [8]). For a mapping  $T : C \rightarrow C$ , we denote by  $F(T)$  the set of *fixed points* of  $T$ , i.e.,  $F(T) = \{z \in C : Tz = z\}$ . For a mapping  $T : C \rightarrow C$  and  $\varepsilon > 0$ , we define the set  $F_\varepsilon(T)$  to be

$$F_\varepsilon(T) = \{x \in C : \|Tx - x\| \leq \varepsilon\}.$$

We denote by  $E^*$  the topological dual space of  $E$ . We denote by  $\mathbb{N}$  and  $\mathbb{Z}^+$  the set of all positive integers and the set of all nonnegative integers, respectively. We also denote by  $\mathbb{R}$  and  $\mathbb{R}^+$  the set of all real numbers and the set of all nonnegative real numbers, respectively.

We write  $x_n \rightarrow x$  (or  $\lim_{n \rightarrow \infty} x_n = x$ ) to indicate that the sequence  $\{x_n\}$  of vectors in  $E$  converges strongly to  $x$ . We also write  $x_n \rightharpoonup x$  (or  $w\text{-}\lim_{n \rightarrow \infty} x_n = x$ ) to indicate that the sequence  $\{x_n\}$  of vectors in  $E$  converges weakly to  $x$ . We also denote by  $\langle y, x^* \rangle$  the value of  $x^* \in E^*$  at  $y \in E$ . For a subset  $A$  of  $E$ ,  $\text{co}A$  and  $\overline{\text{co}}A$  mean the convex hull of  $A$  and the closure of convex hull of  $A$ , respectively.

A Banach space  $E$  is said to be strictly convex if

$$\frac{\|x+y\|}{2} < 1$$

for  $x, y \in E$  with  $\|x\| = \|y\| = 1$  and  $x \neq y$ . In a strictly convex Banach space, we have that if

$$\|x\| = \|y\| = \|(1-\lambda)x + \lambda y\|$$

for  $x, y \in E$  and  $\lambda \in (0, 1)$ , then  $x = y$ . For every  $\varepsilon$  with  $0 \leq \varepsilon \leq 2$ , we define the modulus  $\delta(\varepsilon)$  of convexity of  $E$  by

$$\delta(\varepsilon) = \inf \left\{ 1 - \frac{\|x+y\|}{2} : \|x\| \leq 1, \|y\| \leq 1, \|x-y\| \geq \varepsilon \right\}.$$

A Banach space  $E$  is said to be uniformly convex if  $\delta(\varepsilon) > 0$  for every  $\varepsilon > 0$ . If  $E$  is uniformly convex, then for  $r, \varepsilon$  with  $r \geq \varepsilon > 0$ , we have  $\delta\left(\frac{\varepsilon}{r}\right) > 0$  and

$$\left\| \frac{x+y}{2} \right\| \leq r \left( 1 - \delta\left(\frac{\varepsilon}{r}\right) \right)$$

for every  $x, y \in E$  with  $\|x\| \leq r$ ,  $\|y\| \leq r$  and  $\|x-y\| \geq \varepsilon$ . It is well-known that a uniformly convex Banach space is reflexive and strictly convex.

The following lemmas were proved in [10].

**Lemma 2.1** ([10]). *Let  $E$  be a uniformly convex Banach space and let  $C$  be a nonempty bounded closed convex subset of  $E$ . Then, for each  $\varepsilon > 0$ , there exists  $\delta > 0$  such that*

$$\overline{\text{co}}F_\delta(T) \subset F_\varepsilon(T)$$

*for every nonexpansive mapping  $T$  of  $C$  into itself.*

**Lemma 2.2** ([10]). *Let  $E$  be a uniformly convex Banach space and let  $C$  be a nonempty bounded closed convex subset of  $E$ . Then,*

$$\lim_{n \rightarrow \infty} \sup_{\substack{x \in C \\ T \in N(C)}} \left\| \frac{1}{n} \sum_{i=0}^{n-1} T^i x - T \left( \frac{1}{n} \sum_{i=0}^{n-1} T^i x \right) \right\| = 0$$

*where  $N(C)$  denotes the set of all nonexpansive mappings of  $C$  into itself.*

The following theorem was proved in [9].

**Theorem 2.3** ([9]). *Let  $C$  be a nonempty bounded closed convex subset of a uniformly convex Banach space  $E$  and let  $T$  be a nonexpansive mapping of  $C$  into itself. Let  $\{x_n\}$  be a sequence in  $C$  such that it converges weakly to an element  $u$  in  $C$  and  $\{x_n - Tx_n\}$  converges strongly to 0. Then,  $u$  is a fixed point of  $T$ .*

The following lemma was proved in [24].

**Lemma 2.4** ([24]). *Let  $p > 1$  and  $r > 0$  be two fixed real numbers. Let  $E$  be a uniformly convex Banach space. Then, there is a continuous, strictly increasing and convex function  $g : [0, \infty) \rightarrow [0, \infty)$  such that  $g(0) = 0$  and*

$$\begin{aligned} & \|\lambda x + (1 - \lambda)y\|^p \\ & \leq \lambda \|x\|^p + (1 - \lambda)\|y\|^p - \{\lambda^p(1 - \lambda) + \lambda(1 - \lambda)^p\}g(\|x - y\|) \end{aligned}$$

for all  $x, y \in B_r$  and  $\lambda$  with  $0\lambda < 1$  where  $B_r = \{x \in E : \|x\| \leq r\}$

### 3 Lemmas

In this paper, we consider the following iteration process for approximation of common fixed points of finite nonexpansive mappings in a Banach space: Let  $C$  be a nonempty closed convex subset of an ordered Banach space  $E$ . Let  $T_1, T_2, \dots, T_r$  be finite nonexpansive mappings of  $C$  into itself such that  $T_i T_j = T_j T_i$  for every  $i, j = 1, 2, \dots, r$  and  $\bigcap_{i=1}^r F(T_i)$  is nonempty. For  $x_1 \in C$ ,  $\{x_n\}$  is defined by

$$x_{n+1} = \alpha_n x_n + (1 - \alpha_n) \cdot \frac{1}{(n+1)^r} \sum_{i_1=0}^n \cdots \sum_{i_r=0}^n T_1^{i_1} \cdots T_r^{i_r} x_n$$

for every  $n \in \mathbb{N}$ , where  $\{\alpha_n\}$  is a sequence in  $[0, 1]$  (see also [14]).

Let  $C$  be a nonempty subset of an ordered Banach space  $E$  and let  $T$  be a mapping of  $C$  into itself. A sequence  $\{x_n\}$  in  $C$  is said to be an *approximate fixed point sequence* of a mapping  $T$  of  $C$  into itself if

$$\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$$

(see also [13, 19]). Let  $T_1, T_2, \dots, T_r$  be mappings of  $C$  into itself. A sequence  $\{x_n\}$  in  $C$  is said to be an *approximate common fixed point sequence* of mappings  $T_1, T_2, \dots, T_r$  of  $C$  if for every  $k = 1, 2, \dots, r$ ,

$$\lim_{n \rightarrow \infty} \|x_n - T_k x_n\| = 0.$$

A sequence  $\{x_n\}$  in  $E$  is said to be *monotone increasing* if

$$x_1 \preceq x_2 \preceq x_3 \preceq \cdots$$

(see also [8]).

The following lemma plays an important role in this paper (see also [3]).

**Lemma 3.1.** Let  $E$  be a uniformly convex Banach space, let  $C$  be a nonempty bounded closed convex subset of  $E$ . Let  $T_1, T_2, \dots, T_r$  be nonexpansive mappings of  $C$  into itself such that  $T_i T_j = T_j T_i$  for every  $i, j = 1, 2, \dots, r$ . For any  $n \in \mathbb{N}$ , put

$$S_n^{l_1, \dots, l_r} x = \frac{1}{n^r} \sum_{i_1=0}^{n-1} \dots \sum_{i_r=0}^{n-1} T_1^{i_1+l_1} \dots T_r^{i_r+l_r} x \quad \text{for every } x \in C.$$

Then, for every  $k = 1, 2, \dots, r$ ,

$$\lim_{n \rightarrow \infty} \sup_{\substack{x \in C \\ l_1, \dots, l_r \in \mathbb{Z}^+}} \left\| S_n^{l_1, \dots, l_r} x - T_k (S_n^{l_1, \dots, l_r} x) \right\| = 0. \quad (3.1)$$

**Lemma 3.2** ([2]). Let  $E$  be a Banach space and let  $C$  be a nonempty closed convex subset of  $E$ . Let  $T_1, T_2, \dots, T_r$  be nonexpansive mappings of  $C$  into itself such that  $T_i T_j = T_j T_i$  for every  $i, j = 1, 2, \dots, r$  and  $\bigcap_{i=1}^r F(T_i)$  is nonempty. Suppose  $x_1 \in C$  and  $\{x_n\}$  is given by

$$x_{n+1} = \alpha_n x_n + (1 - \alpha_n) \frac{1}{(n+1)^r} \sum_{i_1=0}^n \dots \sum_{i_r=0}^n T_1^{i_1} \dots T_r^{i_r} x_n \quad \text{for every } n \in \mathbb{N},$$

where  $\{\alpha_n\}$  is a sequence in  $[0, 1]$ . Let  $w$  be a common fixed point of  $T_1, T_2, \dots, T_r$ . Then,  $\lim_{n \rightarrow \infty} \|x_n - w\|$  exists.

**Lemma 3.3** ([2]). Let  $C$  be a nonempty closed convex subset of an ordered uniformly convex Banach space  $E$ . Let  $T_1, T_2, \dots, T_r$  be monotone nonexpansive mappings of  $C$  into itself such that  $T_i T_j = T_j T_i$  for every  $i, j = 1, 2, \dots, r$ , and  $\bigcap_{i=1}^r F(T_i)$  is nonempty. Suppose  $x_1 \in C$  and  $\{x_n\}$  is given by

$$x_{n+1} = \alpha_n x_n + (1 - \alpha_n) \frac{1}{(n+1)^r} \sum_{i_1=0}^n \dots \sum_{i_r=0}^n T_1^{i_1} \dots T_r^{i_r} x_n \quad \text{for every } n \in \mathbb{N},$$

where  $\{\alpha_n\}$  is a sequence in  $[0, a]$  for some  $0 < a < 1$ . Then,  $\{x_n\}$  is an approximate common fixed point sequence of mappings  $T_1, T_2, \dots, T_r$  of  $C$ , i.e., for every  $k = 1, 2, \dots, r$ ,

$$\lim_{n \rightarrow \infty} \|x_n - T_k x_n\| = 0.$$

**Lemma 3.4** ([2]). Let  $C$  be a nonempty closed convex subset of an ordered Banach space  $E$ . Let  $T_1, T_2, \dots, T_r$  be monotone nonexpansive mappings of  $C$  into itself such that  $T_i T_j = T_j T_i$  for every  $i, j = 1, 2, \dots, r$ , and  $\bigcap_{i=1}^r F(T_i)$  is nonempty. Assume that  $x \preceq T_k x$  for every  $k = 1, 2, \dots, r$  and  $x \in C$ . Suppose  $x_1 \in C$  and  $\{x_n\}$  is given by

$$x_{n+1} = \alpha_n x_n + (1 - \alpha_n) \frac{1}{(n+1)^r} \sum_{i_1=0}^n \dots \sum_{i_r=0}^n T_1^{i_1} \dots T_r^{i_r} x_n \quad \text{for every } n \in \mathbb{N},$$

where  $\{\alpha_n\}$  is a sequence in  $[0, 1]$ . Then,  $\{x_n\}$  is monotone increasing.

## 4 Convergence theorems of Mann's type iteration

In this section, we provide the approximation of common fixed points of families of monotone nonexpansive mappings in ordered Banach spaces. It is connected with the theory of differential equations (see [12]). We assume that  $E$  is an ordered Banach space in the sense that

$$\begin{aligned}x \preceq y \text{ implies } x + z &\preceq y + z, \\x \preceq y \text{ implies } \lambda x &\preceq \lambda y\end{aligned}$$

for every  $x, y, z \in E$  and  $\lambda \geq 0$ . Note that this is a very typical situation.

We get a weak convergence theorem for a family of monotone nonexpansive mappings in an ordered uniformly convex Banach space.

**Theorem 4.1** ([2]). *Let  $C$  be a nonempty closed convex subset of an ordered uniformly convex Banach space  $E$ . Let  $T_1, T_2, \dots, T_r$  be monotone nonexpansive mappings of  $C$  into itself such that  $T_i T_j = T_j T_i$  for every  $i, j = 1, 2, \dots, r$  and  $\bigcap_{i=1}^r F(T_i)$  is nonempty. Let  $x \in C$ . Assume that  $x \preceq T_k x$  for every  $k = 1, 2, \dots, r$ . Suppose  $x_1 = x \in C$  and  $\{x_n\}$  is given by*

$$x_{n+1} = \alpha_n x_n + (1 - \alpha_n) \frac{1}{(n+1)^r} \sum_{i_1=0}^n \cdots \sum_{i_r=0}^n T_1^{i_1} \cdots T_r^{i_r} x_n \quad \text{for every } n \in \mathbb{N},$$

where  $\{\alpha_n\}$  is a sequence in  $[0, 1]$ . If  $\{\alpha_n\}$  is chosen so that  $\alpha_n \in [0, a]$  for some  $a$  with  $0 < a < 1$ , then the sequence  $\{x_n\}$  converges weakly to a point of  $\bigcap_{i=1}^r F(T_i)$ .

**Theorem 4.2.** *Let  $C$  be a nonempty closed convex subset of an ordered uniformly convex Banach space  $E$ . Let  $T_1, T_2, \dots, T_r$  be monotone nonexpansive mappings of  $C$  into itself such that  $T_i T_j = T_j T_i$  for every  $i, j = 1, 2, \dots, r$  and  $\bigcap_{i=1}^r F(T_i)$  is nonempty. Let  $x \in C$ . Assume that  $x \preceq T_k x$  for every  $k = 1, 2, \dots, r$ . Suppose  $x_1 = x \in C$  and  $\{x_n\}$  is given by*

$$x_{n+1} = \alpha_n x_n + (1 - \alpha_n) \frac{1}{(n+1)^r} \sum_{i_1=0}^n \cdots \sum_{i_r=0}^n T_1^{i_1} \cdots T_r^{i_r} x_n \quad \text{for every } n \in \mathbb{N},$$

where  $\{\alpha_n\}$  is a sequence in  $[0, 1]$ . Let  $\{\alpha_n\}$  be a sequence in  $[0, 1]$ . Then,  $\{x_n\}$  converges strongly to a point of  $\bigcap_{i=1}^r F(T_i)$  if and only if  $\lim_{n \rightarrow \infty} d\left(x_n, \bigcap_{i=1}^r F(T_i)\right) = 0$

Using Theorem 4.1, we get some convergence theorems for monotone nonexpansive mappings in ordered uniformly convex Banach spaces (see also [18]).

**Theorem 4.3** ([2]). *Let  $C$  be a nonempty closed convex subset of an ordered uniformly convex Banach space  $E$ . Let  $S$  and  $T$  be monotone nonexpansive mappings of  $C$  into itself such that  $ST = TS$  and  $F(S) \cap F(T) \neq \emptyset$ . Let  $x \in C$ . Assume that  $x \preceq Tx$  and  $x \preceq Sx$ . Suppose  $x_1 = x \in C$  and  $\{x_n\}$  is given by*

$$x_{n+1} = \alpha_n x_n + (1 - \alpha_n) \frac{1}{(n+1)^2} \sum_{i=0}^n \sum_{j=0}^n S^i T^j x_n \quad \text{for every } n \in \mathbb{N},$$

where  $\{\alpha_n\}$  is a sequence in  $[0, 1]$ . If  $\{\alpha_n\}$  is chosen so that  $\alpha_n \in [0, a]$  for some  $a$  with  $0 < a < 1$ ,  $\{x_n\}$  converges weakly to a point of  $F(T) \cap F(S)$ .

**Theorem 4.4** ([2]). Let  $C$  be a nonempty closed convex subset of an ordered uniformly convex Banach space  $E$  and let  $T$  be a monotone nonexpansive mapping of  $C$  into itself such that  $F(T) \neq \emptyset$ . Let  $x \in C$ . Assume that  $x \preceq Tx$ . Suppose  $x_1 = x \in C$  and  $\{x_n\}$  is given by

$$x_{n+1} = \alpha_n x_n + (1 - \alpha_n) \frac{1}{n+1} \sum_{i=0}^n T^i x_n \quad \text{for every } n \in \mathbb{N},$$

where  $\{\alpha_n\}$  is a sequence in  $[0, 1]$ . If  $\{\alpha_n\}$  is chosen so that  $\alpha_n \in [0, a]$  for some  $a$  with  $0 < a < 1$ ,  $\{x_n\}$  converges weakly to a point of  $F(T)$ .

## Acknowledgements

The author is supported by Grant-in-Aid for Scientific Research No. 19K03582 from Japan Society for the Promotion of Science.

## References

- [1] S. Atsushiba, *A nonlinear mean convergence theorem for two monotone nonexpansive mappings in Banach space*, Linear and Nonlinear Anal., **8** (2022), 155-162.
- [2] S. Atsushiba, *A nonlinear mean convergence theorem for two monotone nonexpansive mappings in Banach space*, Linear and Nonlinear Anal., to appear in Fixed Point Theory, **26** (2) (2025).
- [3] S. Atsushiba and W. Takahashi, *Approximating common fixed point of two nonexpansive mappings in Banach spaces*, Bull. Austral. Math. Soc. **57** (1998), 117–127.
- [4] J.-B. Baillon, *Un theoreme de type ergodique pour les contractions non lineaires dans un espace de Hilbert*, C. R. Acad. Sci. Paris Ser. A-B, **280** (1975), 1511-1514.
- [5] S. Banach, *Sur les opérations dans les ensembles abstraits et leurs applications*, Fundam. Math. **3** (1922), 133-181.
- [6] M. Bachar and M. A. Khamsi, *Fixed points of monotone mappings and application to integral equations*, Fixed Point Theory Appl., **2015:110**, (2015).
- [7] M. Bachar and M. A. Khamsi, *On common approximate fixed points of monotone non-expansive semigroups in Banach spaces*, Fixed Point Theory Appl., **2015:160**, (2015).
- [8] B.A. Bin Dehaish and M.A. Khamsi, *Mann iteration process for monotone nonexpansive mappings*, Fixed Point Theory Appl., **2015:177**, (2015).

- [9] F. E. Browder, Nonlinear operators and nonlinear equations of evolution in Banach spaces. Nonlinear functional analysis (Proc. Sympos. Pure Math., Vol. XVIII, Part 2, Chicago, Ill., 1968), Amer. Math. Soc., Providence, R.I., 1976.
- [10] R.E. Bruck, *On the convex approximation property and the asymptotic behavior of nonlinear contractions in Banach spaces*, Israel J. Math. **38** (1981), 304-314.
- [11] J. Jachymski, *The contraction principle for mappings on a metric space with a graph*, Proc. Amer. Math. Soc. **136** (2007), 1359-1373.
- [12] W.M. Kozłowski, *Monotone Lipschitzian semigroup in Banach spaces*, J. Aust. Math. Soc. **105** (2018), 417–428.
- [13] E. Matoušková and S. Reich, *Reflexivity and approximate fixed points*, Studia Math. **159** (2003), 403–415.
- [14] W.R. Mann, *Mean value methods in iteration*, Proc. Amer. Math. Soc. **4** (1953), 506–510.
- [15] J.J. Nieto, Rodríguez-López, *Contractive mapping theorems in partially ordered sets and applications to ordinary differential equations*, Order **22** (2005), 223-239.
- [16] A.C.M. Ran and M.C.B. Reurings, *A fixed point theorem in partially ordered sets and some applications to matrix equations*, Proc. Amer. Math. Soc. **132** (2004), 1435-1443.
- [17] S. Reich, *Weak convergence theorems for nonexpansive mappings in Banach spaces*, J. Math. Anal. Appl. **67** (1979), 274–276.
- [18] R. Shukla and A. Wiśnicki, *Iterative methods for monotone nonexpansive mappings in uniformly convex Banach spaces*, Adv. Nonlinear Anal. **10** (2021), 1061–1070
- [19] W. Takahashi, Nonlinear Functional Analysis, Yokohama Publishers, Yokohama, 2000.
- [20] W. Takahashi and K. Shimoji, *Convergence Theorems for Nonexpansive Mappings and Feasibility Problems*, Mathematical and Computer Modelling **32** (2000), 1463-1471.
- [21] W. Takahashi and T. Tamura, *Limit theorems of operators by convex combinations for nonexpansive retractions in Banach spaces*, J. Approximation Theory **91** (1997), 386-397.
- [22] M. Turinici, *Abstract comparison principles and multivariable Gronwall-Bellman inequalities*, J. Math. Anal. Appl. **117** (1986), 100-127.
- [23] M. Turinici, *Fixed points for monotone iteratively local contractions*, Demonstr. Math. **19** (1986), 171-180.
- [24] H.K. Xu, *Inequalities in Banach spaces with applications*, Nonlinear Anal. **16** (1991), 1127-1138.