

Review on Several Nonlinear Mappings

Tae Hwa Kim

Department of Applied Mathematics, College of Natural Sciences

Pukyong National University, Busan 608-737, Korea

E-mail: taehwa@pknu.ac.kr

Abstract

The purpose of this paper is systematically to survey several nonlinear mappings and to classify the implications concerning to them. Also, we suggest several examples of nonlinear mappings which are comparable each other and we finally raise some open questions.

Keywords: non-Lipschitzian mapping, total asymptotically nonexpansive mapping, generalized projection, relatively total asymptotically nonexpansive, generalized total asymptotically nonexpansive.

2010 Mathematics Subject Classification. 46B20, 46N10, 47H09.

1 Introduction

Let X be a real Banach space with norm $\|\cdot\|$ and let X^* be the dual of X . Denote by $\langle \cdot, \cdot \rangle$ the duality product. Also, we denote by $\mathbb{N}$ and $\mathbb{R}$ the set of natural numbers and the set of real numbers, respectively. Let $T : C \rightarrow C$ be a mapping. We denote by $F(T)$ the set of all fixed points of T , namely,

$$F(T) = \{x \in C : Tx = x\}.$$

Recall that the mapping T is said to be *Lipschitzian* if

$$\|Tx - Ty\| \leq L\|x - y\|, \quad x, y \in C,$$

where $L := L_T$ denotes the *Lipschitz constant* of T . Obviously, it is equivalent to the following property: for each $n \in \mathbb{N}$, there exists a constant $k_n > 0$ such that

$$\|T^n x - T^n y\| \leq k_n \|x - y\|, \quad x, y \in C. \quad (1.1)$$

For a Lipschitzian mapping T , we say:

- T is *uniformly k -Lipschitzian* if $k_n = k$ for all $n \in \mathbb{N}$;
- T is *nonexpansive* if $k_n = 1$ for all $n \in \mathbb{N}$;
- T is *asymptotically nonexpansive* [3] if $\lim_{n \rightarrow \infty} k_n = 1$.

The first non-Lipschitzian mapping was introduced by Kirk [10]; we say that T is a mapping of *asymptotically nonexpansive type* if

$$\limsup_{n \rightarrow \infty} \sup_{y \in C} (\|T^n x - T^n y\| - \|x - y\|) \leq 0 \quad (1.2)$$

for every $x \in C$, and T^N is continuous for some $N \geq 1$. In 1993, Bruck et al [2] introduced the stronger definition than (1.2), namely, T is said to be *asymptotically nonexpansive in the intermediate sense* [2] provided T is uniformly continuous and

$$\limsup_{n \rightarrow \infty} \sup_{x, y \in C} (\|T^n x - T^n y\| - \|x - y\|) \leq 0. \quad (1.3)$$

Note that if we define

$$c_n := \sup_{x, y \in C} (\|T^n x - T^n y\| - \|x - y\|) \vee 0, \quad (1.4)$$

where $a \vee b := \max\{a, b\}$, then (1.3) ensures that $c_n \rightarrow 0$ and

$$\|T^n x - T^n y\| \leq \|x - y\| + c_n \quad (1.5)$$

for all $x, y \in C$ and $n \geq 1$. Obviously, (1.5) implies (1.3) in case $c_n \rightarrow 0$. Therefore, we summarize:

Proposition 1.1. (1.3) holds $\Leftrightarrow$ (1.5) holds for some sequence $\{c_n\}$ with $c_n \rightarrow 0$.

Definition 1.2. We say that T is *gradually nonexpansive* whenever (1.5) holds for some sequence $\{c_n\}$ with $c_n \rightarrow 0$.

For the purpose of unifying nonlinear mappings mentioned above, Alber et al [1] introduced a new notion, namely, T is said to be *total asymptotically nonexpansive* [1] if there exist two nonnegative real sequences $\{\alpha_n\}$ and $\{\beta_n\}$ with $\alpha_n, \beta_n \rightarrow 0$, $\tau \in \Gamma[0, \infty)$ such that

$$\|T^n x - T^n y\| \leq \|x - y\| + \alpha_n \tau(\|x - y\|) + \beta_n, \quad x, y \in C, \quad n \geq 1, \quad (1.6)$$

where $\tau \in \Gamma[0, \infty)$ if and only if $\tau : [0, \infty) \rightarrow [0, \infty)$ is strictly increasing, continuous on $[0, \infty)$ and $\tau(0) = 0$.

Remark 1.3. In view of Definition 3.1 of [12], we also say that T is *generalized asymptotically nonexpansive* in case that $\tau(t) = t$ for all $t \geq 0$ in (1.6).

Now it is natural to consider more stronger one than (1.6).

Definition 1.4. T is said to be *square total asymptotically nonexpansive* if (1.6) can be replaced by

$$\|T^n x - T^n y\|^2 \leq \|x - y\|^2 + \tilde{\alpha}_n \tilde{\tau}(\|x - y\|^2) + \tilde{\beta}_n, \quad (1.7)$$

for all $x, y \in C$ and $n \geq m_0$, where $m_0 \in \mathbb{N}$, $\tilde{\alpha}_n, \tilde{\beta}_n \rightarrow 0$ and $\tilde{\tau} \in \Gamma[0, \infty)$.

Remark 1.5. Note that the property (1.6) with $\alpha_n = 0$ for all $n \geq 1$ reduces to (1.5) with $\beta_n = c_n$; moreover, if we take $\tau(t) = t$ for all $t \geq 0$ and $\beta_n = 0$ for all $n \geq 1$ in (1.6), it is reduced to (1.1) with $k_n = 1 + \alpha_n$.

The purpose of this paper is systematically to survey several nonlinear mappings and to classify the implications concerning to them. In section 2, we introduce classes of various nonlinear mappings and suggest their implications. In section 3, we give some counter examples for their implications and some open questions are finally added.

2 Implications of classes

For summarizing the connections between the classes of nonlinear mappings considered above, we use the following notations:

- (N) = the class of nonexpansive mappings
- (L) = the class of Lipschitzian mappings
- (UC) = the class of uniformly continuous mappings
- (UL) = the class of uniformly Lipschitzian mappings
- (AN) = the class of asymptotically nonexpansive mappings
- (GN) = the class of gradually nonexpansive mappings
- (GAN) = the class of generalized asymptotically nonexpansive mappings
- (TAN) = the class of total asymptotically nonexpansive mappings
- (STAN) = the class of square total asymptotically nonexpansive mappings
- (ANIS) = the class of mappings which are asymptotically nonexpansive in the intermediate sense
- (ANT) = the class of mappings of asymptotically nonexpansive type

We say that T is AN, GN, GAN, TAN, STAN, ANIS and ANT, in abbreviated forms, whenever T belongs to its corresponding classes (AN), (GN), (GAN), (TAN), (ANIS) and (ANT). Then, there hold the following implications:

Proposition 2.1. (i) $(N) \subset (AN) \subset (UL) \subset (L) \subset (UC)$.

(ii) $(GN) \cap (UC) = (ANIS) \subset (ANT)$.

(iii) $(AN) \cup (GN) \subset (GAN) \subset (TAN)$.

Proposition 2.2. Let C be bounded. Then:

(i) $(GN) = (GAN) = (TAN) = (STAN)$.

(ii) $(AN) \subset (ANIS)$.

Proof. (i) Since $(TAN) = (STAN)$ is obvious, we claim: $(TAN) \subset (GN)$. Assume $\delta := \text{diam}(C) < \infty$. If T is TAN, then

$$\begin{aligned}
 \|T^n x - T^n y\| &\leq \|x - y\| + \alpha_n \tau(\|x - y\|) + \beta_n \\
 &\leq \|x - y\| + \alpha_n \tau(\delta) + \beta_n \\
 &= \|x - y\| + c_n, \quad x, y \in C, \quad n \geq 1,
 \end{aligned}$$

where $c_n := \alpha_n \tau(\delta) + \beta_n \delta \rightarrow 0$. Hence T is GN. It follows from (iii) of Proposition 2.1 that $(GN) = (GAN) = (TAN)$. Moreover, since $(AN) \cup (GN) = (GN)$, we have $(AN) \subset (GN)$. Then it follows from (i) and (ii) of Proposition 2.1 that $(AN) \subset (ANIS)$. Therefore (ii) is obtained. $\square$

Definition 2.3. Let $f_n, f : C \subset X \rightarrow C$ be mappings. We say that $\{f_n\}$ converges uniformly to f on C if

$$\|f_n - f\| := \sup_{x \in C} \|f_n(x) - f(x)\| \rightarrow 0$$

as $n \rightarrow \infty$. Further, we say that $T^n x$ converges uniformly to a point $p \in C$ on C whenever $\{f_n := T^n\}$ converges uniformly to the function f on C , where $f(x) = p$ for all $x \in C$, that is,

$$\sup_{x \in C} \|T^n x - p\| \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Lemma 2.4. *If $T^n x$ converges uniformly to some point $p \in C$ on C , then T is GN.*

Proof. Let $c_n := \sup_{x,y \in C} \|T^n x - T^n y\|$. Then $c_n \rightarrow 0$ since

$$0 \leq c_n = \sup_{x,y \in C} \|T^n x - T^n y\| \leq \sup_{x \in C} \|T^n x - p\| + \sup_{y \in C} \|p - T^n y\| \rightarrow 0$$

as $n \rightarrow \infty$. From the construction of c_n it easily follows that

$$\|T^n x - T^n y\| \leq c_n, \quad x, y \in C, \quad n \geq 1,$$

which immediately implies (1.5). Hence T is GN. $\square$

Remark 2.5. However, the converse of Lemma 2.4 does not hold in general; see Example 3.9.

3 Counter examples

As counter examples of (i) in Proposition 2.1, it is not hard to see that if $Tx = 2x$ for $x \in C = \mathbb{R}$, then $T \in (L) \setminus (UL)$. Furthermore, if $Tx = \sqrt{x}$ for $x \in C = [0, \infty)$, then $T \in (UC) \setminus (L)$. For an example of $T \in (UL) \setminus (AN)$, see Example 3.7. Also, the following example of $T \in (AN) \setminus (N)$ is originally due to [3] in ℓ^2 spaces.

Example 3.1. ([7]; see Example 3.13). *Let B denote the unit ball in the space $X = \ell^p$, where $1 < p < \infty$. Obviously, X is uniformly convex and uniformly smooth. Let $T : B \rightarrow B$ be defined by*

$$Tx = (0, x_1^2, \lambda_1 x_2, \lambda_2 x_3, \dots)$$

for all $x = (x_1, x_2, x_3, \dots) \in B$, where $0 < \lambda_n < 1$ for all $n \geq 1$ and $\prod_{n=1}^{\infty} \lambda_n = \frac{1}{2}$. Then:

- (i) T is Lipschitzian, i.e., $\|Tx - Ty\| \leq 2\|x - y\|$, $x, y \in C$;
- (ii) T is AN, i.e., $\|T^{n+1}x - T^{n+1}y\| \leq 2 \prod_{i=1}^{n-1} \lambda_i \|x - y\|$, $x, y \in C$, $n \in \mathbb{N}$;
- (iii) T is not nonexpansive.

Proof. Noticing that, for $x = (x_1, x_2, \dots) \in B$,

$$T^n x = \left(\overbrace{0, \dots, 0}^n, \prod_{i=1}^{n-1} \lambda_i x_1^2, \prod_{i=1}^n \lambda_i x_2, \prod_{i=2}^{n+1} \lambda_i x_3, \dots \right).$$

Thus we have $\|T^n x - T^n y\| \leq 2 \prod_{i=1}^{n-1} \lambda_i \|x - y\|$ for all $n \geq 2$. Obviously, since $2 \prod_{i=1}^{n-1} \lambda_i \downarrow 1$, T is AN. On the other hand, since $\|Tx - Ty\| = \frac{3}{4} > \frac{1}{2} = \|x - y\|$ for $x = (1, 0, 0, \dots)$ and $y = (1/2, 0, 0, \dots)$, T is not nonexpansive. $\square$

Remark 3.2. Consider either $\lambda_n := 1 - \frac{1}{(n+1)^2}$ or $\lambda_n := \exp\left(\frac{1}{2n} - \frac{1}{2n-1}\right)$ to get a sequence satisfying that $0 < \lambda_n < 1$ and $\prod_{n=1}^{\infty} \lambda_n = \frac{1}{2}$. Indeed, for the second case, since $0 < \exp(-x) < 1$ for all $x > 0$, we must find a sequence $\{\alpha_n\}$ such that

$$\prod_{n=1}^{\infty} \lambda_n = \prod_{n=1}^{\infty} \exp(-\alpha_n) = \exp\left(-\sum_{n=1}^{\infty} \alpha_n\right) = \frac{1}{2}.$$

This is equivalent to

$$\sum_{n=1}^{\infty} \alpha_n = \ln 2 = \sum_{n=1}^{\infty} \left(\frac{1}{2n-1} - \frac{1}{2n} \right)$$

Several Nonlinear Mappings

because

$$\begin{aligned} s_n &:= \sum_{k=1}^n \left(\frac{1}{2k-1} - \frac{1}{2k} \right) = \frac{1}{n+1} + \frac{1}{n+2} + \cdots + \frac{1}{n+n} \\ &= \frac{1}{n} \sum_{k=1}^n \frac{1}{1+k/n} \rightarrow \int_0^1 \frac{1}{1+x} dx = \ln 2. \end{aligned}$$

Furthermore, since $1-x \leq \exp(-x)$ for all $x \in (0, 1)$, we observe

$$\sum_{n=1}^{\infty} \alpha_n = \infty \Rightarrow \prod_{n=1}^{\infty} (1 - \alpha_n) = 0$$

for all $\alpha_n \in (0, 1)$ since

$$0 \leq \prod_{n=1}^{\infty} (1 - \alpha_n) \leq \prod_{n=1}^{\infty} \exp(-\alpha_n) = \exp\left(-\sum_{n=1}^{\infty} \alpha_n\right) = 0.$$

As a direct consequence of Lemma 2.4, we introduce three non-Lipschitzian mappings $T \in (\text{ANIS}) \setminus (\text{AN})$.

Example 3.3. ([5]). Let $C := [-\frac{1}{\pi}, \frac{1}{\pi}]$ and $0 < |k| < 1$. For each $x \in C$, let $T : C \rightarrow C$ be defined by

$$Tx := \begin{cases} kx \sin \frac{1}{x}, & \text{if } x \neq 0; \\ 0, & \text{if } x = 0. \end{cases}$$

Then $F(T) = \{0\}$ and $T^n x$ converges uniformly to 0 on C (hence T is GN by Lemma 2.4). Since T is clearly uniformly continuous, it follows from (ii) of Proposition 2.1 that T is ANIS. However, T is not Lipschitzian; see Example 4.3 of [5] for the proof. Therefore $T \in (\text{ANIS}) \setminus (\text{AN})$.

Example 3.4. ([9]). Let $X = \mathbb{R}$ and $C = [0, 1]$. For each $x \in C$, let $T : C \rightarrow C$ be defined by

$$Tx = \begin{cases} \alpha, & x \in [0, \alpha]; \\ \frac{\alpha}{\sqrt{1-\alpha}} \sqrt{1-x}, & x \in [\alpha, 1], \end{cases}$$

where $\alpha \in (0, 1)$. Then $F(T) = \{\alpha\}$ and $T^n x = \alpha$ for all $x \in C$, $n \geq 2$; hence T is ANIS as in Example 3.3. However, T is not Lipschitzian; see Example 3.9 of [9] for the proof. Therefore $T \in (\text{ANIS}) \setminus (\text{AN})$.

Example 3.5. ([4]). Let $X = \mathbb{R}$ and $C = [0, 1]$. For each $x \in C$, let $T : C \rightarrow C$ be defined by

$$Tx = \begin{cases} (\sqrt{2}-1)\sqrt{\frac{1}{2}-x} + \frac{1}{\sqrt{2}}, & \text{if } 0 \leq x \leq 1/2; \\ \sqrt{x}, & \text{if } 1/2 \leq x \leq 1. \end{cases}$$

Then $F(T) = \{1\}$ and $T^n x$ converges uniformly to 1 on C ; hence T is ANIS as in Example 3.3. However, T is not Lipschitzian; see Example 1.2 of [4] for more details. Therefore $T \in (\text{ANIS}) \setminus (\text{AN})$.

A mapping satisfying the property (1.3) do not always guarantee its *non-Lipschitz*. The following two examples are *uniformly Lipschitzian* ANIS mappings.

Example 3.6. ([8]). Let $X = \mathbb{R}$ and $C = [0, 1]$. For each $x \in C$, let $T : C \rightarrow C$ be defined by

$$Tx = \begin{cases} kx, & \text{if } 0 \leq x \leq 1/2; \\ \frac{k}{2k-1}(k-x), & \text{if } 1/2 \leq x \leq k; \\ 0, & \text{if } k \leq x \leq 1, \end{cases}$$

where $1/2 < k < 1$. Then $F(T) = \{0\}$ and $T^n x$ converges uniformly to 0 on C . Obviously, T is uniformly continuous. It follows from Lemma 2.4 and (ii) of Proposition 2.1 that T is ANIS. Furthermore, T is uniformly Lipschitzian. Indeed, if $0 \leq x \leq 1/2$ and $1/2 \leq y \leq k$, then $T^n x = k^n x$ and $T^n y = \frac{k^n}{2k-1}(k-y)$. Therefore, we see that

$$\begin{aligned} |T^n x - T^n y| &= \left| k^n x - \frac{k^n}{2} + \frac{k^n}{2} - \frac{k^n}{2k-1}(k-y) \right| \\ &= \left| k^n \left(x - \frac{1}{2} \right) + \frac{k^n}{2k-1} \left[\left(k - \frac{1}{2} \right) - (k-y) \right] \right| \\ &\leq k^n \left| x - \frac{1}{2} \right| + \frac{k^n}{2k-1} \left| y - \frac{1}{2} \right| \\ &\leq \frac{k^n}{2k-1} |x - y| \leq \frac{k}{2k-1} |x - y|. \end{aligned}$$

The remaining cases are obvious. Hence T is uniformly $\frac{k}{2k-1}$ -Lipschitzian.

The following example of $T \in (\text{UL}) \cap (\text{ANIS}) \setminus (\text{AN})$ is originally due to Example 1.3 (with $k = 4$) of [4].

Example 3.7. ([4]). For any $k > 0$, let $\{a_n\}$ be a sequence of positive numbers such that $a_n \downarrow 0$ and $\prod_{n=1}^{\infty} (1 + a_n) = k$. Set

$$b_n := \frac{1}{2^{n+1}(1 + a_n)}, \quad n \geq 1.$$

Let $T : C \rightarrow C$ be defined by

$$Tx = \begin{cases} (1 + a_1)x + 1/2, & \text{if } x \in [0, b_1]; \\ 1/2 + 1/4, & \text{if } x \in [b_1, 1/2] \end{cases}$$

and

$$Tx = \begin{cases} (1 + a_n) \left(x - \sum_{i=1}^{n-1} \frac{1}{2^i} \right) + \sum_{i=1}^n \frac{1}{2^i}, & \text{if } x \in \left[\sum_{i=1}^{n-1} \frac{1}{2^i}, \sum_{i=1}^{n-1} \frac{1}{2^i} + b_n \right]; \\ \sum_{i=1}^{n+1} \frac{1}{2^i}, & \text{if } x \in \left[\sum_{i=1}^{n-1} \frac{1}{2^i} + b_n, \sum_{i=1}^n \frac{1}{2^i} \right], \quad n \geq 2 \end{cases}$$

and $T1 = 1$. Then $F(T) = \{1\}$ and $T^n x$ converges uniformly to 1 on C . Since T is continuous on C , T is also uniformly continuous on C . It follows from Lemma 2.4 and (ii) of Proposition 2.1 that T is ANIS. Furthermore, T is uniformly k -Lipschitzian. Note that if $k > 1$, we conclude : $T \in (\text{UL}) \cap (\text{ANIS}) \setminus (\text{AN})$.

Proof. It suffices to show that T is uniformly k -Lipschitzian. Indeed, (i) if we take

$$x := \prod_{i=1}^n b_i = \frac{1}{\prod_{i=1}^n 2^{i+1} \cdot \prod_{i=1}^n (1 + a_i)}, \quad y := \frac{b_n}{\prod_{i=1}^{n-1} (1 + a_i)}$$

for each $n \geq 1$, then, since

$$T^n x = \prod_{i=1}^n (1 + a_i) x + \sum_{i=1}^n \frac{1}{2^i}, \quad T^n y = \sum_{i=1}^{n+1} \frac{1}{2^i}$$

Several Nonlinear Mappings

and $\prod_{k=1}^n (1 + a_k)y = b_n(1 + a_n) = \frac{1}{2^{n+1}}$, we obtain that

$$\begin{aligned} |T^n x - T^n y| &= \left| \prod_{i=1}^n (1 + a_i)x + \sum_{i=1}^n \frac{1}{2^i} - \sum_{i=1}^{n+1} \frac{1}{2^i} \right| \\ &= \left| \prod_{i=1}^n (1 + a_i)x - \frac{1}{2^{n+1}} \right| \\ &= \left| \prod_{i=1}^n (1 + a_i)x - b_n(1 + a_n) \right| = \prod_{i=1}^n (1 + a_i)|x - y|. \end{aligned}$$

(ii) if we replace y with a point $p \in [b_1, 1/2]$ in the case (i), then $T^n p = \sum_{i=1}^{n+1} \frac{1}{2^i}$. Now use $1/2^2 = (1 + a_1)b_1$ and $\prod_{i=1}^n (1 + a_i)x = 1/(\prod_{i=1}^n 2^{i+1})$ to derive

$$\begin{aligned} |T^n x - T^n p| &= |T^n x - T^n b_1| \\ &= \left| \prod_{i=1}^n (1 + a_i)x + \sum_{i=1}^n \frac{1}{2^i} - \sum_{i=1}^{n+1} \frac{1}{2^i} \right| \\ &= \left| \prod_{i=1}^n (1 + a_i)x - \frac{1}{2^{n+1}} \right| = \frac{1}{2^{n+1}} - \frac{1}{\prod_{i=1}^n 2^{i+1}} \\ &\leq \frac{\prod_{i=2}^n (1 + a_i)}{2^2} - \frac{1}{\prod_{i=1}^n 2^{i+1}} \\ &= \prod_{i=1}^n (1 + a_i)b_1 - \prod_{i=1}^n (1 + a_i)x = \prod_{i=1}^n (1 + a_i)|x - b_1| \\ &\leq \prod_{i=1}^n (1 + a_i)|x - p|. \end{aligned}$$

The remaining cases are obvious. We conclude that

$$|T^n x - T^n y| \leq \prod_{i=1}^n (1 + a_i)|x - y|, \quad x, y \in C, \quad n \geq 1,$$

which ensures that T is uniformly k -Lipschitzian since $\prod_{n=1}^{\infty} (1 + a_n) = k$. $\square$

Remark 3.8. (i) Since $1 + x \leq e^x$ for all $x \in \mathbb{R}$, we easily find a sequence $\{a_n\}$ of positive numbers such that $a_n \downarrow 0$ and $\prod_{n=1}^{\infty} (1 + a_n) = k (> 1)$. Indeed, since the sequence $\{s_n\}$, $s_n := \prod_{k=1}^n (1 + a_k)$, of its n th partial sums is strictly increasing and

$$\prod_{n=1}^{\infty} (1 + a_n) \leq \prod_{n=1}^{\infty} e^{a_n} = e^{\sum_{n=1}^{\infty} a_n} = e^{\ln k} = k,$$

it suffices to find a (convergent) geometric series replaced with $a_n := r^n$, $0 \leq r < 1$ such that

$$\begin{aligned} \sum_{n=1}^{\infty} a_n &= \sum_{n=1}^{\infty} r^n = \frac{r}{1 - r} = \ln k \\ \Leftrightarrow r &= \frac{\ln k}{1 + \ln k}. \end{aligned}$$

(ii) Note that if we take $a_n \equiv 0$ in Example 3.7, then T is clearly nonexpansive.

As a slight modification of Example 3.7, we shall give an example of a uniformly Lipschitzian ANIS mapping defined a (unbounded) closed convex subset C on which is not converges uniformly.

Example 3.9. Consider $C := [0, \infty) \subset \mathbb{R}$. Let T be defined on $[0, 1]$ as in Example 3.7 and define $Tx = x$ on $[1, \infty)$. Since $T^n x$ converges uniformly to 1 on $[0, 1]$, setting $c_n := \sup\{|T^n x - T^n y| : x, y \in [0, 1]\} \rightarrow 0$, then it is not hard to see that

$$|T^n x - T^n y| \leq |x - y| + c_n, \quad x, y \in C, \quad n \geq 1.$$

Therefore T is GN. In view of Example 3.7, T is uniformly k -Lipschitzian. Therefore, $T : C \rightarrow C$ is ANIS. However, note that $T^n x$ does not converge uniformly to a point $p \in C$ on C .

Proof. We claim : $|T^n x - T^n y| \leq |x - y| + c_n$ for all $x, y \in C$ and $n \geq 1$. Consider the case of $x \in [0, 1]$ and $y \in [1, \infty)$. Then

$$\begin{aligned} |T^n x - T^n y| &\leq |T^n x - T^n 1| + |T^n 1 - T^n y| \\ &\leq c_n + |1 - y| \leq |x - y| + c_n \end{aligned}$$

for all $n \geq 1$. The remaining cases are obvious. □

The following example of $T \in (\text{UL}) \setminus (\text{ANT})$ is also interesting.

Example 3.10. ([6]). Let $X := \mathbb{R}$, $C := [-\frac{1}{k}, 1]$, where $1 < k < 2$. Define a mapping $T : C \rightarrow C$ by

$$Tx := \begin{cases} -kx, & \text{if } -\frac{1}{k} \leq x \leq 0; \\ -\frac{1}{k}x, & \text{if } 0 \leq x \leq 1. \end{cases}$$

Then:

- (i) $T^2 x = x$ for all $x \in C$ (hence, $T^{2n-1} = T$ for all $n \geq 1$);
- (ii) T is uniformly k -lipschitzian;
- (iii) T does not satisfy (1.2); hence it is not ANT.

Indeed, it suffice to show: T is not ANT. To this end, for each $x \in C$,

$$\begin{aligned} &\limsup_{n \rightarrow \infty} \sup_{y \in C} \{|T^n x - T^n y| - |x - y|\} \\ &\geq \sup\{|Ty| - |y| : y \in [-1/k, 1]\} \\ &= \sup\{(k-1)|y| : -1/k \leq y \leq 0\} \\ &= (k-1) \frac{1}{k} = 1 - \frac{1}{k} > 0. \end{aligned}$$

Now we introduce a discontinuous mapping $T \in (\text{GN})$; see Example 1.6 of [11].

Example 3.11. ([11]). Let $X := \mathbb{R}$, $C := [0, 1]$ and $0 < k < 1$. Define a mapping $T : C \rightarrow C$ by

$$Tx := \begin{cases} kx, & \text{if } 0 \leq x \leq \frac{1}{2}; \\ 0, & \text{if } \frac{1}{2} < x \leq 1. \end{cases}$$

Then T is not continuous at $\frac{1}{2}$ but $T^n x$ converges uniformly to 0 on C . By Lemma 2.4 we readily see that T is GN.

Finally we raise the following questions.

Question 3.12. Find examples of the following mapping T :

- (i) $T \in (\text{ANT}) \setminus (\text{ANIS})$.
- (ii) $T \in (\text{ANT}) \setminus (\text{ANIS})$.
- (iii) $T \in (\text{GAN}) \setminus (\text{GN})$.

Question 3.13. Find the corresponding analogue examples in infinite dimensional spaces.

References

- [1] Ya.I. Alber, C.E. Chidume and H. Zegeye, Approximating fixed points of total asymptotically nonexpansive mappings, *Fixed Point Theory and Appl.* **2006** (2006) article ID 10673, 20 pages.
- [2] R.E. Bruck, T. Kuczumow and S. Reich, Convergence of iterates of asymptotically non-expansive mappings in Banach spaces with the uniform Opial property, *Colloq. Math.* **65** (1993) 169–179.
- [3] K. Goebel and W.A. Kirk, A fixed point theorem for asymptotically nonexpansive mappings, *Proc. Amer. Math. Soc.* **35** (1972), 171–174.
- [4] C.S. Hu and G. Gai, Convergence theorems for equilibrium problems and fixed point problems of a finite family of asymptotically k -strictly pseudocontractive mappings in the intermediate sense, *Comput. Math. Appl.* **61** (2011), 79–93.
- [5] T.H. Kim and J.W. Choi, Asymptotic behavior for almost-orbits of non-Lipschitzian mappings in Banach spaces, *Mathematica Japonica* **38** (1993), 191–197.
- [6] T.H. Kim and K.M. Park, Some results on metric fixed point theory and open problems, *Comm. Korean Math. Soc.* **11** (1996), 725–742.
- [7] T.H. Kim and H.J. Lee, Strong convergence of modified iteration processes for relatively nonexpansive mappings, *Kyungpook Math. J.* **48** (2008), 685–703.
- [8] T.H. Kim and D.H. Kim, Demiclosedness principle for continuous TAN mappings, “Nonlinear Analysis and Convex Analysis” W. Takahashi and T. Tanaka, eds., RIMS Kôyokûroku 1821, 2013, p. 90–106.
- [9] G.E. Kim and T.H. Kim, Strong convergence theorems of total asymptotically nonexpansive mappings, Proceedings of the 7th International Conference on Nonlinear Analysis and Convex Analysis (Busan, Korea, 2011), 1–12.
- [10] W.A. Kirk, Fixed point theorems for non-Lipschitzian mappings of asymptotically non-expansive type, *Israel J. Math.* **17** (1974), 339–346.
- [11] D.R. Sahu, H.K. Xu and J.C. Yao, Asymptotically strict pseudocontractive mappings in the intermediate sense, *Nonlinear Anal.* **70** (2009), 3502–3511.
- [12] N. Shahzad and H. Zegeye, Strong convergence of an implicit iteration process for a finite family of generalized asymptotically quasi-nonexpansive maps, *Appl. Math. Comput.* **189** (2007), 1058–1065.