

The Role of Disorder in Producing Supersolid Phases

Derek Larson¹ and Ying-Jer Kao¹

¹ Department of Physics, National Taiwan University, Taipei, TAIWAN

Recent experimental evidence [1] supports a picture where the glass state can play an important role in creating superflow, and some simulation results [2] find the existence of a novel "superglass" state. Furthermore, there are also findings that a *lack* of disorder and frustration cause the supersolid phase to become unstable [3, 4]. In order to better understand the relationship between disorder, frustration, and the superglass state, we consider a disordered Bose-Hubbard model and tune parameters controlling the level of disorder. We solve the model through quantum Monte Carlo simulations using the worm algorithm. A first study involves scanning the bimodal distribution probability $p(V_{ij} = +V)$ to see the transition from the ordered states $p \rightarrow 0, 1$ to the glassy intermediate states. Initial results suggest paired onset of superflow and glassy behavior for the parameters under consideration.

References

- [1] A.S.C. Rittner, J.D. Reppy, Phys. Rev. Lett. **98**, 175302 (2007).
- [2] Ka-Ming Tam, Scott Geraedts, Stephen Inglis, Michel J.P. Gingras, Roger G. Melko, Phys. Rev. Lett. **104**, 215301 (2010).
- [3] Stefan Wessel, Matthias Troyer, Phys. Rev. Lett. **95**, 127205 (2005).
- [4] Takahumi Suzuki, Naoki Kawashima, Phys. Rev. B **75**, 180502(R) (2007).