

Note


**A Rapid Enzymatic Preparation of [^{32}P]AMP
from [α - ^{32}P]ATP**

Jun OKA* and Kunihiro UEDA**

Received October 7, 1977

Adenosine 5'-mono[^{32}P]phosphate ([^{32}P]AMP) is used as a substrate for sensitive assays of 5'-nucleotidase (1, 2). It is commercially available from the Radiochemical Centre, Amersham, but the commercial product has a relatively low specific activity (0.5~3 Ci/mmol) and was provided only in a large package (≥ 1 mCi). When a relatively small amount of highly labeled [^{32}P]AMP is a demand, it seems convenient to prepare it from adenosine 5'-[α - ^{32}P]triphosphate ([α - ^{32}P]ATP) which is available with higher specific activity and in a relatively small quantity (0.5~250 Ci/mmol, ≥ 250 μCi from the Radiochemical Centre or 10~30 Ci/mmol, ≥ 100 μCi from New England Nuclear, Boston).

For the preparation of AMP from ATP, a chemical method (3) does not appear applicable to small-scale preparations. We recently developed a rapid and simple method using myokinase. Myokinase (adenylate kinase, ATP: AMP phosphotransferase, EC 2.7.4.3) catalyzes the following reaction;



The equilibrium constant at pH 7.4 and at 25° is reported to be 2.26 M (4). When the reaction starts with [α - ^{32}P]ATP and 10 or 100 times more nonradioactive AMP, the equilibrium is expected to be attained when approximately 83% or 98%, respectively, of the radioactivity is distributed to AMP. The procedure is very simple and applicable to any scale, albeit lowering of specific activity by about one or two orders.

EXPERIMENTAL

[α - ^{32}P]ATP (sodium salt, 250 μCi , 10 Ci/mmol) was obtained from the Radiochemical Centre. Myokinase (from rabbit muscle) was purchased from Boehringer/Mannheim-Yamanouchi, Tokyo. Polyethyleneimine cellulose (Polygram cel 300) was from Mesherey-Nagel and Co., Düren, and Dowex AG 1 $\times$ 2 from Bio-Rad Laboratories. Determination of radioactivity on thin-layer chromatograms was performed with Packard Radiochromatogram Scanner Model 7201.

[α - ^{32}P]ATP, freed of vehicle solvents by evaporation, was dissolved in 1.0 ml

* 岡 純: Department of Medical Chemistry, Faculty of Medicine, Kyoto University, Sakyo-ku, Kyoto.

** 上田国寛: Laboratory of Molecular Biology, Institute for Chemical Research, Kyoto University, Uji, Kyoto.

of distilled water. A 0.7-ml portion (3×10^8 cpm, 26 nmol) was transferred to a small test tube, and again evaporated to dryness. The dried material was dissolved, in the same tube, in the mixture (0.1 ml) containing 10 μ mol of Tris-HCl (pH 7.5), 2.6 μ mol of AMP, 1 μ mol of MgCl_2 and 7 units of myokinase. Incubation was carried out for 3 hours at 37°. Figure 1 shows radiochromatograms of aliquots




Fig. 1. Analysis of reaction products at various stages. Either the reaction mixture at zero time (A) or after 3 hour incubation (B) or AMP purified on Dowex AG 1 $\times$ 2 (C) was applied to a PEI cellulose sheet. Chromatography was performed with 1.0 M LiCl as a solvent system at room temperature (5). Authentic AMP, ADP, and ATP were cochromatographed as markers, and located on the chromatogram under UV light.

taken at zero-time (A) and after three hours of incubation (B). As judged by the distribution of radioactivity among three nucleotides, the reaction appeared to reach equilibrium within 3 hours. The reaction was terminated by adding 0.2 ml of 10% perchloric acid and the mixture neutralized with KOH. The supernatant fraction after brief centrifugation was applied to a Dowex AG 1 $\times$ 2 column (0.7 $\times$ 6 cm). The column was washed with 15 ml of water, and eluted with 30 ml of 0.3 N formic acid. Under this condition, neither ADP nor ATP elutes out from the column. The fractions containing [^{32}P]AMP were pooled and lyophilized. The recovery of radioactivity from [α - ^{32}P]ATP to AMP was 79%. Purity of [^{32}P]AMP thus prepared was verified by PEI cellulose thin-layer chromatography (Fig. 1C); no radioactive impurity was detectable.

Preparation of [^{32}P]AMP

A similar method is applicable to a preparation of [β - ^{32}P]ADP from [γ - ^{32}P]ATP (6, 7).

REFERENCES

- (1) C. C. Widnell, *Methods Enzymol.*, **32B**, 368 (1974).
- (2) J. Oka, K. Ueda, and O. Hayaishi, *Biochem. Biophys. Res. Commun.*, **80**, 841 (1978).
- (3) L. Berger, *Methods Enzymol.*, **3**, 866 (1957).
- (4) L. V. Eggleston and R. Hems, *Biochem. J.*, **52**, 156 (1952).
- (5) K. Randerath and E. Randerath, *Methods Enzymol.*, **12A**, 323 (1967).
- (6) S. Narindrasorasak and W. A. Bridger, *J. Biol. Chem.*, **252**, 3121 (1977).
- (7) K. Ueda and J. Oka, manuscript in preparation.