

Calculations for Broué's abelian defect group conjecture ブルエの可換不足群予想の計算

千葉大学 理学部 越谷 重夫 (Shigeo Koshitani)

Department of Mathematics and Informatics,

Faculty of Science, Chiba University

e-mail koshitan@math.s.chiba-u.ac.jp

This is a joint work with **Naoko Kunugi** and **Katsushi Waki**, and a detailed version of a result presented here is in [6].

It has been conjectured by Michel Broué that a block algebra of a finite group should be derived (Rickard) equivalent to a block algebra of the normalizer of a common defect group which correspond each other via the Brauer correspondence provided the defect group is abelian, see [2, 6.2.Question]. This is known as *Broué's Abelian Defect Group Conjecture*, (ADGC) for short. We have been continuing a project on Broué's ADGC for a specific defect group, say the elementary abelian group of order nine, see [3], [4], [5]. Our main result here is the following:

Theorem (Koshitani-Kunugi-Waki, 2005). *Let G be the Janko simple group J_4 , and let $(\mathcal{O}, \mathcal{K}, k)$ be a splitting 3-modular system for all subgroups of G , namely, $\mathcal{O}$ is a complete discrete valuation ring of rank one such that $\mathcal{K}$ is the quotient field of $\mathcal{O}$ with $\text{char}(\mathcal{K}) = 0$ and such that k is the residue field of $\mathcal{O}$, namely $k = \mathcal{O}/\text{rad}(\mathcal{O})$, with $\text{char}(k) = 3$, and $\mathcal{K}$ and k are both splitting fields for all subgroups of G . Let A be a unique block algebra of $\mathcal{O}G$ whose defect group P is elementary abelian of order 9, and let B be the Brauer correspondent of A in $\mathcal{O}H$ where $H = N_G(P)$. Then, A and B are derived (Rickard) equivalent. In fact, even stronger fact is proved, namely, A and B are splendidly derived (Rickard) equivalent, see [9] and [10].*

Remark. In our proof results in papers of Okuyama [7] and [8] are important.

Corollary. *It turns out that Broué's ADGC holds for any prime p and any block algebra of G . This means that Broué's ADGC is settled for all primes and all block algebras of J_4 .*

Proof. This follows immediately from Theorem and [1, Lemma 5.1].

Acknowledgment. The author is grateful to Professor Hiroki Sasaki for the nice meeting held in Kyoto, October 29 – September 2, 2005.

REFERENCES

- [1] J. An, E.A. O'Brien, R.A. Wilson, The Alperin's weight conjecture and Dade's conjecture for the simple group J_4 , LMS J. Comput. Math. **6** (2003), 119–140.
- [2] M. Broué, Isométries de caractères et équivalences de Morita ou dérivées, Publ. Math. IHES **71** (1990), 45–63.
- [3] S. Koshitani, N. Kunugi, Broué's conjecture holds for principal 3-blocks with elementary abelian defect group of order 9, J. Algebra **248** (2002), 575–604.
- [4] S. Koshitani, N. Kunugi, K. Waki, Broué's conjecture for non-principal 3-blocks of finite groups, J. Pure Appl. Algebra **173** (2002), 177–211.
- [5] S. Koshitani, N. Kunugi, K. Waki, Broué's abelian defect group conjecture for the Held group and the sporadic Suzuki group, J. Algebra **279** (2004), 638–666.
- [6] S. Koshitani, N. Kunugi, K. Waki, Broué's abelian defect group conjecture holds for the Janko simple group J_4 , 2005, in preparation.
- [7] T. Okuyama, Some examples of derived equivalent blocks of finite groups, preprint (1997).
- [8] T. Okuyama, Remarks on splendid tilting complexes, in: Representation theory of finite groups and related topics, edited by S. Koshitani, RIMS Kokyuroku **1149**, Proc. Research Institute for Mathematical Sciences, Kyoto University, 2000, 53–59.
- [9] J. Rickard, Splendid equivalences: derived categories and permutation modules, Proc. London Math. Soc. (3) **72** (1996), 331–358.
- [10] J. Rickard, Equivalences of derived categories for symmetric algebras, J. Algebra **257** (2002), 460–481.