

**Development of a regulatable low-temperature protein expression system using the
psychrotrophic bacterium, *Shewanella livingstonensis* Ac10, as the host**

Soichiro KAWAI, Jun KAWAMOTO, Takuya OGAWA, Tatsuo KURIHARA*

Institute for Chemical Research, Kyoto University, Kyoto, Japan

* Corresponding author: Gokasho, Uji, Kyoto 611-0011, Japan, Tel: +81-774-38-4710,
fax: +81-774-38-3248, e-mail: kurihara@scl.kyoto-u.ac.jp

Running Title: Regulatable low-temperature protein production system

Keywords: *trp* promoter, *Shewanella*, heterologous protein production, regulatable gene
expression

Abstract

A low-temperature protein expression system is useful for the production of thermolabile proteins. We previously developed a system that enables constitutive protein production at low temperatures, using the psychrotrophic bacterium *Shewanella livingstonensis* Ac10 as the host. To increase the utility of this system, in the present study, we introduced a repressible promoter of the *trp* operon of this bacterium into the system. When β -lactamase was produced under the control of this promoter at 18 °C and 4 °C, the yields were 75 and 33 mg/L-culture, respectively, in the absence of L-Trp, and the yields were decreased by 72% and 77%, respectively, in the presence of L-Trp. We also found that 3-indoleacrylic acid, a competitive inhibitor of the *Escherichia coli* *trp* repressor, increased the expression of the reporter gene. This repressible gene expression system would be useful for regulatable recombinant protein production at low temperatures.

1 **Introduction**

2 A number of recombinant protein expression systems have been constructed
3 using various organisms including bacteria [1], eukaryotic microorganisms [2], and
4 mammalian cells [3] as the hosts. However, there are still many proteins that are
5 difficult to overproduce using conventional recombinant protein expression systems.
6 These include low-stability [4] and toxic [5] proteins. To achieve efficient production of
7 these proteins, a low-temperature protein-expression system is expected to be useful. At
8 low temperatures, thermal denaturation of proteins is alleviated. Moreover, as the low
9 temperature expression system can suppress enzyme activity by lowering the
10 temperature, it is also effective for the production of enzymes that may harm the host
11 cells [6]. To date, cold-adapted microorganisms, such as Antarctic sea bacteria,
12 *Shewanella livingstonensis* Ac10 [7] and *Pseudoalteromonas haloplanktis* TAC125 [8],
13 have been employed for heterologous protein expression at low temperatures.

14 *S. livingstonensis* Ac10 grows well at low temperatures close to 0 °C but
15 cannot grow at temperatures above 30 °C [9]. In our previous work, we found a
16 promoter, named LI3, that leads to strong expression of heterologous proteins in this

strain and developed a low-temperature expression system using this strain as a host [7].

The LI3 promoter was obtained from the upstream noncoding region of the gene encoding a putative alkyl hydroperoxide reductase (AhpC), which is highly produced in this strain both at 18 °C and 4 °C. Under the control of this promoter, the maximum yield of the reporter protein β -lactamase (BLA) was 139 and 91 mg/L-culture at 18 °C and 4 °C, respectively, in the late stationary phase [7]. We also found that this protein production system is suitable for the production of thermolabile peptidases, PepF and PepQ, from a psychrophilic strain, *Desulfotalea psychrophila* DSM12343 [7,10]. The production of these proteins in *S. livingstonensis* Ac10 under the LI3 promoter system was higher than that under the *Escherichia coli* T7 promoter system. These results demonstrated that the LI3 promoter is useful for constitutive production of foreign proteins, especially those with low thermostability. Nevertheless, there is a drawback of this system: the LI3 promoter is not regulatable; thus, the expression of the gene under the control of this promoter cannot be repressed. This could be problematic in the production of highly toxic proteins, the toxicity of which cannot be sufficiently suppressed even at low temperatures. In this study, we introduced a gene regulatory

system of the *trp* operon into the heterologous protein production system operating at low temperatures using *S. livingstonensis* Ac10. We assessed its validity and found that the newly developed system is useful for regulatable expression of recombinant proteins at low temperatures.

Materials and methods

Bacterial strains, plasmids, and culture conditions

Bacterial strains used in this study are listed in Table 1. As the host, *S. livingstonensis* Ac10-Rif^r, a rifampicin resistant mutant of *S. livingstonensis* Ac10, was used. *S. livingstonensis* Ac10-Rif^r was grown under aerobic conditions in Luria-Bertani (LB) medium consisting of 10 g/L tryptone (BD Difco, Detroit, MI), 10 g/L NaCl (Nacalai Tesque, Kyoto, Japan), and 5 g/L yeast extract (BD Difco) for 24 h at 18 °C, and then transferred to modified DSMZ medium 79 (a minimal synthetic sea water medium containing 1 g/L KH₂PO₄, 1 g/L NH₄NO₃, 10 g/L NaCl, 0.2 g/L MgSO₄ · 7H₂O, 10 mg/L FeSO₄, and 10 mg/L CaCl₂ · 2H₂O) supplemented with 0.5% w/v casamino acids as carbon source (BD Difco), which was used as the basal medium

throughout the present study [11]. The cells grown in this medium at 18 °C and 4 °C were harvested in the log phase. As needed, 50 µg/mL of rifampicin (Sigma Aldrich, St. Louis, MO) and/or 30 µg/mL of chloramphenicol (Nacalai Tesque) were added. For regulation of gene expression, 0.1% (w/v) L-Trp (Kyowa Hakko Bio, Tokyo, Japan) or 40 µg/mL of 3-indoleacrylic acid (3-IAA) (Tokyo Chemical Industry, Tokyo, Japan) was added. *E. coli* DH5α was used as the host for plasmid construction, and *E. coli* S17-1/λ_{pir} was used as the donor strain in the conjugation experiments. *E. coli* DH5α and *E. coli* S17-1/λ_{pir} were grown in LB broth in the presence or absence of chloramphenicol (30 µg/mL). A derivative of a broad-host-range vector, pJRD215 [12], containing the chloramphenicol selection cassette, pJRD-Cm^r [13], was used to construct plasmids for protein expression in *S. livingstonensis* Ac10-Rif^r. Promoter-assay plasmids, pT1_{bla} to pT10_{bla}, constructed as described below were introduced into *S. livingstonensis* Ac10-Rif^r to obtain the strains BLAT1 to BLAT10, respectively. BLAT0 was prepared as a control strain harboring pJRD-Cm^r.

Construction of plasmids

Plasmids and primers used in this study are listed in Table 1 and Table S1, respectively. Q5 High-Fidelity Polymerase and NEBuilder HiFi DNA Assembly Master Mix were purchased from New England BioLabs, Ipswich, MA. Construction of the plasmids for the promoter assay is summarized in Figure 1.

Each of the predicted promoter regions was amplified from the upstream region of the *trpE* gene (Figure 2) by using the following primer sets: Trp1-Fw/Trp-Rv.1 for the T1 region, Trp2-Fw/Trp-Rv.1 for the T2 region, Trp3-Fw/Trp-Rv.1 for the T3 region, Trp4-Fw/Trp-Rv.1 for the T4 region, Trp5-Fw/Trp-Rv.1 for the T5 region, Trp6-Fw/Trp-Rv.2 for the T6 region, Trp7-Fw/Trp-Rv.1 for the T7 region, Trp8-Fw/Trp-Rv.1 for the T8 region, Trp9-Fw/Trp-Rv.2 for the T9 region, and Trp10-Fw/Trp-Rv.1 for the T10 region (Table S1 “Amplification of predicted promoter regions” section). By using the primer set, pJRD-Fw/pJRD-Rv, a linear plasmid fragment of pJRD-Cm^r was amplified with Q5 High-Fidelity DNA Polymerase. BLA-coding gene was amplified from pBR322 by using the primer set, β -lac-Fw.1/ β -lac-Rv.1 or β -lac-Fw.2/ β -lac-Rv.2. The purified linear plasmid fragment and each of the insert gene fragments consisting of the

predicted promoter regions and the BLA-coding gene were subjected to the NEBuilder HiFi DNA Assembly Cloning System to generate promoter-assay plasmids pT1*bla* to pT10*bla*.

Conjugative transformation of *S. livingstonensis* Ac10-Rif^r

pJRD-Cm^r and its derivatives were introduced into *E. coli* S17-1/ λ *pir* and then transferred to *S. livingstonensis* Ac10-Rif^r by conjugation as follows. *E. coli* S17-1/ λ *pir* harboring pJRD-Cm^r and its derivatives was cultivated at 37 °C aerobically in 5 mL of LB medium with 30 µg/ml chloramphenicol until the optical density at 600 nm (OD₆₀₀) reached about 1.0. *S. livingstonensis* Ac10-Rif^r was cultivated in 5 mL of LB medium at 18 °C aerobically until the OD₆₀₀ reached about 2.0. These cells were washed twice with fresh LB medium without antibiotics. After washing, 100 µL of the cultures of each of the donor cells and the recipient cells were mixed, and the mixture was incubated on LB plate for 24 h at 18 °C. The bacteria grown on the plate were collected with LB medium and spread onto an LB plate containing 30 µg/mL

chloramphenicol and 50 µg/mL rifampicin. The plates were incubated at 18 °C, and the transformants were selected.

Quantitative real-time RT-PCR analysis

Total RNA was extracted using the High Pure RNA Isolation Kit (Roche, Basel, Switzerland) from the cells cultivated in modified DSMZ medium 79 with 0.5% w/v casamino acids supplementation and with or without L-Trp supplementation. Total RNA was dissolved in 0.1% diethyl pyrocarbonate (Nacalai Tesque)-treated water and stored at –80 °C until use. Quantitative real-time RT-PCR was performed with SuperScript III Platinum SYBR Green One-Step qRT-PCR Kit (Invitrogen, Waltham, CA) and the Mx3000P Multiple Quantitative RT-PCR system (Agilent Technologies, Santa Clara, CA). The amplification curve was obtained from RT-PCR reaction mixtures containing 0.1 to 1,000 pg of the purified total RNA extract of *S. livingstonensis* Ac10. Primers listed in the “Real-time RT-PCR” section of Table S1 were used to amplify the internal region of the genes involved in tryptophan synthesis

(*trpA* to *trpE*). The reaction conditions were as follows: 95 °C for 10 min and 50 cycles of 95 °C for 30 s, 55 °C for 30 s, and 72 °C for 30 s. The threshold cycle (Ct) value for each sample was normalized with the Ct value for 16S rRNA.

BLA assay

The recombinant *S. livingstonensis* Ac10-Rif^r strains harboring each of the promoter-assay plasmids were cultivated aerobically in liquid modified DSMZ medium 79 with 0.5% w/v casamino acids supplementation containing 30 µg/mL of chloramphenicol in the presence or absence of 0.1% L-Trp at 18 °C and 4 °C. The cells were harvested by centrifugation at a temperature identical to the cultivation temperature and resuspended in 500 µL of 100 mM potassium phosphate buffer (pH 7.0). The cells were homogenized by sonication, and the soluble protein extracts were used for the BLA assay. The BLA activity was measured at 25 °C with 100 µM nitrocefin (Cayman Chemical, Ann Arbor, MI), a chromogenic substrate, in 100 mM potassium phosphate buffer (pH 7.0). The formation of the product was monitored by

measuring the increase of the absorbance at 486 nm with SPECTRA MAX 190 (Molecular Devices, San Jose, CA). The molar extinction coefficient of hydrolyzed nitrocefin is $20,500 \text{ M}^{-1} \cdot \text{cm}^{-1}$. One unit of the enzyme was defined as the amount of the enzyme that catalyzed the formation of 1 μmol of the product in 1 min.

Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and protein quantification

We performed SDS-PAGE to quantify the expression levels of BLA. Cells harvested at stationary phase ($\text{OD}_{600} = 1.8\text{-}2.0$) were applied to 12.5% polyacrylamide gel. Gels were stained with Coomassie Brilliant Blue G-250 in 10% acetic acid, and protein bands of BLA were quantified with ImageJ version 1.52o (imagej.nih.gov/ij/download.html). A standard curve of protein band intensities versus protein amounts was generated from band intensities of protein molecular weight marker, Precision Plus ProteinTM Unstained Protein Standards (Bio-Rad, Hercules, CA).

Cultivation in the presence of 3-IAA

After the cells were grown in LB medium or LB medium containing 0.1% L-Trp until the log phase ($OD_{600} = 0.7-0.8$), the cells were harvested and washed once with LB medium containing or absent of 3-IAA. After washing, the cells were transferred to fresh LB medium containing or absent of 3-IAA, which is used as a gratuitous inducer of the *trp* promoter [14], at a final concentration of 0.1%. After a 2 h incubation at 18 °C or 4 °C, the cells were harvested by centrifugation. Harvested cells were homogenized by sonication, and the crude extracts were collected by centrifugation for the BLA assay.

Results and discussion

Characterization of the *trp* operon of *S. livingstonensis* Ac10

To develop a regulatable protein production system using *S. livingstonensis* Ac10 as the host, we focused on a gene cluster (DDBJ: LC474873) containing Trp biosynthesis genes assigned as *trpA*, *trpB*, *trpC*, *trpD*, *pabA*, and *trpE* (Figure 2(a)). The amino acid sequence identities between TrpA, TrpB, TrpC, TrpD, and TrpE of *S.*

1 *livingstonensis* Ac10 and the corresponding proteins of *E. coli* were 63%, 77%, 45%,
2 52%, and 50%, respectively. Trp-biosynthesis genes are known to be included in a gene
3 cluster called the *trp* operon and transcribed together during Trp biosynthesis [15–17].
4 Expression of the *trp* operon is suppressed in the presence of excess L-Trp [18,19], and
5 this repressive regulation is found in many organisms [20–22]. Compared with the *E.*
6 *coli trp* operon, the putative Trp-biosynthesis gene cluster of *S. livingstonensis* Ac10
7 lacks a *trpL* coding for a leader peptide, which is involved in the regulation of Trp
8 synthesis by attenuation [23]. This suggests that expression of the *trp* operon of *S.*
9 *livingstonensis* Ac10 is mainly regulated by a repressor protein [18,19]. We examined
10 whether the transcriptional regulation system for the *trp* operon is applicable to
11 regulatable gene expression of recombinant protein production at low temperatures in
12 the following experiments.

14 **Analysis of the transcription level of the Trp-biosynthesis genes in *S.*** 15 ***livingstonensis* Ac10-Rif^r**

16 We investigated the effects of L-Trp supplementation on the growth of *S.*

1 *livingstonensis* Ac10-Rif^r using modified DSMZ medium 79 supplemented with 0.5%
2 w/v casamino acids as a basal medium. Both at 18 °C and 4 °C, this strain grew better in
3 the absence of L-Trp than in the presence of 0.1% L-Trp (Figure 3(a,b)). The maximum
4 OD₆₀₀ were 3.5 and 1.7, respectively, at 18 °C and 2.0 and 1.0, respectively, at 4 °C.

5 Next, to examine whether the expression of Trp-biosynthesis genes is
6 transcriptionally regulated by L-Trp supplementation, relative amounts of mRNAs of
7 Trp-biosynthesis genes were determined by real-time RT-PCR. In the presence of 0.1%
8 L-Trp, relative amounts of mRNA of *trpA*, *trpB*, *trpC*, *trpD*, and *trpE* declined to less
9 than 20% and 40% of those in the absence of L-Trp supplementation at 18 °C and 4 °C,
10 respectively (Figure 3(c,d)). These results indicated that supplementation of L-Trp
11 suppresses the expression of Trp-biosynthesis enzymes of *S. livingstonensis* Ac10, and
12 this repressive gene expression system may be applicable to regulatable heterologous
13 protein production using *S. livingstonensis* Ac10 as the host.

14
15 **Identification and characterization of a promoter of the *trp* operon of *S.***
16 ***livingstonensis* Ac10**

Using the promoter predictor (Neural Network Promoter Prediction Ver. 2.2, www.fruitfly.org/seq_tools/promoter.html), we found ten putative promoter regions from the 5'-UTR of the Trp-biosynthesis gene cluster (3,443 bp) (Figure 2(b)). The predicted promoter regions, which have conserved sequences of transcriptional regulation elements [24], were named T1, T2, T3, T4, T5, T6, T7, T8, T9, and T10. To investigate their function as promoters, we constructed ten plasmids (pT1*bla*, pT2*bla*, pT3*bla*, pT4*bla*, pT5*bla*, pT6*bla*, pT7*bla*, pT8*bla*, pT9*bla*, and pT10*bla*) harboring these predicted promoter regions upstream of the BLA-coding gene (Figure 2(b)). These plasmids were introduced into *S. livingstonensis* Ac10-Rif^r, and the transformants (named as shown in Table 1) were cultured in the basal medium with or without L-Trp supplementation at 18 °C and 4 °C to examine the expression of the BLA gene.

All the transformants showed BLA activities both at 18 °C and 4 °C (Figure 4). Higher specific activities were observed at 18 °C than at 4 °C in the absence of L-Trp. At 18 °C, the specific activities were in the range of 5.3 to 27.9 U/mg. The highest specific activity was observed for BLAT9 both at 18 °C and 4 °C. The specific activity of BLAT9 at 18 °C was more than double of that at 4 °C. The higher specific

activity at 18 °C may be partially due to a higher copy number of pJRD-Cm^r at this temperature than at 4 °C [7]. The predicted promoter T9 has a putative transcription initiation site at -1,186 position from *trpE* start codon; the site neither was empirically determined nor could be validated from sequence inspection (e.g. no apparent *trp* operator sequence in the vicinity [25]). Further experiments will be needed in the future to identify the mode of action of T9 promoter.

SDS-PAGE analysis of the proteins of BLAT9 showed that the production of the reporter protein was 75 and 33 mg/L-culture at 18 °C and 4 °C, respectively (Figure 5(a,b)), and was comparable to that achieved by using the LI3 promoter, which was previously developed for low-temperature overproduction of foreign proteins in *S. livingstonensis* Ac10 [7]. By supplementation with L-Trp, the production of this protein was significantly decreased both at 18 °C and 4 °C (Figure 5(a,b)). Real-time RT-PCR analysis demonstrated that, in the presence of L-Trp, the expression of *bla* was suppressed to about 15% and 30% of that without L-Trp supplementation at 18 °C and 4 °C, respectively (Figure 5(c)). These results suggest that the T9 promoter is useful for regulatable heterologous protein production at low temperatures.

Regulation of heterologous gene expression using the T9 promoter

We examined the validity of the T9 promoter in regulation of heterologous gene expression under various conditions. First, we examined whether the expression level can be modulated by L-Trp in a short incubation period after exchanging the culture medium. BLAT9 cells were grown in the basal medium without L-Trp supplementation until the log, late-log, and early stationary phases at 18 °C or 4 °C, and the whole cells obtained were transferred to the basal medium with or without 0.1% L-Trp supplementation. After a 2 h incubation at 18 °C or 4 °C, cell extracts were prepared, and BLA activities were measured. When the cells grown to the late-log phase were transferred to the medium supplemented with 0.1% L-Trp, both total activity and specific activity were suppressed to less than 50% of those transferred to the medium without L-Trp supplementation at 18 °C (Figure S1(a,b)). At 18 °C, a significant suppressive effect was also observed for the cells grown to the log phase. Suppression of the BLA activity by L-Trp supplementation was also observed for the cells cultivated at 4 °C (Figure S1(c,d)). Real-time RT-PCR analysis of *bla* mRNA in the cells

1 incubated in the presence or absence of L-Trp supplementation demonstrated that
2 transcription of the gene was significantly decreased in the presence of L-Trp (Figure
3 S1(e)), indicating that production of BLA is regulated at the transcriptional level. Thus,
4 expression of genes under the control of the T9 promoter can be suppressed by
5 incubation with L-Trp for a short period at low temperatures after exchanging the
6 culture medium. Since BLA was detected as one of the major protein bands in the cells
7 incubated without L-Trp supplementation (Figure S1(f)), the T9 promoter can be
8 considered as a powerful and regulatable promoter useful for production of
9 heterologous proteins in *S. livingstonensis* Ac10.

10 We also performed a similar experiment by using the BLAT9 cells
11 pre-cultivated in the basal medium supplemented with L-Trp to suppress expression of
12 the heterologous protein. The cells grown to the late-log phase at 18 °C or 4 °C were
13 transferred to the basal medium with or without 0.1% L-Trp supplementation and
14 incubated for 2 h at the respective temperature. In this case, the total activity of BLA
15 was about 3.7 and 1.9 times higher when the cells were incubated in the absence of
16 L-Trp than in the presence of L-Trp at 18 °C and 4 °C, respectively (Figure S2(a,c)).

1 The specific activity was about 3.2 and 2.4 times higher when the cells were incubated
2 in the absence of L-Trp at 18 °C and 4 °C, respectively (Figure S2(b,d)). Induction of
3 expression of heterologous proteins by incubation in the absence of L-Trp after
4 obtaining a large amount of the cells by pre-cultivation in the presence of L-Trp would
5 be useful for production of toxic proteins of which production inhibits the growth of the
6 host cells.

7 We next examined the effects of 3-IAA, which is used as a gratuitous inducer
8 of the *trp* promoter of *E. coli* [14], on expression of genes under the control of the T9
9 promoter. For this experiment, LB medium was used instead of the synthetic minimal
10 medium. The cells grown in LB medium to the log ($OD_{600} = 0.7-0.8$), late-log ($OD_{600} =$
11 $0.8-1.0$), and stationary ($OD_{600} = 1.8-2.0$) phases at 18 °C and 4 °C were transferred to
12 LB medium containing 0.1% 3-IAA and incubated for 2 h (Figure 6(a-d)). As a control,
13 cells were transferred to LB medium without 3-IAA. The highest total activity and
14 specific activity were observed when the cells grown to log phase were transferred to
15 the 3-IAA-containing medium both at 18 °C and 4 °C. The total activity in the
16 3-IAA-containing medium was 3.2 and 3.3 times higher than that in the medium

1 without 3-IAA at 18 °C and 4 °C, respectively. The specific activity in the
2 3-IAA-containing medium was 2.6 and 3.8 times higher than that in the medium
3 without 3-IAA at 18 °C and 4 °C, respectively. Thus, 3-IAA is expected to be useful for
4 inducible expression of heterologous genes under the control of the T9 promoter.

5 Through this study, we succeeded in applying a promoter of the
6 Trp-biosynthesis operon, named the T9 promoter, of *S. livingstonensis* Ac10 to
7 regulatable gene expression at low temperatures. In the absence of L-Trp, 75 mg/L and
8 33 mg/L cultures of BLA were produced as one of the major proteins at 18 °C and 4 °C,
9 respectively. These yields are comparable to those obtained using the LI3 promoter,
10 which was previously developed for overproduction of foreign proteins in *S.*
11 *livingstonensis* Ac10 [7]. We also demonstrated that gene expression under the control
12 of the T9 promoter can be regulated by 3-IAA supplementation. Thus, the T9 promoter
13 is expected to be useful for the production of various proteins including toxic proteins
14 whose production should be suppressed during pre-cultivation to obtain a large enough
15 number of cells for their production. Finding of the T9 promoter as a new regulatable
16 promoter could expand the utility of the low-temperature protein expression systems

with *S. livingstonensis* Ac10 as the host.

Author Contribution

S.K., J.K., and T.K. designed the study. S.K. performed the experiments. S.K., J.K., and T.K. discussed the results. S.K., J.K., T.O., and T.K. wrote the paper.

Acknowledgements

This work was supported in part by JSPS KAKENHI (17H04598 and 18K19178 to T.K. and 16K14885 to J.K.).

References

- [1] Sørensen HP, Mortensen KK. Soluble expression of recombinant proteins in the cytoplasm of *Escherichia coli*. Microb Cell Fact. 2005;4:1–8.
- [2] Rasala BA, Mayfield SP. Photosynthetic biomanufacturing in green algae; Production of recombinant proteins for industrial, nutritional, and medical uses. Photosynth Res. 2015;123(3):227–239.
- [3] Ritacco FV, Wu Y, Khetan A. Cell culture media for recombinant protein expression in Chinese hamster ovary (CHO) cells: History, key components, and optimization strategies. Biotechnol Prog. 2018;34(6):1407–1426.
- [4] Gerday C, Aittaleb M, Bentahir M, et al. Cold-adapted enzymes: from fundamentals to biotechnology. Trends Biotechnol. 2000;18(3):103–107.
- [5] Cronan JE. Improved plasmid-based system for fully regulated off-to-on gene expression in *Escherichia coli*: Application to production of toxic proteins. Plasmid. 2013;69(1):81–89.
- [6] Mujacic M, Cooper KW, Baneyx F. Cold-inducible cloning vectors for low-temperature protein expression in *Escherichia coli*: application to the

- 1 production of a toxic and proteolytically sensitive fusion protein. *Gene*.
2 1999;238(2):325-332.
- 3 [7] Miyake R, Kawamoto J, Wei YL, et al. Construction of a low-temperature
4 protein expression system using a cold-adapted bacterium, *Shewanella* sp. strain
5 Ac10, as the host. *Appl Environ Microbiol*. 2007;73(15):4849–4856.
- 6 [8] Sannino F, Giuliani M, Salvatore, U, et al. A novel synthetic medium and
7 expression system for subzero growth and recombinant protein production in
8 *Pseudoalteromonas haloplanktis* TAC125. *Appl Microbiol Biotechnol*.
9 2017;101(2):725-734.
- 10 [9] Kawamoto J, Kurihara T, Kitagawa M, et al. Proteomic studies of an Antarctic
11 cold-adapted bacterium, *Shewanella livingstonensis* Ac10, for global
12 identification of cold-inducible proteins. *Extremophiles*. 2007;11:819–826.
- 13 [10] Rabus R, Ruepp A, Frickey T, et al. The genome of *Desulfotalea psychrophila*, a
14 sulfate-reducing bacterium from permanently cold Arctic sediments. *Environ*
15 *Microbiol*. 2004;6(9):887–902.
- 16 [11] Papa R, Glagla S, Danchin A, et al. Proteomic identification of a two-component

- 1 regulatory system in *Pseudoalteromonas haloplanktis* TAC125. *Extremophiles*.
2 2006;10(6):483–491.
- 3 [12] Davison J, Heusterspreute M, Chevalier N, et al. Vectors with restriction site
4 banks. V. pJRD215, a wide-host-range cosmid vector with multiple cloning sites.
5 *Gene*. 1987;51(2,3):275–280.
- 6 [13] Toyotake Y, Cho HN, Kawamoto J, et al. A novel
7 1-acyl-*sn*-glycerol-3-phosphate *O*-acyltransferase homolog for the synthesis of
8 membrane phospholipids with a branched-chain fatty acyl group in *Shewanella*
9 *livingstonensis* Ac10. *Biochem Biophys Res Commun*. 2018;500(3):704–709.
- 10 [14] Somerville RL. The *trp* promoter of *Escherichia coli* and its exploitation in the
11 design of efficient protein production systems. *Biotechnol Genet Eng Rev*.
12 1988;6:1–41.
- 13 [15] Liubetskaia EV, Leont’ev LA, Gel’fand MS, et al. Search for alternative
14 secondary structures of RNA, regulating expression of bacterial genes. *Mol Biol*.
15 2003;37(5):834–842.
- 16 [16] Das A, Crawford IP, Yanofsky C. Regulation of tryptophan operon expression by

1 attenuation in cell-free extracts of *Escherichia coli*. J Biol Chem.

2 1982;257(15):8795–8798.

3 [17] Kelley RL, Yanofsky C. Trp aporepressor production is controlled by autogenous

4 regulation and inefficient translation. Proc Natl Acad Sci U S A.

5 1982;79(10):3120–3124.

6 [18] Jeeves M, Evans PD, Parslow RA, et al. Studies of the *Escherichia coli* Trp

7 repressor binding to its five operators and to variant operator sequences. Eur J

8 Biochem. 1999;265(3):919–928.

9 [19] Arvidson DN, Grove Arvidson C, Lawson CL, et al. The tryptophan repressor

10 sequence is highly conserved among the *Enterobacteriaceae*. Nucleic Acids Res.

11 1994;22(10):1821–1829.

12 [20] Palego L, Betti L, Rossi A, et al. Tryptophan biochemistry: Structural, nutritional,

13 metabolic, and medical aspects in humans. J Amino Acids. 2016;2016:1-13.

14 [21] Abbas SH, Sohail M, Saleem M, et al. Effect of L-tryptophan on plant weight and

15 pod weight in chickpea under rainfed conditions. Sci Tech Dev. 2013;32(4):277–

16 280.

- 1 [22] Xie Y, Reeve JN. Regulation of tryptophan operon expression in the archaeon
2 *Methanothermobacter thermautotrophicus*. J Bacteriol. 2005;187(18):6419–
3 6429.
- 4 [23] Bertrand K, Yanofsky C. Regulation of transcription termination in the leader
5 region of the tryptophan operon of *Escherichia coli* involves tryptophan or its
6 metabolic product. J Mol Biol. 1976;103(2):339–349.
- 7 [24] Hook-Barnard IG, Hinton DM. Transcription initiation by mix and match
8 elements: flexibility for polymerase binding to bacterial promoters. Gene Regul
9 Syst Biol. 2007(1);275–293.
- 10 [25] Mironov AA, Koonin EV, Roytberg MA, et al. Computer analysis of
11 transcription regulatory patterns in completely sequenced bacterial genomes.
12 Nucleic Acids Res. 1999;27(14):2981-2989.
- 13 [26] Hanahan D, Jessee J, Bloom FR. Plasmid transformation of *Escherichia coli* and
14 other bacteria. Methods Enzymol. 1991;204:63–113.
- 15 [27] Simon R, Priefer U, Pühler A, et al. A broad host range mobilization system for
16 *in vivo* genetic engineering: transposon mutagenesis in gram negative bacteria.

1 Bio Technol. 1983;1:784–791.

2

3

4

1 Table 1. Strains and plasmids used in this study

Strains	Descriptions	Reference
<i>Escherichia coli</i>		
DH5α	Host for plasmids constructions	[26]
S17-1/λ _{pir}	S17-1 derivative, host for <i>pir</i> -dependent plasmids	[27]
<i>Shewanella livingstonensis</i> Ac10		
Ac10-Rif ^r	Parent strain, rifampicin-resistant mutant of Ac10	[7]
BLAT0	Ac10-Rif ^r harboring <i>pbla</i>	This study
BLAT1	Ac10-Rif ^r harboring pT1 <i>bla</i>	This study
BLAT2	Ac10-Rif ^r harboring pT2 <i>bla</i>	This study
BLAT3	Ac10-Rif ^r harboring pT3 <i>bla</i>	This study
BLAT4	Ac10-Rif ^r harboring pT4 <i>bla</i>	This study
BLAT5	Ac10-Rif ^r harboring pT5 <i>bla</i>	This study
BLAT6	Ac10-Rif ^r harboring pT6 <i>bla</i>	This study
BLAT7	Ac10-Rif ^r harboring pT7 <i>bla</i>	This study
BLAT8	Ac10-Rif ^r harboring pT8 <i>bla</i>	This study
BLAT9	Ac10-Rif ^r harboring pT9 <i>bla</i>	This study
BLAT10	Ac10-Rif ^r harboring pT10 <i>bla</i>	This study
Plasmids		
pJRD-Cm ^r	A broad-host-range vector	[13]
<i>pbla</i>	pJRD-Cm ^r containing <i>bla</i> from pBR322 without promoter sequence	This study
pT1 <i>bla</i>	pJRD-Cm ^r containing T1 and <i>bla</i>	This study
pT2 <i>bla</i>	pJRD-Cm ^r containing T2 and <i>bla</i>	This study
pT3 <i>bla</i>	pJRD-Cm ^r containing T3 and <i>bla</i>	This study
pT4 <i>bla</i>	pJRD-Cm ^r containing T4 and <i>bla</i>	This study
pT5 <i>bla</i>	pJRD-Cm ^r containing T5 and <i>bla</i>	This study
pT6 <i>bla</i>	pJRD-Cm ^r containing T6 and <i>bla</i>	This study
pT7 <i>bla</i>	pJRD-Cm ^r containing T7 and <i>bla</i>	This study
pT8 <i>bla</i>	pJRD-Cm ^r containing T8 and <i>bla</i>	This study
pT9 <i>bla</i>	pJRD-Cm ^r containing T9 and <i>bla</i>	This study
pT10 <i>bla</i>	pJRD-Cm ^r containing T10 and <i>bla</i>	This study

2

3

1 **Figure legends**

2 **Figure 1. Construction of plasmids for the promoter assay.**

3 *rep*: replication-related gene, *mob*: mobilization-related gene, *oriV*: replication origin,

4 *Cm^r*: chloramphenicol resistance gene, *MluI* and *SpeI*: restriction enzyme sites

5

6 **Figure 2. The *trp* operon of *S. livingstonensis* Ac10 and the 5'-untranslated region**

7 **(5'-UTR) of *trpE* tested for its promoter activity.**

8 (a) Genetic region containing the *trp* operon of *S. livingstonensis* Ac10. The scale bar

9 represents 1 kb. (b) Schematic illustration of 5'-UTR of *trpE* tested for its promoter

10 activity by using the BLA gene as a reporter. DNA fragments of the 5'-UTR of *trpE*

11 (T1-T10) were fused with the BLA-coding gene and introduced into the promoter-assay

12 plasmid as shown in Figure 1. Putative promoter sequences were predicted by Neural

13 Network Promoter Prediction (ver.2.2)

14 (http://www.fruitfly.org/seq_tools/promoter.html), and the locations of the sequences

15 with a prediction score over 0.95 are indicated with triangles.

16

Figure 3. Effects of L-Trp supplementation on the growth of *S. livingstonensis*

Ac10-Rif^r and expression of its Trp-biosynthesis genes.

(a, b) Growth curves of *S. livingstonensis* Ac10-Rif^r. The cells were grown at 18 °C (a) and 4 °C (b) in modified DSMZ medium 79 with 0.5% w/v casamino acids supplementation containing (■) and absent of L-Trp (□) at a final concentration of 0.1%.

The statistical analysis was performed using a Student's *t*-test from three independent experiments. Error bars indicate SD. An asterisk (*) indicates a statistically significant

difference (Student's *t*-test, $P < 0.01$). (c, d) Transcription levels of the Trp-biosynthesis

genes. The ratios of the amounts of mRNA of the genes in the *trp* operon in the cells

grown with or without 0.1% L-Trp are shown. The cells were grown at 18 °C (c) or 4 °C

(d) and harvested in the log phase. All values were normalized to the amounts of 16S

rRNA. Error bars represent SD from three independent experiments.

Figure 4. Production of BLA by recombinant *S. livingstonensis* Ac10-Rif^r

harboring various promoter-assay plasmids.

Specific activities of BLA produced by recombinant *S. livingstonensis* Ac10-Rif^r cells

(BLAT 1 to BLAT10) harboring the promoter-assay plasmids, pT1*bla* to pT10*bla*, respectively, are shown. BLAT0 is a negative control strain harboring *pbla*, which contains the BLA gene without a promoter sequence. The cells were grown at 18 °C (a) and 4 °C (b) in the presence (open bars) or absence (closed bars) of 0.1% L-Trp and harvested in the stationary phase ($OD_{600} = 1.8-2.0$). Error bars represent SD from three independent experiments. An asterisk (*) indicates a statistically significant difference (Student's *t*-test, $P < 0.01$).

Figure 5. Expression of *bla* in recombinant *S. livingstonensis* Ac10-Rif^r harboring pT9*bla* (BLAT9).

(a, b) SDS-PAGE analysis of proteins of BLAT9 grown with or without L-Trp at 18 °C (a) and 4 °C (b). Arrowheads indicate the position of the BLA protein band. M: protein marker; Cont.: BLAT0 grown without L-Trp. (c, d) Relative amounts of *bla* mRNA in BLAT9 grown with 0.1% L-Trp in comparison with that in the cells grown without L-Trp. The cells were grown at 18 °C (c) and 4 °C (d). All values were normalized to the amounts of 16S rRNA. Error bars represent SD from three independent experiments.

Figure 6. Effects of 3-indoleacrylic acid (3-IAA) on expression of the gene under the control of the T9 promoter.

Total activity (a and c) and specific activity (b and d) of BLA from a 5 mL culture of BLAT9. The cells grown at 18 °C (a and b) and 4 °C (c and d) to the log ($OD_{600} = 0.7-0.8$), late-log ($OD_{600} = 0.8-1.0$), and stationary ($OD_{600} = 1.8-2.0$) phases were transferred to LB medium with (open bars) or without (closed bars) 0.1% 3-IAA supplementation. As a control, BLAT0 harboring *pbla*, which contains the BLA gene without a promoter, was used. Error bars represent SD values from three independent experiments. Asterisks (*) indicate statistically significant differences (Student's *t*-test, $P < 0.01$).

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6





Table S1. Primers used in this study

Primer	Sequence
Real-time RT-PCR	
16S-Fw	5'-AGAGTTTGATCMTGGCTCAG-3'
16S-Rv	5'-TACGGYTACCTTGTTACGACTT-3'
TrpA-Fw	5'-CATTAGCGGCAGGTACAAC-3'
TrpA-Rv	5'-AGATCGGTGCAATACCATGT-3'
TrpB-Fw	5'-TCCGCTGGTGAAAATCTACC-3'
TrpB-Rv	5'-GCAAAAGACATTGAGCGTCA-3'
TrpC-Fw	5'-ACAGATCCCAGCAGACAGAG-3'
TrpC-Rv	5'-CAAGCCGCAGACTTTATTGT-3'
TrpD-Fw	5'-GCGTATCAAGCAAGTCTGGT-3'
TrpD-Rv	5'-GGTACTGCGTGTTTCACTCC-3'
TrpE-Fw	5'-GTGTTACCGGTCAATTACGC-3'
TrpE-Rv	5'-AGCTGAACGAATAACGATGC-3'
BLA-Fw	5'-GCCTTCCTGTTTTGCTCAC-3'
BLA-Rv	5'-ATAATACCGCGCCACATAGC-3'
Amplification of pJRD-Cm	
pJRD-Fw	5'-CGTAATCCATGGATCAAGAG-3'
pJRD-Rv	5'-TAGTATAGTCTATAGTCCGTGG-3'
Amplification of predicted promoter regions	
Trp1-Fw	5'-CCTGTCTCTTGATCCATGGATTACGATTGTGCGTTGTATCACTGC-3'
Trp2-Fw	5'-CCTGTCTCTTGATCCATGGTTACGAATATGACTAATCGCAAC-3'
Trp3-Fw	5'-CCTGTCTCTTGATCCATGGATTACGTCTTGAGTTTGGAACGATAG-3'
Trp4-Fw	5'-CCTGTCTCTTGATCCATGGATTACGGATTTTAACTAACTGTACGCC-3'
Trp5-Fw	5'-CCTGTCTCTTGATCCATGGATTACGTGATTTGTTGAATAGGCAAATC-3'
Trp6-Fw	5'-TTGATCCATGGATTACGAATTTTTGTTTACGGATTAAATATAACAATAAATC-3'
Trp7-Fw	5'-CCTGTCTCTTGATCCATGGATTACGGTTTGATTTTCAGGTAAATAACC-3'
Trp8-Fw	5'-CCTGTCTCTTGATCCATGGATTACGATTTTTTTCACCAATCATACAGC-3'
Trp9-Fw	5'-CCTGTCTCTTGATCCATGGTTACGTGGTTGGAATAAGTTTTTACC-3'
Trp10-Fw	5'-TTGATCCATGGATTACGGTTATTTGCTTACTTAATGCAATAA-3'
Trp-Rv.1	5'-GTTGAATACTCATCTGTTGGCCTTTAATATTATGTAACG-3'
Trp-Rv.2	5'-GAATACTCATCTGTTGGCCTTTAATATTATGTAAC-3'
Amplification of BLA-coding gene	
β -lac-Fw.1	5'-TAAAGGCCAACAGATGAGTATTCAACATTTCCGTG-3'
β -lac-Rv.1	5'-ATTCCACCGACTATAGACTATACTATACTATTACCAATGCTTAATCAGTGA-3'
β -lac-Fw.2	5'-ATTCCACCGACTATAGACTATACTATTACCAATGCTTAATCAGTGAG-3'
β -lac-Rv.2	5'-GGACTATAGACTATTACCAATTGCTTAATCAGTGA-3'

Figure S1






Figure S1. Effects of L-Trp on the T9 promoter-controlled gene expression in the cells pre-cultivated in the absence of L-Trp.

(a-d) Total activity of 5-mL culture (a, c) and specific activity (b, d) of BLA from BLAT9 incubated in the presence (open bars) or absence (closed bars) of L-Trp for 2 h after pre-cultivation in the absence of L-Trp to the log ($OD_{600} = 0.7-0.8$), late-log ($OD_{600} = 0.8-1.0$), and early stationary ($OD_{600} = 1.8-2.0$) phases at 18 °C (a, b) and 4 °C (c, d). Asterisk (*) indicates a statistically significant difference ($P < 0.01$). (e) The amount of *bla* mRNA in the cells incubated in the absence or presence of L-Trp for 2 h after growth to the late-log phase at 18 °C were quantified by real-time RT-PCR. Error bars represent SD values from three independent experiments. (f) The protein composition of the BLAT9 cells was analyzed by SDS-PAGE. After the cells were grown to log phase in the basal medium without L-Trp supplementation, the cells were incubated in the presence (+) or absence (-) of 0.1% L-Trp for 2 h at 18 °C. Arrowheads indicate the position of the BLA protein band. M: protein marker; Cont.: BLAT0, harboring a control plasmid, *pbla*. After BLAT0 was grown to the late-log phase, the cells were transferred to the basal medium without L-Trp supplementation and incubated for 2 h.

Figure S2




Figure S2. Effects of L-Trp on the T9 promoter-controlled gene expression in cells pre-cultivated in the presence of L-Trp.

Total activity of 5-mL culture (a, c) and specific activity (b, d) of BLA from BLAT9 incubated in the presence (open bars) or absence (closed bars) of L-Trp for 2 h after pre-cultivation in the presence of 0.1% L-Trp to the late-log ($OD_{600} = 0.8-1.0$) phase at 18 °C (a, b) and 4 °C (c, d). Error bars represent SD values from three independent experiments. Asterisks (*) indicate statistically significant differences ($P < 0.01$).