

Original Article

Synthesis of the ABCDG ring skeleton of communesin F based on carboborylation of 1,3-diene and Bi(OTf)₃-catalyzed cyclizations

Motoyuki Nakajima, Chihiro Tsukano,* Motohiro Yasui, and Yoshiji Takemoto*

Graduate School of Pharmaceutical Sciences, Kyoto University, Yoshida, Sakyo-ku, Kyoto 606-8501, Japan

ABSTRACT: Communesins, isolated from the mycelium of a strain of *Penicillium* sp., are cytotoxic heptacyclic indole alkaloids bearing a bis-aminal structure and two contiguous quaternary carbon centers. Towards a total synthesis of communesin F, we synthesized a pentacyclic ABCDG ring skeleton via carboborylation of 1,3-diene and a Friedel-Crafts-type cyclization, resulting in the formation of an azepine ring through a Bi(OTf)₃-catalyzed S_N2' reaction.

Keywords: communesin/ carboborylation/ amidine/ Bi(OTf)₃

Introduction

Communesins A and B, which were originally isolated by Numata and co-workers from the mycelium of a strain of *Penicillium* sp. attached to the marine alga *Enteromorpha intestinalis*, are heptacyclic indole alkaloids (Figure 1).¹ Spectroscopic analyses, including nuclear magnetic resonance (NMR) spectroscopy (¹H NMR, ¹³C NMR including 2D NMR) and high-resolution mass spectrometry, have revealed that their structures are quite unique. They are characterized by a heptacyclic skeleton bearing two aminals and two contiguous quaternary carbon centers. To date, nine congeners have been reported,²⁻⁶ and perophoramidine is also known as a structurally related bis-amidine indole alkaloid.⁷ Recently, Tang and co-workers confirmed that communesins can be biosynthetically produced through the coupling of aurantioclavine and tryptamine based on genetic inactivation studies.⁸ Communesins show cytotoxicity against P388 lymphocytic leukemia cells (ED₅₀ A: 3.5 µg/mL, B: 0.45 µg/mL) and potent insecticidal activity towards silkworms (LD₅₀ D: 300 µg/g, E: 80 µg/g). Because of their unique structure and biological activity, many research groups have conducted synthetic studies of communesins in which various synthetic methods were developed.⁹⁻¹⁴ The first racemic total synthesis of communesin F was achieved by Qin and co-workers based on an intramolecular cyclopropanation strategy.¹⁵ Weinreb and Funk also reported total synthesis of communesin F, independently.^{16,17} The first asymmetric total syntheses of communesins A, B and F were accomplished by Ma and co-workers.^{18,19} Asymmetric total syntheses were also reported by Stoltz, Movassaghi, Yang, and Chen, independently.²⁰⁻²³ We have also engaged in the development of synthetic strategies for this class of alkaloids including communesins, perophoramidine and aurantioclavine.²⁴⁻³¹





Figure 1. Communesins and related alkaloids.


Results and Discussion

Recently, we have developed palladium(Pd)-catalyzed carbosilylation of 1,3-diene with carbamoyl chloride for the synthesis of several spirooxindoles.³² Extending this reaction, a Pd-catalyzed carboborylation of 1,3-diene was developed for a synthesis of iminoindoline.³⁰ Considering our developed method, it was envisioned that communesin F would be accessed from a pentacyclic skeleton **II** through intermediate **I** by the introduction of an aminoethyl unit and the formation of amidine. The pentacyclic skeleton **II** would be constructed from a tetracyclic compound **IV** via **III** by the introduction of an allyl alcohol unit, resulting in an S_N2' reaction for the formation of an azepine ring and a reduction of amidine. The tetracyclic compound **IV** can be synthesized by a carboborylation of 1,3-diene **VI** and an intramolecular Friedel-Crafts-type reaction of a resultant iminoindoline **V**.³⁰ Following this retrosynthetic analysis, we have recently succeeded in the construction of tetracyclic skeleton **IV** (R = OMe) from diene **VI** (R = OMe) through iminoindoline **V** (R = OMe). However, compound **1** could not be converted to compound **2** through removal of the methyl group, although we tried various conditions including BBr₃, BCl₃, AlCl₃, LiCl, Ph₂PLi and *p*MeC₆H₄SLi (Scheme 1b).³³ These reaction conditions resulted in the removal of a Cbz group or the decomposition of compound **1**. Therefore, we needed to revise our initial synthetic route and planned to employ a 1,3-diene-containing triflate (R = OTf) to avoid a protecting group manipulation. The use of a substrate bearing a triflate group for Pd-catalyzed carboborylation would extend its reaction scope, and it might react itself under the reaction conditions. In the current manuscript, we report the construction of a pentacyclic skeleton of communesin F by extending our strategy based on carboborylation of 1,3-diene.

(a) Retrosynthesis of communesin F



(b)




Scheme 1. (a) Retrosynthesis of communesin F and (b) failed attempt at removing a methyl group from compound 1.

The synthesis started with a removal of a methyl group on a phenolic hydroxyl group. A methoxy aniline derivative **3**, which was prepared from *t*-butyl(3-methoxyphenyl)carbamate in four steps,³⁰ was treated using BBr₃ to give a phenol (Scheme 2). The resultant phenolic hydroxy group was silylated with *tert*-butyldimethylsilyl (TBS) chloride and imidazole to give compound **4**. A half reduction of a lactone with diisobutylaluminum hydride (DIBAL-H) was followed by Wittig olefination, which gave diene **5** through internal transfer of a TBS group. After the formation of urea by a treatment of phenyl isocyanate, a phenolic hydroxy group was protected as a triflate. A removal of a TBS group was followed by a Mitsunobu reaction with *p*NsNHBOc³⁴ to give compound **8**, which was converted to carbodiimide **9** through dehydration with CBr₄, PPh₃ and Et₃N.

With carbodiimide **9** containing a diene moiety, we investigated whether the triflate is intact under the reaction conditions of Pd-catalyzed carboborylation of 1,3-diene. In the previous literature, there is no report concerning


69 Pd(II)-catalyzed Miyaura borylation of triflates and diborane without a ligand, but reactions using
 70 diphenylphosphinoferrocene³⁵ or the reaction of arylbromide have been reported.³⁶ Therefore, it was expected that a
 71 triflate group would be intact during the carboborylation of 1,3-diene. As expected, the reaction of **9** proceeded
 72 smoothly under the established conditions (Pd(OAc)₂, (pinB)₂, xylene, 50°C) to give an allyl borane, which was
 73 treated with NaBO₃·4H₂O to give allyl alcohol **10**. After silylation of allyl alcohol **10**, a *tert*-butoxyoxycarbonyl (Boc)
 74 group was introduced to an amidine nitrogen for further transformation. The treatment of compound **11** using
 75 tetrabutylammonium fluoride gave an allyl alcohol along with the removal of a triflate group, which was converted
 76 to allyl bromide **12** under standard conditions. Unfortunately, the resultant allyl bromide **12** could not be converted
 77 to compound **13** through a treatment of Tf₂O and pyridine. On the other hand, when HF·pyridine was used, a
 78 triethylsilyl group was selectively removed with the triflate group intact. The resultant allyl alcohol was also
 79 converted to allyl bromide **13** containing a triflate group, while a small amount of compound **14** was also obtained
 80 through the removal of a Boc group.



81
 82 Scheme 2. Synthesis of 3,3-disubstituted iminoindoline **10** based on the Pd-catalyzed carboborylation of 1,3-diene

and its derivatization.

Next, we investigated Friedel-Crafts-type cyclization of allyl bromides **12** and **13** to construct a tetracyclic ABCD ring skeleton. Previously, we have reported the cyclization of compound **15** containing a methoxy group using 10 mol% of Bi(OTf)₃ and 3.5 equivalents of AgOTf (Table 1, entry 1).^{30,37–39} The reaction gave compound **16a** in 49% yield along with **16b** in 30% yield. We initially applied these conditions to a cyclization of compound **13** containing a triflate group. However, the reaction gave a complex mixture instead of any cyclized products **17a** and **17b** (entry 2). On the other hand, the cyclization of compound **12** containing a phenolic hydroxy group proceeded under the same conditions to give compounds **18a** and **18b** in 63% and 30% yields with excellent stereochemistry, respectively (entry 3). The stereochemistry was determined by a comparison with our previous results³⁰ and a NOESY experiment of a derivatized compound **28** (Scheme 4, *vide infra*). When 3.5 equivalents of AgOTf was reduced to 1.2 equivalents, the formation of byproduct **18b** was suppressed to 17% yield (entry 4). Finally, the yield of the desired product **18a** was improved to 80% yield using 1.05 equivalents of AgOTf (entry 5). AgOTf was essential for this Friedel-Crafts-type reaction (entry 6)




entry	starting material	X equiv	product	
1	15 (R = OMe)	3.5	16a : 49%	16b : 30%
2	13 (R = OTf)	3.5	17a : 0%	17b : 0%*
3	12 (R = OH)	3.5	18a : 63%	18b : 30%
4	12 (R = OH)	1.2	18a : 74%	18b : 17%
5	12 (R = OH)	1.05	18a : 80%	18b : 13%
6	12 (R = OH)	0	18a : 0%	18b : 0%**

*complex mixture. ** Starting material **12** was recovered in 77% yield.

Table 1. Formation of a tetracyclic ABCD skeleton through a Friedel-Crafts-type reaction.

After the construction of a tetracyclic ABCD ring skeleton containing an amidine, we turned our attention to the

formation of an azepine ring (G ring). A treatment of compound **18a** with TiF_2O and pyridine gave compound **17a** in 91% yield (Scheme 3). To introduce an allyl alcohol unit, Suzuki-Miyaura coupling with vinyl boronic ester **19** was examined. When compound **17a** and vinyl boronic ester **19** were treated with a catalytic amount of $\text{Pd}(\text{dba})_2$, SPhos and K_3PO_4 , or $\text{Pd}(\text{PPh}_3)_4$ and Na_2CO_3 in *N,N*-dimethylformamide (DMF) at 100°C , respectively, these reactions gave the desired product **20** in low yields (Table 2, entries 1 and 2). However, conditions involving $\text{Pd}(\text{PPh}_3)_4$ and Na_2CO_3 in toluene and ethanol at 100°C improved the yield to 56% (entry 3). The removal of the *p*Ns and trimethylsilyl (TMS) group gave allyl alcohol **21** in 66% yield over two steps. To construct the azepine ring, mesylation of a tertiary alcohol was initially attempted through a treatment using methanesulfonyl chloride (MsCl) and Et_3N .¹⁸ However, a dehydration occurred to give diene **23** instead of the desired cyclized product **22**. Interestingly, when compound **21** was treated with pyridinium *p*-toluenesulfonate (PPTS),¹⁵ ortho-amide **24** was observed (as assessed using ^1H NMR analysis). A related structure was observed in synthetic studies of dehaloperophoramidine reported by Somfai and co-workers.^{13,14} We considered the thermodynamic stability of possible equilibrium products such as simplified compounds **25**, **26** and **27** through density functional theory (DFT) calculations (Figure 2). These calculations revealed that ortho-amide **26** was the most stable isomer among these compounds. These results indicate that the formation of the ortho-amide through acid activation using PPTS from amidine would be a competitive process with the formation of the azepine ring via the $\text{S}_{\text{N}}2$ reaction of the tertiary alcohol, and the equilibrium tends to be biased towards the ortho-amides such as compounds **24** and **26**. Therefore, we expected that it would be difficult to achieve the formation of azepine **22** from compound **21** containing the amidine moiety.



Scheme 3. Failed attempt at the formation of an azepine ring.

Table 2. Suzuki-Miyaura coupling of compound **17a** and boronic acid **19**.

entry	cat.	ligand	base	solvent	temp.	yield
1	Pd(dba) ₂	SPhos	K ₃ PO ₄	DMF	100 °C	10%
2	Pd(PPh ₃) ₄	—	aq. Na ₂ CO ₃	DMF	100 °C	28%
3	Pd(PPh ₃) ₄	—	aq. Na ₂ CO ₃	toluene/EtOH	100 °C	56%





Figure 2. Comparison of the thermodynamic stability of formable compounds **25**, **26** and **27**, calculated using Gaussian '09 at the B3LYP/6-31G(d) level of theory (DFT).

Therefore, a reduction of amidine **20** was investigated prior to the formation of the azepine ring to avoid the formation of the ortho-amide (Scheme 4). When compound **20** was treated using NaBH₄, the desired product was not obtained. In the case of DIBAL-H, the removal of a Boc group occurred instead of the reduction of the amidine. However, in sharp contrast, treatment using catecholborane⁴⁰ gave the desired product **28** in 65% yield as a 3.3:1 mixture of diastereomers. A NOESY experiment indicated that the stereochemistry of the major isomer was a *trans*-fused structure, which would be epimerized to a *cis*-fused structure later. Because a reducing reagent approached from the less hindered face, the *trans* isomer was obtained as a major product in this reaction. After the removal of Boc and the TMS groups, the formation of an azepine ring was investigated again. When compound **29** was treated using MsCl and Et₃N,¹⁸ the reaction gave diene **31** in 48% yield and the desired cyclized product **30** was not detected at all (Table 3, entry 1). When Bi(OTf)₃ was employed at –15°C as a Lewis acid, the reaction proceeded to give the desired product **30** as a major product albeit in low yield (entry 2).^{41,42} The reaction using Bi(OTf)₃ at –40°C gave the desired product **30** in 17% yield with recovery of the starting material (entry 3). However, under room temperature reaction

conditions the starting material **29** was consumed completely to give the desired azepine **30** in 55% yield, while diene **31** was obtained in 34% yield (entry 4). The obtained pentacyclic compound **30** would be useful for further derivatization, and now we are investigating further transformations to achieve a total synthesis of communesin F.



Scheme 4. Synthesis of the ABCDG ring skeleton **30**.

Table 3. Investigation of the formation of the azepine ring.


entry	conditions	yields
1	MsCl, Et ₃ N, CH ₂ Cl ₂ , 0 °C	30 : 0%, 31 : 48%
2	Bi(OTf) ₃ (10 mol%), MS4A CH ₂ Cl ₂ , -15 °C	30 : 23%, 31 : trace
3	Bi(OTf) ₃ (10 mol%), MS4A CH ₂ Cl ₂ , -40 °C	30 : 17%, 31 : 0%, starting material 29 67%
4	Bi(OTf) ₃ (10 mol%), MS4A CH ₂ Cl ₂ , 0 °C to rt	30 : 55%, 31 : 34%

In summary, we have investigated the synthesis of a pentacyclic ABCDG ring skeleton of communesin F based on carboborylation of 1,3-diene, a Bi(OTf)₃-catalyzed Friedel-Crafts-type reaction and azepine ring formation. It is interesting that a triflate group was intact under the conditions required for Pd-catalyzed carboborylation of 1,3-diene. Additionally, it was essential that the resultant amidine was reduced prior to the formation of the azepine ring through Bi(OTf)₃-catalyzed cyclization to avoid an undesired formation of ortho-amide. We are currently investigating further transformation of the pentacyclic compound to complete the synthesis of communesin F.

Experimental Procedure


General. All non-aqueous reactions were carried out under a positive pressure of argon in over-dried glassware. Analytical thin-layer chromatography was performed using Silica gel 60 plates (Merck, Darmstadt, Germany). Silica gel column chromatography was performed using Kanto silica gel 60 (particle size 63–210 μm , Kanto, Tokyo, Japan) and Chromatorex BW-300 (Fuji silysia, Aichi, Japan). Proton nuclear magnetic resonance (^1H NMR) spectra were recorded using a JNM-ECA 500 (JEOL, Tokyo, Japan) at 500 MHz or a JNM-AL 400 (JEOL) at 400 MHz. Chemical shifts were reported relative to Me_4Si (δ 0.00) in CDCl_3 or the residual solvent peak in C_6D_6 (δ 7.16). Multiplicity was indicated by one or more of the following: s (singlet); d (doublet); t (triplet); q (quartet); m (multiplet); br (broad). Carbon nuclear magnetic resonance (^{13}C NMR) spectra were recorded using a JNM-ECA 500 at 126 MHz or a JNM-AL 400 at 100 MHz. Chemical shifts were reported relative to CDCl_3 (δ 77.0) or C_6D_6 (δ 128.0). Infrared spectra were recorded using a FT/IR-4100 Fourier-transform infrared spectrometer (JASCO, Tokyo, Japan) with ATR (attenuated total reflectance). Low and high resolution mass spectra were recorded using a JMS-700 mass spectrometer (JEOL) for FAB-MS and a LCMS-IT-TOF (Shimadzu, Kyoto, Japan) for ESI-MS.

Experimental procedures and spectroscopic data.




Silylether 4: To a solution of aniline **3** (2.06 g, 9.40 mmol) in CH_2Cl_2 (94.0 mL) was added a solution of BBr_3 (25.0 g, 94.0 mmol) in CH_2Cl_2 (94.0 mL) at -78°C . The mixture was stirred at -78°C for 20 min, and then warmed to room temperature. After 2 h, saturated aqueous NaHCO_3 and 1M aqueous NaOH were added to the reaction mixture until the mixture became basic. The mixture was extracted with EtOAc . The combined organic layers were washed with brine, and dried over Na_2SO_4 . Concentration under reduced pressure gave a crude demethylated lactone.

To a solution of the above crude lactone in anhydrous DMF (20.0 mL) were added TBSCl (2.80 g, 18.8 mmol) and imidazole (1.90 g, 28.2 mmol) at 0°C . The mixture was stirred at room temperature for 3 h. After addition of water, the mixture was extracted with extracted with Et_2O . The combined organic layers were washed with brine and dried over Na_2SO_4 . After concentration under reduced pressure, purification by flash column chromatography on silica gel (5–40% EtOAc /hexane) gave silylether **4** (1.88 g, 63% in 2 steps) as a pale yellow solid: ^1H NMR (500 MHz, CDCl_3) δ 6.99 (dd, 1H, $J = 8.0, 8.0$ Hz), 6.35 (dd, 1H, $J = 8.0, 1.1$ Hz), 6.27 (dd, 1H, $J = 8.0, 1.1$ Hz), 6.06 (dd, 1H, $J = 1.7, 1.2$ Hz), 4.53 (dd, 2H, $J = 6.3, 5.8$ Hz), 3.76 (br, 2H), 2.72 (dd, 2H, $J = 6.3, 5.7$ Hz), 0.94 (s, 9H), 0.21 (s, 6H); ^{13}C NMR (126 MHz, CDCl_3) δ 164.4, 156.1, 153.1, 144.0, 129.8, 120.3, 115.5, 108.7, 108.6, 66.5, 28.2, 25.6, 18.1, -4.1 ; IR (ATR, cm^{-1}) 3369, 2954, 2891, 2857, 1716, 1625, 1580, 1462, 1398, 1302, 1257, 1219, 1081, 1020; MS (FAB) m/z 320 $[\text{M} + \text{H}]^+$; HRMS calcd for $\text{C}_{17}\text{H}_{26}\text{NO}_3\text{Si}$ $[\text{M} + \text{H}]^+$ 320.1682; Found: m/z 320.1685.



(*E*)-Dienylaniline **5**: To a solution of silyl ether **4** (1.25 g, 3.91 mmol) in CH₂Cl₂ (40.0 mL) was added DIBAL-H (1M in toluene, 7.80 mL, 7.80 mmol) at –78 °C. After the mixture was stirred at –78 °C for 2 h, saturated aqueous Na/K tartrate was added to the reaction solution. The resultant mixture was stirred vigorously at room temperature for 2 h, and extracted with EtOAc. The combined organic layers were washed with brine and dried over Na₂SO₄. Concentration under reduced pressure gave a crude acetal.


To a suspension of MePPh₃Br (4.89 g, 13.7 mmol) in anhydrous THF (25.0 mL) was added KHMDS (1M solution in THF; 12.0 mL, 11.7 mmol) at 0 °C, and the mixture was stirred at 0 °C for 30 min. To the yellow mixture was then added a solution of the above crude acetal in anhydrous THF (15 mL) via cannula. The reaction mixture was stirred at room temperature for 2 h. After addition of water, the resultant mixture was extracted with EtOAc. The combined organic layers were washed with brine and dried over Na₂SO₄. After concentration under reduced pressure, purification by flash column chromatography on silica gel (5-40% EtOAc/hexane) gave (*E*)-dienylaniline **5** (963.1 mg, 77% in 2 steps) as a yellow oil: ¹H NMR (500 MHz, CDCl₃) δ 6.95 (dd, 1H, *J* = 8.0, 8.0 Hz), 6.77 (ddd, 1H, *J* = 16.9, 10.9, 10.3 Hz), 6.36 (dd, 1H, *J* = 8.0, 0.8 Hz), 6.29 (d, 1H, *J* = 11.1 Hz), 6.25 (dd, 1H, *J* = 10.3 Hz), 5.27 (dd, 1H, *J* = 16.9, 1.2 Hz), 5.24 (d, 1H, *J* = 10.3 Hz), 3.75 (br, 1H), 3.63 (br, 1H), 3.60 (br, 2H), 2.98 (br, 1H), 2.38 (br, 1H), 0.88 (s, 9H), 0.08 (s, 6H); ¹³C NMR (126 MHz, CDCl₃) δ 154.9, 144.7, 136.6, 132.4, 132.1, 128.7, 119.3, 115.8, 106.9, 106.0, 60.6, 33.3, 25.7, 18.1, –5.5; IR (ATR, cm^{–1}) 3375, 2955, 2924, 2857, 1618, 1581, 1464, 1234, 1088; MS (FAB) *m/z* 320 [M + H]⁺; HRMS calcd for C₁₈H₃₀NO₂Si [M + H]⁺ 320.2046; Found: *m/z* 320.2045.




(*E*)-Dienylurea **6**: To a solution of (*E*)-dienylaniline **5** (847.9 mg, 2.65 mmol) in CH₂Cl₂ (26.0 mL) was added phenyl isocyanate (317.0 μL, 2.92 mmol) at 0 °C. The mixture was stirred at 0 °C for 13 h. After addition of water, the reaction mixture was extracted with EtOAc. The combined organic layers were washed with brine and dried over Na₂SO₄. After concentration under reduced pressure, purification by short column chromatography on silica gel (10-20% EtOAc/hexane) gave a crude urea as a white solid.

To a solution of the above crude urea in CH₂Cl₂ (50.0 mL) were added Et₃N (2.10 mL, 15.1 mmol) and PhNTf₂ (6.15 g, 17.2 mmol) in some portions. The resultant solution was refluxed at 55 °C for 3 days. The reaction mixture was then cooled to room temperature. After addition of saturated aqueous NH₄Cl, the mixture was extracted with EtOAc. The combined organic layers were washed with brine and dried over Na₂SO₄. After concentration under reduced pressure, purification by flash column chromatography on silica gel (5-20% EtOAc/hexane) gave (*E*)-dienylurea **6** (1.37 g, 90% in 2 steps) as a pale yellow solid: ¹H NMR (500 MHz, CDCl₃) δ 8.21 (dd, 1H, *J* = 8.3, 0.9 Hz), 7.39 (br, 1H), 7.34-7.30 (m, 5H), 7.13-7.09 (m, 1H), 9.97 (dd, 1H, *J* = 8.3, 0.9 Hz), 6.69 (ddd, 1H, *J* = 16.6, 10.9, 10.3 Hz), 6.65 (br, 1H), 6.21 (d, 1H, *J* = 11.1 Hz), 5.38-5.34 (m, 2H), 3.71-3.69 (m, 1H), 3.55-3.51 (m, 1H), 2.97-2.95 (m, 1H), 2.39-2.36 (m, 1H), 0.83 (s, 9H), 0.02 (s, 6H); ¹³C NMR (126 MHz, CDCl₃) δ 152.6, 147.2, 138.8, 137.6, 137.1, 131.4, 129.7, 129.3, 128.9, 126.8, 124.4, 121.4, 121.1, 120.6, 119.7, 118.4 (q, *J* = 321 Hz), 115.1, 61.6, 35.0, 25.9, 18.5, –5.5; IR (ATR, cm^{–1}) 3332, 2954, 2857, 1659, 1550, 1524, 1446, 1420, 1296, 1250, 1207, 1139, 1054, 962; MS


(FAB) m/z 571 $[M + H]^+$; HRMS calcd for $C_{26}H_{34}F_3N_2O_5SSi$ $[M + H]^+$ 571.1910; Found: m/z 570.1910.




(*E*)-Dienylalcohol **7**: To a solution of (*E*)-dienylurea **6** (42.7 mg, 0.0748 mmol) in THF (1.0 mL) was added TBAF (1M in THF, 83.0 μ L, 0.0823 mmol) at 0 °C. The mixture was stirred at 0 °C for 1 h. After addition of saturated aqueous NH_4Cl , the mixture was extracted with EtOAc. The combined organic layers were washed with brine and dried over Na_2SO_4 . After concentration under reduced pressure, purification by flash column chromatography on silica gel (10-40% EtOAc/hexane) gave (*E*)-dienylalcohol **7** (35.3 mg, quant.) as a pale yellow solid: 1H NMR (500 MHz, $CDCl_3$) δ 8.25 (d, 1H, J = 8.6 Hz), 7.77 (br, 1H), 7.32-7.18 (m, 5H), 7.18 (br, 1H), 7.07 (dd, 1H, J = 7.1, 6.9 Hz), 6.94 (d, 1H, J = 8.3 Hz), 6.72 (ddd, 1H, J = 16.9, 10.9, 10.3 Hz), 6.24 (d, 1H, J = 10.9 Hz), 5.39-5.33 (m, 2H), 3.81 (br, 1H), 3.46 (br, 1H), 3.04 (br, 1H), 2.35 (d, 1H, J = 14.6 Hz), 2.23 (br, 1H); ^{13}C NMR (126 MHz, $CDCl_3$) δ 153.2, 147.3, 139.3, 137.9, 137.8, 131.3, 129.2, 129.1, 129.1, 126.1, 124.1, 121.4, 120.9, 119.6, 118.4 (q, J = 321 Hz), 114.8, 60.2, 34.2; IR (ATR, cm^{-1}) 3337, 3010, 2926, 1670, 1579, 1550, 1446, 1420, 1297, 1210, 1138, 1051, 963; MS (FAB) m/z 457 $[M + H]^+$; HRMS calcd for $C_{20}H_{20}F_3N_2O_5S$ $[M + H]^+$ 457.1045; Found: m/z 457.1042.




(*E*)-Dienylurea **8**: To a solution of (*E*)-dienylalcohol **7** (992.0 mg, 2.17 mmol), *p*NsNHBoc (786.0 mg, 2.60 mmol) and PPh_3 (682.0 mg, 2.60 mmol) in THF (12.0 mL) was added a solution of di-*tert*-butyl azodicarboxylate (598.7 mg, 2.60 mmol) in THF (10.0 mL). The mixture was stirred at room temperature for 13.5 h. After addition of saturated aqueous NH_4Cl , the mixture was extracted with EtOAc. The combined organic layers were washed with brine and dried over Na_2SO_4 . After concentration under reduced pressure, purification by flash column chromatography on silica gel (10-40% EtOAc/hexane) gave the mixture of (*E*)-dienylurea **8** and *p*NsNHBoc. The mixture was dissolved in $CHCl_3$, washed with 1M aqueous $NaOH$ and brine, dried over Na_2SO_4 . Concentration under reduced pressure gave (*E*)-dienylurea **8** (1.43 g, 89%) as a pale yellow solid: 1H NMR (500 MHz, $CDCl_3$) δ 8.35-8.32 (m, 3H), 8.04 (d, 2H, J = 9.1 Hz), 7.47 (br, 1H), 7.37-7.29 (m, 5H), 7.17 (br, 1H), 7.13-7.09 (m, 1H), 6.98 (d, 1H, J = 8.3 Hz), 6.79 (ddd, 1H, J = 16.3, 10.9, 10.6 Hz), 6.24 (d, 1H, J = 10.9 Hz), 5.42-5.38 (m, 2H), 3.88-3.78 (m, 2H), 3.15-3.09 (m, 1H), 2.76-2.70 (m, 1H), 1.32 (s, 9H); ^{13}C NMR (126 MHz, $CDCl_3$) δ 152.6, 150.9, 150.5, 147.3, 145.1, 138.5, 138.4, 137.7, 131.3, 129.4, 129.4, 129.1, 128.1, 125.9, 124.6, 124.1, 122.6, 121.3, 120.0, 118.4 (q, J = 321 Hz), 115.1, 86.5, 46.2, 33.4, 27.9; IR (ATR, cm^{-1}) 3349, 2929, 2854, 1732, 1668, 1534, 1446, 1420, 1368, 1351, 1291, 1249, 1212, 1139, 1055, 961; MS (FAB) m/z 741 $[M + H]^+$; HRMS calcd for $C_{31}H_{32}F_3N_4O_{10}S_2$ $[M + H]^+$ 741.1512; Found: m/z 741.1512.




(*E*)-Dienylcarbodiimide **9**: To a solution of (*E*)-dienylurea **8** (62.5 mg, 0.0844 mmol) and PPh₃ (73.5 mg, 0.270 mmol) in CH₂Cl₂ (2.0 mL) were added Et₃N (47.0 μL, 0.338 mmol) and CBr₄ (83.9 mg, 0.253 mmol) at 0 °C. The mixture was stirred at 0 °C for 2.5 h. After concentration of the mixture under reduced pressure, purification of the residue by flash column chromatography on neutral silica gel (5-20% EtOAc/hexane) gave (*E*)-dienylcarbodiimide **9** (56.4 mg, 92%) as a pale-yellow oil. The product was not stable, thus it was used for the next reaction immediately: ¹H NMR (500 MHz, CDCl₃) δ 8.33 (d, 2H, *J* = 8.9 Hz), 8.07 (d, 2H, *J* = 8.8 Hz), 7.37-7.27 (m, 4H), 7.19 (dd, 1H, *J* = 7.5, 7.4 Hz), 7.16-7.13 (m, 3H), 6.80 (ddd, 1H, *J* = 16.6, 10.6, 10.6 Hz), 6.25 (d, 1H, *J* = 11.2 Hz), 5.37-5.32 (m, 2H), 3.89 (dd, 2H, *J* = 7.2, 7.2 Hz), 2.99 (br, 2H), 1.31 (s, 9H); ¹³C NMR (126 MHz, CDCl₃) δ 150.3, 150.2, 147.6, 145.4, 139.3, 137.3, 137.2, 132.4, 132.1, 131.1, 129.5, 129.3, 129.1, 127.9, 125.9, 124.9, 124.4, 123.9, 121.6, 118.4 (q, *J* = 321 Hz), 118.2, 85.2, 45.9, 33.4, 27.7; IR (ATR, cm⁻¹) 3105, 2938, 2857, 2141, 1731, 1591, 1563, 1533, 1476, 1452, 1421, 1366, 1351, 1285, 1250, 1213, 1137, 909. (Compound **9** was too unstable to measure HRMS)



2-Iminoindoline **10**: To a solution of carbodiimide **9** (56.4 mg, 0.0780 mmol) in anhydrous xylene (1.0 mL) were added bis(pinacolato)diboron (39.6 mg, 0.156 mmol) and Pd(OAc)₂ (3.5 mg, 0.0156 mmol) and the reaction atmosphere was replaced by the Ar atmosphere. The mixture was stirred at 50 °C for 1 h, and then cooled to 0 °C. After addition of water (1.0 mL) and sodium perborate tetrahydrate (72.0 mg, 0.468 mmol), the mixture was stirred vigorously at room temperature for 1 h. The mixture was then extracted with EtOAc. The combined organic layers were washed with brine and dried over Na₂SO₄. After concentration under reduced pressure, purification by flash column chromatography on silica gel (20-60% EtOAc/hexane) gave 2-iminoindoline **10** (42.4 mg, 73%) as a yellow oil: ¹H NMR (500 MHz, CDCl₃) δ 8.26 (d, 2H, *J* = 8.9 Hz), 7.94 (d, 2H, *J* = 8.9 Hz), 7.79 (d, 2H, *J* = 8.0 Hz), 7.40-7.34 (m, 4H), 7.13 (dd, 1H, *J* = 7.5, 7.4 Hz), 6.98 (br, 1H), 6.93-6.90 (m, 1H), 6.07 (ddd, 1H, *J* = 15.8, 5.2, 4.8 Hz), 5.78 (d, 1H, *J* = 16.0 Hz), 4.23 (d, 2H, *J* = 4.9 Hz), 3.45 (ddd, 1H, *J* = 14.0, 11.8, 4.0 Hz), 3.22 (ddd, 1H, *J* = 14.3, 12.0, 4.3 Hz), 2.89 (ddd, 1H, *J* = 12.6, 12.6, 4.3 Hz), 2.54 (ddd, 1H, *J* = 12.6, 12.4, 4.0 Hz), 1.94 (br, 1H), 1.31 (s, 9H); ¹³C NMR (126 MHz, CDCl₃) δ 170.5, 158.9, 150.3, 150.0, 145.1, 144.9, 138.6, 133.2, 131.1, 129.2, 129.1, 127.6, 127.5, 124.1, 123.9, 120.0, 118.5 (q, *J* = 320 Hz), 117.9, 114.4, 85.7, 62.8, 59.0, 43.4, 33.0, 27.7; IR (ATR, cm⁻¹) 3380, 3106, 2936, 2877, 1732, 1561, 1534, 1439, 1420, 1349, 1247, 1213, 1138, 1083, 1014, 907; MS (FAB) *m/z* 741 [M + H]⁺; HRMS calcd for C₃₁H₃₂F₃N₄O₁₀S₂ [M + H]⁺ 741.1512; Found: *m/z* 741.1508.




N-Boc-iminoindoline **11**: To a solution of 2-iminoindoline **10** (39.4 mg, 0.0532 mmol) in CH₂Cl₂ (1.0 mL) were added Et₃N (23.0 μL, 0.160 mmol) and TESCl (16.0 μL, 0.106 mmol) at 0 °C. The mixture was stirred at room temperature for 2 h. After addition of water, the mixture was extracted with EtOAc. The combined organic layers were washed with brine and dried over Na₂SO₄. After concentration under reduced pressure, purification by short column chromatography on neutral silica gel (10-30% EtOAc/hexane) gave a crude TES-protected iminoindoline. To a solution of the above crude iminoindoline in CH₂Cl₂ (1.0 mL) were added *i*Pr₂NEt (37.0 μL, 0.213 mmol), Boc₂O (34.9 mg, 0.160 mmol) and DMAP (6.5 mg, 0.0532 mmol) at 0 °C. The mixture was stirred at room temperature for 1.5 h. After concentration of the resultant mixture under reduced pressure, purification of the residue by flash column chromatography on silica gel (10-30% EtOAc/hexane) gave *N*-Boc-iminoindoline **11** (33.7 mg, 84% in 2 steps) as a colorless oil: ¹H NMR (500 MHz, CDCl₃) δ 8.26 (d, 2H, *J* = 8.9 Hz), 8.05 (d, 2H, *J* = 8.8 Hz), 7.73 (d, 1H, *J* = 8.0 Hz), 7.41 (dd, 1H, *J* = 8.6, 8.3 Hz), 7.31 (dd, 2H, *J* = 7.8, 7.7 Hz), 7.12 (d, 1H, *J* = 8.3 Hz), 7.06-7.02 (m, 3H), 5.93 (d, 1H, *J* = 15.5 Hz), 5.66 (d, 1H, *J* = 15.5 Hz), 4.17 (d, 2H, *J* = 4.3 Hz), 3.85-3.79 (m, 1H), 3.65-3.62 (m, 1H), 2.70-2.63 (m, 2H), 1.27 (s, 9H), 1.18 (s, 9H), 0.92 (dd, 9H, *J* = 8.0, 7.8 Hz), 0.57 (q, 6H, *J* = 7.8 Hz); ¹³C NMR (126 MHz, CDCl₃) δ 153.2, 150.2, 150.1, 149.0, 147.9, 145.9, 145.6, 143.5, 131.5, 130.5, 129.4, 129.1, 128.7, 123.9, 123.8, 122.1, 120.5, 118.3 (q, *J* = 321 Hz), 115.8, 114.1, 85.3, 84.8, 62.6, 54.0, 43.2, 34.7, 27.7, 27.4, 6.7, 4.3; IR (ATR, cm⁻¹) 2955, 2876, 1731, 1698, 1617, 1594, 1535, 1456, 1421, 1370, 1348, 1287, 1251, 1218, 1141, 1046, 1014, 917, 822; MS (FAB) *m/z* 955 [M + H]⁺; HRMS calcd for C₄₂H₅₄F₃N₄O₁₂S₂Si [M + H]⁺ 955.2901; Found: *m/z* 955.2900.




Allyl bromide **12**: To a solution of *N*-Boc-iminoindoline **11** (21.9 mg, 0.0229 mmol) in THF (0.5 mL) was added TBAF (48.1 μL, 0.0481 mmol) at 0 °C. The mixture was stirred at room temperature for 30 min. After addition of saturated aqueous NH₄Cl, the mixture was extracted with EtOAc. The combined organic layers were washed with brine and dried over Na₂SO₄. After concentration under reduced pressure, purification by flash column chromatography on silica gel (20-60% EtOAc/hexane) gave an allyl alcohol (15.1 mg, 93%) as a pale yellow oil: ¹H NMR (500 MHz, CDCl₃) δ 8.24 (d, 2H, *J* = 8.8 Hz), 8.07 (d, 2H, *J* = 8.8 Hz), 7.29 (dd, 2H, *J* = 7.5, 7.2 Hz), 7.19 (d, 1H, *J* = 7.7 Hz), 7.13-7.09 (m, 1H), 7.04-6.99 (m, 3H), 6.59 (d, 1H, *J* = 8.0 Hz), 6.03 (d, 1H, *J* = 15.1 Hz), 5.77 (d, 1H, *J* = 15.4 Hz), 4.09 (br, 2H), 3.86-3.80 (m, 1H), 3.69-3.63 (m, 1H), 2.78 (ddd, 1H, *J* = 12.3, 12.1, 4.6 Hz), 2.58 (ddd, 1H, *J* = 12.1, 12.0, 4.3 Hz), 1.27 (s, 9H), 1.21 (s, 9H); ¹³C NMR (126 MHz, CDCl₃) δ 156.0, 152.7, 150.3,

150.2, 149.2, 148.4, 145.5, 142.2, 131.3, 129.8, 129.6, 129.4, 129.1, 123.8, 123.7, 120.4, 115.2, 112.7, 106.6, 85.2, 84.2, 62.9, 53.5, 43.9, 35.2, 27.7, 27.5; IR (ATR, cm^{-1}) 3445, 2980, 1729, 1695, 1535, 1450, 1360, 1352, 1270, 1085, 910, 730; MS (FAB) m/z 709 $[\text{M} + \text{H}]^+$; HRMS calcd for $\text{C}_{35}\text{H}_{41}\text{N}_4\text{O}_{10}\text{S}$ $[\text{M} + \text{H}]^+$ 709.2543; Found: m/z 709.2543.


To a solution of the above allyl alcohol (169.6 mg, 0.239 mmol) and PPh_3 (157.4 mg, 0.598 mmol) in CH_2Cl_2 (2.5 mL) was added CBr_4 (158.5 mg, 0.478 mmol) at 0 °C. The mixture was stirred at 0 °C for 30 min. After addition of water, the mixture was extracted with EtOAc. The combined organic layers were washed with brine and dried over Na_2SO_4 . After concentration under reduced pressure, purification by flash column chromatography on silica gel (10-40% EtOAc/hexane) gave allylbromide **12** (180.3 mg, 98%) as a white solid: ^1H NMR (500 MHz, CDCl_3) δ 8.26 (d, 2H, $J = 8.6$ Hz), 8.08 (d, 2H, $J = 8.8$ Hz), 7.32 (dd, 2H, $J = 8.0, 7.7$ Hz), 7.32-7.27 (m, 1H), 7.21-7.17 (m, 1H), 7.07-7.02 (m, 3H), 6.63 (d, 1H, $J = 8.0$ Hz), 6.05 (d, 1H, $J = 15.2$ Hz), 5.87-5.81 (m, 1H), 3.96-3.89 (m, 2H), 3.82 (ddd, 1H, $J = 14.7, 11.1, 4.6$ Hz), 3.67 (dd, 1H, $J = 12.0, 11.2$ Hz), 2.77 (dd, 1H, $J = 12.3, 10.9$ Hz), 2.58 (ddd, 1H, $J = 12.3, 12.0, 4.3$ Hz), 1.29 (s, 9H), 1.20 (s, 9H); ^{13}C NMR (126 MHz, CDCl_3) δ 154.8, 152.2, 150.3, 150.2, 149.2, 148.3, 145.5, 142.4, 134.6, 130.1, 129.4, 129.1, 126.8, 123.8, 120.5, 114.8, 112.8, 107.4, 107.3, 85.3, 84.2, 53.5, 43.8, 34.8, 32.1, 27.8, 27.4; IR (ATR, cm^{-1}) 3445, 2980, 1729, 1695, 1599, 1532, 1460, 1366, 1348, 1277, 1250, 1143, 1085, 1061, 968, 909, 852, 730, 605, 578; MS (FAB) m/z 771 $[\text{M} + \text{H}]^+$; HRMS calcd for $\text{C}_{35}\text{H}_{40}\text{BrN}_4\text{O}_9\text{S}$ $[\text{M} + \text{H}]^+$ 771.1699; Found: m/z 771.1696.



Tetracyclic compound **18a**: A suspension of allylbromide **12** (300.0 mg, 0.389 mmol), $\text{Bi}(\text{OTf})_3$ (25.5 mg, 0.0389 mmol), AgOTf (104.8 mg, 0.408 mmol), MS4A (300 mg) and K_2CO_3 (161.7 mg, 1.17 mmol) in CH_2Cl_2 (40.0 mL) was stirred at room temperature for 15 min. After addition of water, the mixture was then filtered through Celite pad. The filtrate was extracted with EtOAc. The combined organic layers were dried over Na_2SO_4 . After concentration under reduced pressure, purification by flash column chromatography on silica gel (10-60% EtOAc/hexane) gave a tetracyclic compound **18a** (215.8 mg, 80%) as a yellow oil: ^1H NMR (500 MHz, CDCl_3) δ 8.21 (d, 2H, $J = 8.8$ Hz), 7.89 (d, 2H, $J = 8.9$ Hz), 7.52 (d, 1H, $J = 8.3$ Hz), 7.41 (dd, 1H, $J = 7.8, 1.1$ Hz), 7.36 (dd, 1H, $J = 7.5, 7.4$ Hz), 7.24 (ddd, 1H, $J = 8.3, 8.3, 0.8$ Hz), 7.18 (dd, 1H, $J = 7.5, 7.4$ Hz), 7.13 (d, 1H, $J = 7.4$ Hz), 6.69 (d, 1H, $J = 8.0$ Hz), 6.48 (ddd, 1H, $J = 17.4, 10.0, 9.1$ Hz), 5.81 (d, 1H, $J = 9.1$ Hz), 5.67 (d, 1H, $J = 17.5$ Hz), 4.00 (d, 1H, $J = 9.7$ Hz), 3.57-3.50 (m, 1H), 3.26-3.20 (m, 1H), 2.19-2.11 (m, 2H), 1.70 (s, 9H), 1.26 (s, 9H); ^{13}C NMR (126 MHz, CDCl_3) δ 163.0, 152.9, 150.2, 149.9, 149.3, 145.0, 143.6, 142.9, 136.2, 130.7, 129.5, 128.6, 126.5, 126.0, 125.8, 125.1, 124.7, 123.8, 114.5, 114.2, 108.0, 85.1, 84.2, 50.1, 47.9, 43.8, 28.2, 27.7, 27.3; IR (ATR, cm^{-1}) 3449, 2979, 2919, 1731, 1654, 1599, 1533, 1460, 1368, 1348, 1282, 1236, 1148, 1088, 889; MS (FAB) m/z 691 $[\text{M} + \text{H}]^+$; HRMS calcd for $\text{C}_{35}\text{H}_{39}\text{N}_4\text{O}_9\text{S}$ $[\text{M} + \text{H}]^+$ 691.2438; Found: m/z 691.2439.




Triflate **17a**: To a solution of tetracyclic compound **18a** (213.0 mg, 0.308 mmol) in CH_2Cl_2 (5.0 mL) were added pyridine (87.3 μL , 1.08 mmol) and Tf_2O (103.5 μL , 0.616 mmol) at 0 °C. The mixture was stirred at 0 °C for 2.5 h. After addition of water, the mixture was extracted with EtOAc. The combined organic layers were washed with brine and dried over Na_2SO_4 . After concentration under reduced pressure, purification by flash column chromatography on silica gel (5-20% EtOAc/hexane) gave triflate **17a** (230.3 mg, 91%) as a yellow oil: ^1H NMR (500 MHz, CDCl_3) δ 8.24 (d, 2H, $J = 8.6$ Hz), 7.97 (d, 1H, $J = 8.3$ Hz), 7.91 (d, 2H, $J = 8.9$ Hz), 7.45 (dd, 1H, $J = 8.6, 8.3$ Hz), 7.37-7.32 (m, 2H), 7.23-7.14 (m, 3H), 6.28 (ddd, 1H, $J = 16.9, 10.0, 9.8$ Hz), 5.55 (d, 1H, $J = 10.0$ Hz), 5.30 (d, 1H, $J = 16.9$ Hz), 3.88 (d, 1H, $J = 9.7$ Hz), 3.53-3.49 (m, 1H), 3.38-3.32 (m, 1H), 2.26-2.22 (m, 2H), 1.70 (s, 9H), 1.27 (s, 9H); ^{13}C NMR (126 MHz, CDCl_3) δ 162.3, 150.2, 149.7, 149.1, 147.1, 145.2, 144.5, 143.4, 133.6, 131.0, 129.5, 128.4, 126.8, 126.3, 125.7, 125.5, 123.8, 121.5, 119.5, 118.2 (q, $J = 318$ Hz), 114.9, 114.3, 85.3, 84.8, 51.0, 48.1, 43.2, 28.1, 27.8, 27.6; IR (ATR, cm^{-1}) 2982, 2933, 1729, 1661, 1613, 1534, 1455, 1423, 1369, 13647, 1291, 1217, 1143, 1086, 1033, 922; MS (FAB) m/z 823 $[\text{M} + \text{H}]^+$; HRMS calcd for $\text{C}_{36}\text{H}_{38}\text{F}_3\text{N}_4\text{O}_{11}\text{S}_2$ $[\text{M} + \text{H}]^+$ 823.1931; Found: m/z 823.1929.




Coupling product **20**: To a solution of triflate **17a** (30.0 mg, 0.0365 mmol) and vinyl boronate **19** (20.8 mg, 0.0730 mmol) in toluene (1.0 mL) and EtOH (0.1 mL) were added 0.5 M aqueous Na_2CO_3 (220.0 μL , 0.110 mmol) and $\text{Pd}(\text{PPh}_3)_4$ (4.2 mg, 3.65×10^{-3} mmol). The reaction atmosphere was replaced by the Ar atmosphere, and the mixture was stirred at 100 °C for 7 h. After the reaction mixture was then cooled to room temperature, the resultant mixture was extracted with EtOAc. The combined organic layers were washed with brine and dried over Na_2SO_4 . After concentration under reduced pressure, purification by flash column chromatography on silica gel (5-20% EtOAc/hexane) gave coupling product **20** (16.9 mg, 56%) as a yellow oil: ^1H NMR (500 MHz, CDCl_3) δ 8.21 (d, 2H, $J = 8.9$ Hz), 7.86 (d, 2H, $J = 9.2$ Hz), 7.35-7.31 (m, 5H), 7.18-7.17 (m, 2H), 6.90 (d, 1H, $J = 15.5$ Hz), 6.30 (ddd, 1H, $J = 16.9, 10.1, 10.0$ Hz), 6.13 (d, 1H, $J = 15.7$ Hz), 5.48 (dd, 1H, $J = 10.0, 1.5$ Hz), 5.27 (d, 1H, $J = 16.9$ Hz), 3.85 (d, 1H, $J = 10.0$ Hz), 3.41 (ddd, 1H, $J = 14.3, 13.7, 4.0$ Hz), 3.26-3.19 (m, 1H), 2.29 (ddd, 1H, $J = 12.9, 12.8, 5.5$ Hz), 2.16 (ddd, 1H, $J = 12.9, 12.0, 4.0$ Hz), 1.67 (s, 9H), 1.68 (s, 3H), 1.30 (s, 3H), 1.29 (s, 9H), 0.15 (s, 9H); ^{13}C NMR (126 MHz, CDCl_3) δ 164.3, 149.8, 149.6, 149.7, 145.2, 144.1, 142.6, 139.0, 136.1, 134.6, 129.4, 129.0, 128.2, 126.7, 126.3, 125.7, 125.3, 125.3, 125.0, 123.8, 123.7, 122.5, 122.4, 113.5, 85.1, 84.0, 74.0, 51.5, 48.1, 44.3, 30.1,

30.1, 28.2, 27.9, 2.6; IR (ATR, cm^{-1}) 2978, 1727, 1655, 1597, 1575, 1533, 1474, 1452, 1368, 1347, 1291, 1249, 1150, 1087, 1034, 840, 748, 713, 685, 628, 602; MS (FAB) m/z 831 $[\text{M} + \text{H}]^+$; HRMS calcd for $\text{C}_{43}\text{H}_{55}\text{N}_4\text{O}_9\text{SSi}$ $[\text{M} + \text{H}]^+$ 831.3459; Found: m/z 831.3448.




Aminal 28: To a solution of coupling product **20** (50.0 mg, 0.0602 mmol) in THF (6.0 mL) was added catechol borane solution (1M in THF, 75.3 μL , 0.0753 mmol) at 0 °C. The mixture was stirred at 0 °C for 2 h. After addition of water, the resultant mixture was extracted with EtOAc. The combined organic layers were washed with brine and dried over Na_2SO_4 . After concentration under reduced pressure, purification by flash column chromatography on silica gel (5-20% EtOAc/hexane) gave aminal **28** (32.6 mg, 65%, dr = 3.3:1) as a yellow oil: (major diastereomer) ^1H NMR (500 MHz, CDCl_3) δ 8.14 (d, 2H, J = 8.9 Hz), 7.93 (d, 2H, J = 8.9 Hz), 7.75 (br, 1H), 7.30 (d, 1H, J = 8.0 Hz), 7.27-7.24 (m, 1H), 7.20-7.14 (m, 2H), 7.06 (d, 1H, J = 15.8 Hz), 6.89 (dd, 1H, J = 7.8, 7.4 Hz), 6.83 (d, 1H, J = 8.0 Hz), 6.07 (d, 1H, J = 15.8 Hz), 6.05-5.98 (m, 2H), 5.61 (dd, 1H, J = 10.0, 1.7 Hz), 5.35 (dd, 1H, J = 16.9, 1.5 Hz), 4.89 (s, 1H), 4.15 (d, 1H, J = 10.3 Hz), 4.13-4.08 (m, 1H), 3.29 (ddd, 1H, J = 14.1, 14.1, 4.0 Hz), 2.08 (ddd, 1H, J = 12.6, 12.6, 4.3 Hz), 1.86 (ddd, 1H, J = 12.9, 12.9, 4.3 Hz), 1.65 (s, 9H), 1.64 (s, 3H), 1.28 (s, 3H), 1.25 (s, 9H), 0.15 (s, 9H); ^{13}C NMR (126 MHz, CDCl_3) δ 150.4, 150.2, 145.7, 144.7, 140.6, 137.8, 137.1, 131.5, 129.5, 129.2, 128.9, 127.8, 127.7, 127.0, 125.4, 123.8, 123.7, 123.5, 121.9, 120.1, 116.9, 113.7, 84.6, 83.3, 78.3, 74.1, 54.8, 50.6, 44.8, 30.6, 30.4, 28.6, 27.9, 2.8; IR (ATR, cm^{-1}) 2997, 2918, 1731, 1696, 1534, 1467, 1370, 1347, 1235, 1089, 887, 627 ; MS (FAB) m/z 833 $[\text{M} + \text{H}]^+$; HRMS calcd for $\text{C}_{43}\text{H}_{57}\text{N}_4\text{O}_9\text{SSi}$ $[\text{M} + \text{H}]^+$ 833.3616; Found: m/z 833.3616.



Aminal 29: To a solution of aminal **28** (10.8 mg, 0.0130 mmol) in THF (1.3 mL) was added TBAF (1M in THF, 15.6 μL , 0.0156 mmol) at 0 °C. The mixture was stirred at 0 °C for 4 h. After addition of saturated aqueous NH_4Cl , the resultant mixture was extracted with EtOAc. The combined organic layers were washed with brine and dried over Na_2SO_4 . After concentration under reduced pressure, purification by flash column chromatography on silica gel (5-30% EtOAc/hexane) gave an alcohol (7.6 mg, 77%) as a yellow oil. To a solution of the above alcohol (7.6 mg, 9.99×10^{-3} mmol) in MeCN (1.0 mL) were added K_2CO_3 (9.6 mg, 0.0695 mmol) and PhSH (6.3 μL , 0.0614 mmol). The mixture was stirred at room temperature for 12 h, and then diluted with EtOAc. The organic layer was washed with water and brine, and then dried over Na_2SO_4 . After concentration under reduced pressure, purification by flash column chromatography on silica gel (5-30% EtOAc/hexane) gave

aminal **29** (4.0 mg, 70%) as a yellow oil: ^1H NMR (500 MHz, CDCl_3) δ 7.70 (br, 1H), 7.33-7.26 (m, 2H), 7.21 (dd, 1H, $J = 8.0, 8.0$ Hz), 7.14-7.10 (m, 2H), 6.84 (ddd, 1H, $J = 7.8, 7.8, 1.2$ Hz), 6.78 (d, 1H, $J = 7.7$ Hz), 6.14-6.07 (m, 2H), 5.97 (br, 1H), 5.61 (dd, 1H, $J = 10.0, 1.7$ Hz), 5.40 (dd, 1H, $J = 17.2, 1.8$ Hz), 4.85 (s, 1H), 4.40 (br, 1H), 4.15 (d, 1H, $J = 9.7$ Hz), 2.94 (br, 1H), 2.73 (br, 1H), 1.93 (br, 1H), 1.86 (br, 1H), 1.63 (s, 9H), 1.44 (s, 3H), 1.40 (s, 3H), 1.32 (s, 9H); ^{13}C NMR (126 MHz, CDCl_3) δ 155.6, 144.7, 142.9, 140.7, 139.3, 137.2, 131.3, 129.9, 129.1, 128.8, 127.6, 127.3, 124.0, 122.9, 120.9, 120.1, 117.2, 113.9, 83.2, 78.9, 78.4, 71.2, 55.5, 50.8, 37.1, 30.2, 29.3, 28.53, 28.49; IR (ATR, cm^{-1}) 2978, 2916, 1469, 1384, 1283, 1234, 1089, 888, 628; MS (FAB) m/z 576 $[\text{M} + \text{H}]^+$; HRMS calcd for $\text{C}_{34}\text{H}_{45}\text{N}_3\text{O}_5$ $[\text{M}]^+$ 575.3359; Found: m/z 575.3359.



Pentacyclic compound **30**: To a mixture of aminal **29** (6.9 mg, 0.0120 mmol) and MS4A (7.0 mg) in CH_2Cl_2 (1.2 mL) was added $\text{Bi}(\text{OTf})_3$ (0.8 mg, 1.2×10^{-3} mmol) at 0 °C. The mixture was stirred at 0 °C for 2 h, and then warmed to room temperature and stirred for 1 h. After addition of saturated aqueous NaHCO_3 , and the mixture was diluted with EtOAc. The organic layer was washed with brine and dried over Na_2SO_4 . After concentration under reduced pressure, purification by flash column chromatography on silica gel (5-20% EtOAc/hexane) gave a pentacyclic compound **30** (3.7 mg, 55%) as a yellow oil: ^1H NMR (500 MHz, CDCl_3) δ 7.74 (br, 1H), 7.23-7.18 (m, 2H), 7.11 (dd, 1H, $J = 7.2, 7.1$ Hz), 7.00 (d, 1H, $J = 7.8$ Hz), 6.81 (dd, 1H, $J = 8.3, 7.9$ Hz), 6.75 (d, 1H, $J = 8.0$ Hz), 5.94 (d, 1H, $J = 9.2$ Hz), 5.92-5.88 (m, 1H), 5.84 (br, 1H), 5.42 (dd, 1H, $J = 16.6, 2.3$ Hz), 5.38 (dd, 1H, $J = 9.5, 2.3$ Hz), 5.05 (s, 1H), 5.00 (d, 1H, $J = 8.3$ Hz), 4.10 (d, 1H, $J = 10.0$ Hz), 3.90 (dd, 1H, $J = 14.0, 4.0$ Hz), 2.10 (ddd, 1H, $J = 14.6, 11.4, 5.4$ Hz), 2.03-1.96 (m, 2H), 1.85 (s, 3H), 1.65 (s, 3H), 1.63 (s, 9H), 1.46 (s, 9H); ^{13}C NMR (126 MHz, CDCl_3) δ 155.4, 153.8, 144.7, 142.8, 138.5, 137.7, 132.7, 131.2, 130.7, 128.2, 127.6, 126.0, 124.7, 122.8, 120.0, 118.2, 116.7, 114.6, 82.8, 79.2, 78.7, 58.7, 58.2, 50.7, 41.0, 28.5, 28.4, 25.2, 23.5, 18.4; IR (ATR, cm^{-1}) 2977, 2919, 1691, 1466, 1391, 1341, 1279, 1235, 1089, 889, 756, 628, 523; MS (FAB) m/z 558 $[\text{M} + \text{H}]^+$; HRMS calcd for $\text{C}_{34}\text{H}_{42}\text{N}_3\text{O}_4$ $[\text{M} - \text{H}]^-$ 556.3175; Found: m/z 556.3177. (ESI) HRMS calcd for $\text{C}_{34}\text{H}_{44}\text{N}_3\text{O}_4$ $[\text{M} + \text{H}]^+$ 558.3332; Found: m/z 558.3311.

Acknowledgements

This work was supported by a Grant-in-Aid from JSPS KAKENHI (Grant Nos. JP17H05051 (CT), JP18H04407 (CT) and JP16H06384 (YT)) and a Grant-in-Aid from the Uehara Memorial Foundation, Japan (CT).

References

- Numata A, et al. Communesins, cytotoxic metabolites of a fungus isolated from a marine alga. *Tetrahedron Lett.* 1993;34:2355–2358.

2. Jadulco R, et al. New Communesin Derivatives from the Fungus *Penicillium* sp. Derived from the Mediterranean Sponge *Axinella verrucosa*. *J. Nat. Prod.* 2004;67:78–81.
3. Hayashi H, Matsumoto H, Akiyama K. New Insecticidal Compounds, Communesins C, D and E, from *Penicillium expansum* Link MK-57. *Biosci. Biotechnol. Biochem.* 2004;68:753–756.
4. Dalsgaard PW, Blunt JW, Munro MHG, Frisvad JC, Christophersen C. Communesins G and H, New Alkaloids from the Psychrotolerant Fungus *Penicillium rivulum*. *J. Nat. Prod.* 2005;68:258–261.
5. Kerzaon I, et al. Structural investigation and elucidation of new communesins from a marine-derived *Penicillium expansum* Link by liquid chromatography/electrospray ionization mass spectrometry. *Rapid Commun. Mass Spectrom.* 2009;23:3928–3938.
6. Fan YQ, et al. Alkaloids with Cardiovascular Effects from the Marine-Derived Fungus *Penicillium expansum* Y32. *Mar. Drugs.* 2015;13:6489–6504.
7. Verbitski SM, Mayne CL, Davis RA, Concepcion GP, Ireland CM. Isolation, Structure Determination, and Biological Activity of a Novel Alkaloid, Perophoramidine, from the Philippine Ascidian *Perophora namei*. *J. Org. Chem.* 2002;67:7124–7126.
8. Lin HC, et al. Elucidation of the concise biosynthetic pathway of the communesin indole alkaloids. *Angew. Chem. Int. Ed.* 2015;54:3004–3007.
9. Trost BM, Osipov M. Recent Advances on the Total Syntheses of Communesin Alkaloids and Perophoramidine. *Chem. Eur. J.* 2015;21:16318–16343.
10. Siengalewicz P, Gaich T, Mulzer J. It All Began with an Error: The Nomofungin/Communesin Story. *Angew. Chem. Int. Ed.* 2008;47:8170–8276.
11. Johnston CA, et al. Polycyclic ethers and an unexpected dearomatisation reaction during studies towards the bioactive alkaloid, perophoramidine. *Tetrahedron* 2018;74:3339–3347.
12. Shao W, You SL. Highly Diastereo- and Enantioselective Synthesis of Tetrahydro-5*H*-Indolo[2,3-*b*]quinolines through Copper-Catalyzed Propargylic Dearomatization of Indoles. *Chem. Eur. J.* 2017;23:12489–12493.
13. Hoang A, Popov K, Somfai P. An Efficient Synthesis of (±)-Dehaloperophoramidine. *J. Org. Chem.* 2017;82:2171–2176.
14. Popov K, Hoang A, Somfai P. Concise Total Synthesis of Dehaloperophoramidine. *Angew. Chem. Int. Ed.* 2016;55:1801–1804.
15. Yang J, Wu H, Shen L, Qin Y. Total Synthesis of (±)-Communesin F. *J. Am. Chem. Soc.* 2007;129:13794–13795.
16. Liu P, Seo JH, Weinreb SM. Total Synthesis of the Polycyclic Fungal Metabolite (±)-Communesin F. *Angew. Chem. Int. Ed.* 2010;49:2000–2003.
17. Belmar J, Funk RL. Total Synthesis of (±)-Communesin F via a Cycloaddition with Indol-2-one. *J. Am. Chem. Soc.* 2012;134:16941–16943.
18. Zuo Z, Xie W, Ma D. Total Synthesis and Absolute Stereochemical Assignment of (–)-Communesin F. *J. Am. Chem. Soc.* 2010;132:13226–13228.
19. Zuo Z, Ma D. Enantioselective Total Syntheses of Communesins A and B. *Angew. Chem. Int. Ed.* 2011;50:12008–12011.

- 480 20. Han S-J, et al. A Diastereodivergent Synthetic Strategy for the Syntheses of Communesin F and Perophoramidine.
481 *Org. Lett.* 2014;16:3316–3319.
- 482 21. Lathrop SP, Pompeo M, Chang W-T, Movassaghi M. Convergent and Biomimetic Enantioselective Total
483 Synthesis of (–)-Communesin F. *J. Am. Chem. Soc.* 2016;138:7763–7769.
- 484 22. Liang X, et al. Ir-Catalyzed Asymmetric Total Synthesis of (–)-Communesin F. *J. Am. Chem. Soc.*
485 2017;139:3364–3367.
- 486 23. Park J, Jean A, Chen DY-K. Asymmetric Total Syntheses of Communesin F and a Putative Member of the
487 Communesin Family. *Angew. Chem. Int. Ed.* 2017;56:14237–14240.
- 488 24. Ishida T, Tsukano C, Takemoto Y. Synthesis of 2-Iminoindolines via Samarium Diiodide Mediated Reductive
489 Cyclization of Carbodiimides. *Chem. Lett.* 2012;41:44–46.
- 490 25. Ishida T, Ikota H, Kurahashi K, Tsukano C, Takemoto Y. Dearomatizing Conjugate Addition to Quinoliny
491 Amidines for the Synthesis of Dehaloperophoramidine through Tandem Arylation and Allylation. *Angew. Chem.*
492 *Int. Ed.* 2013;52:10204–10207.
- 493 26. Ishida T, Takemoto Y. Synthetic study of perophoramidine: construction of pentacyclic core structure via SmI₂-
494 mediated reductive cyclization. *Tetrahedron* 2013;69:4517–4523.
- 495 27. Nanjo T, Tsukano C, Takemoto Y. Synthesis of 3,3-Disubstituted 2-Aminoindolenines by Palladium-Catalyzed
496 Allylic Amidination with Isocyanide. *Synlett* 2014;25:1473–1477.
- 497 28. Suetsugu S, Nishiguchi H, Tsukano C, Takemoto Y. Asymmetric Synthesis of (–)-Aurantoclavine via Palladium-
498 Catalyzed Intramolecular Allylic Amination. *Org. Lett.* 2014;16:996–999.
- 499 29. Suetsugu S, Tsukano C, Takemoto Y. Synthetic Studies towards Communesins: Diastereoselective Oxidative
500 Rearrangement of Aurantoclavine Derivatives. *Eur. J. Org. Chem.* 2016;2016:108–115.
- 501 30. Tsukano C, Nakajima M, Hande SM, Takemoto Y. Palladium-catalyzed intramolecular carboborylation of 1,3-
502 diene and synthesis of ABCD ring of communesins. *Org. Biomol. Chem.* In press 2018, 10.1039/C8OB02224K.
- 503 31. Ikota H, Tsukano C, Takemoto Y. Concise Synthesis of (±)-Aurantoclavine. *Heterocycles.* In press 2018.
- 504 32. Hande SM, Nakajima M, Kamisaki H, Tsukano C, Takemoto Y. Flexible Strategy for Syntheses of
505 Spirooxindoles using Palladium-Catalyzed Carbosilylation and Sakurai-Type Cyclization. *Org. Lett.*
506 2011;13:1828–1831.
- 507 33. Also see Supplementary Information.
- 508 34. Fukuyama T, Cheung M, Kan T. *N*-Carboalkoxy-2-Nitrobenzenesulfonamides: A Practical Preparation of *N*-Boc-,
509 *N*-Alloc-, and *N*-Cbz-Protected Primary Amines. *Synlett* 1999;1999:1301–1303.
- 510 35. Ishiyama T, Itoh Y, Kitano T, Miyaura N. Synthesis of arylboronates via the palladium(0)-catalyzed cross-
511 coupling reaction of tetra(alkoxy)diborons with aryl triflates. *Tetrahedron Lett.* 1997;38:3447–3450.
- 512 36. Papagni A, et al. Cross-Coupling of 5,11-Dibromotetracene Catalyzed by a Triethylammonium Ion Tagged
513 Diphenylphosphine Palladium Complex in Ionic Liquids. *Organometallics* 2011;30:4325–4329.
- 514 37. Hayashi R, Cook GR. Bi(OTf)₃-catalyzed 5-exo-trig cyclization via halide activation. *Tetrahedron Lett.*
515 2008;49:3888–3890.
- 516 38. Giera DS, Schneider C. InCl₃-Catalyzed Allylic Friedel–Crafts Reactions toward the Stereocontrolled Synthesis
517 of 1,2,3,4-Tetrahydroquinolines. *Org. Lett.* 2010;12:4884–4887.

- 518 39. Kargbo RB, Hashemi ZS, Roy S, Jin X, Herr RJ. Synthesis of 3-benzazepines and azepino[4,5-*b*]heterocyclic
519 ring systems via intramolecular Friedel–Crafts cyclization. *Tetrahedron Lett.* 2013;54:2018–2021.
- 520 40. Okada M, Sato I, Cho SJ, Dubnau D, Sakagami Y. Chemical synthesis of ComX pheromone and related peptides
521 containing isoprenoidal tryptophan residues. *Tetrahedron* 2006;62:8907–8918.
- 522 41. Kawai N, Abe R, Uenishi J. Lewis acid-catalyzed intramolecular amination via 1,3-chirality transfer. *Tetrahedron*
523 *Lett.* 2009;50:6580–6583.
- 524 42. Kawai N, Abe R, Matsuda M, Uenishi J. Synthesis of Chiral 1-Substituted Tetrahydroisoquinolines by the
525 Intramolecular 1,3-Chirality Transfer Reaction Catalyzed by Bi(OTf)₃. *J. Org. Chem.* 2011;76:2102–2114.
- 526