

The limiting absorption principle
for Dirac operators

Ritsumeikan Univ. Osanobu YAMADA

In the present paper we are concerned with the Dirac operator

$$L = -i \sum_{j=1}^3 \alpha_j \frac{\partial}{\partial x_j} + \beta + Q(x) \quad (x \in \mathbb{R}^3),$$

which appears in relativistic quantum mechanics. The matrices α_j and β (called the Dirac matrices) are 4×4 Hermitian matrices with the anti-commutation relation

$$\alpha_j \alpha_k + \alpha_k \alpha_j = 2 \delta_{jk} I, \quad j, k = 1, 2, 3, 4$$

($\alpha_4 = \beta$, I is the unit matrix). The potential $Q(x)$ is a 4×4 Hermitian matrix-valued function.

The unperturbed operator L_0 (as $Q(x) \equiv 0$) defined on $\mathcal{C}_0^\infty = \left[\mathcal{C}_0^\infty(\mathbb{R}^3) \right]^4$ is essentially selfadjoint in $\mathcal{L}^2 = \left[L^2(\mathbb{R}^3) \right]^4$, that is, the closure $H_0 = (L_0)^{**}$ is the unique selfadjoint extension of L_0 .

Then it turns out that the domain of H_0

$$D(H_0) = \mathcal{H}^1 = \left[H^1(\mathbb{R}^3) \right]^4$$

$$(H^1(\mathbb{R}^3) = \left\{ u(x) \in L^2(\mathbb{R}^3) ; \frac{\partial}{\partial x_j} u(x) \in L^2(\mathbb{R}^3), j=1,2,3 \right\},$$

where the derivatives are taken in the distribution sense), and the essential spectrum

$$\sigma_e(H_0) = (-1, 1)^c$$

(the complement of the interval $(-1, 1)$ in the real line).

Proposition 1. Let $Q(x)$ satisfy

$$(1) \quad |Q(x)| \longrightarrow 0 \quad (|x| \rightarrow \infty)$$

and

$$(2) \quad \sup_{x \in \mathbb{R}^3} \int_{|x-y| \leq 1} |Q(y)|^2 |x-y|^{-1-\delta} dy < +\infty$$

for some $\delta > 0$, where $|M|$ for a matrix M indicates the square root of the maximum eigenvalue of $M^* M$. Then $L = L_0 + Q(x)$ has the unique selfadjoint realization $H = H_0 + Q$ with the domain $D(H) = \mathcal{H}^1$ and the essential spectrum $\sigma_e(H) = (-1, 1)^c$.

For the proof of the above proposition see, e.g., Jörgens [1].

Remark 1. Coulomb potentials do not fulfill the condition (2). But

the above result holds also by replacing (2) by a condition

$$(3) \quad \begin{cases} |Q(x)| \leq \frac{c_1}{|x|} & (|x| \leq 1), \quad \frac{1}{2} > c_1 > 0 \\ |Q(x)| \leq c_2 & (|x| \geq 1), \quad c_2 > 0 \end{cases}$$

(see Jörgen [1] and Arai [2]) .

There are many works related to the spectral and scattering theory for the Dirac operator (e.g., Birman [3], Titchmarsh [4], Prosser [5], Roze [6], Evans [7], Thompson [8], Mochizuki [9], Eckardt [10], [11]). Prosser [5] shows that the wave operator

$$W_{\pm} = s\text{-}\lim_{t \rightarrow \pm \infty} \exp(itH) \exp(-itH_0)$$

exists under the main assumption

$$(4) \quad |Q(x)| = O(|x|^{-1-h}) \quad (h > 0)$$

at infinity, and that the scattering operator

$$S = W_{+}^{*} W_{-}$$

is unitary for a class of potentials with compact support. Eckardt [10] proves the existence of wave operators under a weaker condition

$$|Q(x)| (1 + |x|)^{-1/2 + \delta} \in L^2(\mathbb{R}^3) \quad (\delta > 0).$$

In [12] and [13] we assume (4) and

$$Q(x) \in \mathcal{B}^1(\mathbb{R}^3)$$

(i.e., every component of $Q(x)$ is bounded and has bounded continuous first derivatives). We can assume some local singularities of $Q(x)$, but for the sake of simplicity we omit them. In [12] we show that the limiting absorption principle holds on $[-1, 1]^c$. The limiting absorption principle is, roughly speaking, to investigate the resolvent of H

near the spectrum. Let $R(z) = (H - z)^{-1}$ be the resolvent of H for non-real z . As z tends to the spectrum, the limit of $R(z) f$ for $f \in \mathcal{L}^2$ does not exist generally in $\mathcal{L}^2$. The limit of $R(z) f$, however, exists for appropriate functions $f(x)$ in some weighted Hilbert spaces (the method is called the limiting absorption principle). We introduce two weighted functional spaces

$$\begin{aligned} \mathcal{L}_t^2 &= \left\{ u ; \int_{\mathbb{R}^3} (1 + |x|)^{2t} |u(x)|^2 dx < +\infty \right\}, \\ \mathcal{H}_{-t}^1 &= \left\{ u ; \int_{\mathbb{R}^3} (1 + |x|)^{-2t} (|u(x)|^2 + \sum_{j=1}^3 \left| \frac{\partial}{\partial x_j} u(x) \right|^2) dx \right. \\ &\quad \left. < +\infty \right\}. \end{aligned}$$

Theorem 1. (the limiting absorption principle) Let $t > 1/2$.

Then for every real λ such that $|\lambda| > 1$, there exist bounded operators $R^+(\lambda)$, $R^-(\lambda)$ on $\mathcal{H}_{-t}^1$ to $\mathcal{L}_t^2$ such that

$$\begin{aligned} \text{s-lim}_{z \rightarrow \lambda \pm 0i} R(z) f &= R_{\pm}^+(\lambda) f \quad \text{in } \mathcal{H}_{-t}^1 \end{aligned}$$

for $f \in \mathcal{L}_t^2$. For every $f \in \mathcal{L}_t^2$, $R(z) f$ is strongly continuous

in the topology of $\mathcal{H}_{-t}^1$ with respect to z with the boundary values

$$R^+(\lambda) f, R^-(\lambda) f.$$

The following assertion follows directly from Theorem 1.

Corollary 1 . $[-1, 1]^c$ is absolutely continuous spectrum of H .

In [13] we see eigenfunction expansions and scattering theory under the same condition as in [12].

We shall summarize the results in [13].

There exist 4×4 matrix-valued functions $\Phi_{\nu}^{\pm}(x, r)$ ($\nu = p, n$) for $x \in \mathbb{R}^3$ and $r > 0$. Every component of $\Phi_{\nu}^{\pm}(x, r)$ is a $L^2(S)$ -valued function, locally Hölder continuous in $L^2(S)$ with respect to x and locally bounded in $L^2(S)$ with respect to r (S is the unit surface about the origin). $\Phi_{\nu}^{\pm}(x, r)$ might be called generalized eigenfunctions in the following sense :

$$(L_0 + Q(x)) \int_S (\Phi_{\nu}^{\pm}(x, r))(\omega) h(\omega) d\omega = \tau_{\nu} \sqrt{r^2 + 1} \int_S (\Phi_{\nu}^{\pm}(x, r))(\omega) h(\omega) d\omega$$

for any $h \in \mathcal{L}^2(S) = (L^2(S))^4$, where $\tau_p = 1$, $\tau_n = -1$. Let

$$(Z_{\nu}^{\pm} f)(r \cdot) = (2\pi)^{-3/2} \text{ l.i.m. } \int_{\mathbb{R}^3} \Phi_{\nu}^{\pm}(x, r)^* f(x) dx$$

for $f \in \mathcal{L}^2$. Then $Z_{\nu}^{\pm}$ is a partially isometric operator in $\mathcal{L}^2$ with the initial set $(I - E(1)) \mathcal{L}^2$ ($\nu = p$), $E(-1-0) \mathcal{L}^2$ ($\nu = n$)

For $f \in \mathcal{L}^2$

- 1) $E(\cdot)$ is the right-continuous resolution of the identity associated with H .

$$\|f\|^2 = \|Z_p^+ f\|^2 + \|Z_n^+ f\|^2 + \sum_j |(f, \varphi_j)|^2,$$

where $\{\varphi_j\}$ is the set of the orthonormalized eigenfunctions for the discrete eigenvalues in $[-1, 1]$ (it may be empty). We can construct the stationary wave operator

$$U_{\pm} f = (Z_p^+)^*(\hat{f}) + (Z_n^+)^*(\hat{f})$$

isometric from $\mathcal{L}^2$ onto $(I - E(1) + E(-1-0))\mathcal{L}^2$, where $\hat{f}$ is the Fourier image of f . Then we have

$$U_{\pm} = W_{\pm}$$

(that is, the above stationary wave operator coincides with the time-dependent wave operator $W_{\pm}$), and that the scattering operator $S = W_+^* W_-$ is unitary.

We shall now give the result for the long range potential. The potential $Q(x)$ is assumed to satisfy the following condition (A):

- (A) each component of $Q(x)$ is continuously differentiable except at a finite number of singularities, satisfying (2) or (3), and

$$(4) \quad |Q(x)| = O(|x|^{-\delta})$$

$$(5) \quad \sum_{j=1}^3 \left| \frac{\partial}{\partial x_j} Q(x) \right| = O(|x|^{-1-\delta})$$

at infinity for some $\delta > 0$.

Theorem 2. Let $Q(x)$ satisfy the condition (A). Then the number of eigenvalues of $H = H_0 + Q$ is, if it exists, is at most finite in $[-1-\delta, 1+\delta]^c$ for every $\delta > 0$. $\{\lambda_n\}$ denotes the set of eigenvalues in $[-1, 1]^c$ (it may be empty). Then each λ_n is of finite multiplicity. $[-1, 1]^c - \{\lambda_n\}$ is the absolutely continuous spectrum of $H = H_0 + Q$.

Proof. Let us take any $f \in \mathcal{H}_t^1$ ($t > 1/2$) and non-real z . Then $u = (H - z)^{-1} f$ fulfills

$$(L_0 + Q(x)) u(x) - z u(x) = f(x)$$

and, by $L_0^2 = (-\Delta + 1) I$ (which is easily checked by the anti-commutation relation of α_j),

$$\begin{aligned} -\Delta u(x) + L_0 (Q(x) u(x)) + z Q(x) u(x) - (z^2 - 1) u(x) \\ = L_0 f(x) + z f(x). \end{aligned}$$

In view of this fact we notice that a result on Schrödinger operators with long range potentials, obtained by Ikebe-Saito [14], will be applicable to our assertion. The above theorem is proved along the same line of Ikebe-Saito [14].

A sufficient condition for the non-existence of the eigenvalues in $[-1, 1]^c$ is given as follows.

Theorem 3. Assume that

$$\begin{aligned} L &= L_0 + Q(x) \\ &\equiv -i \sum_{j=1}^3 \alpha_j \left(\frac{\partial}{\partial x_j} + i A_j(x) \right) + (\beta + q(x) I) \\ &\quad \left(Q(x) \equiv \sum_j A_j(x) \alpha_j + q(x) I \right) \end{aligned}$$

such that the scalar potential $A_j(x)$ and $q(x)$ belong to $C^1(\mathbb{R}^3 - \mathcal{E})$

($\mathcal{E}$ is a set of a finite number of points), satisfying

$$\begin{aligned} \sum_{j=1}^3 \left(|A_j(x)| + |x| |\operatorname{grad} A_j(x)| \right) \\ + |q(x)| + |x| |\operatorname{grad} q(x)| = o(1) \quad (|x| \rightarrow \infty). \end{aligned}$$

Then the selfadjoint extension H has no $\mathcal{L}^2$ -eigenfunctions in $[-1, 1]^c$.

Proof. Let $Hu = \lambda u$ ($|\lambda| > 1$, $u \in \mathcal{L}^2$). Then u is a solution of a Schrödinger equation

$$\begin{aligned} -\Delta u(x) - 2i \sum_{j=1}^3 A_j(x) \frac{\partial u}{\partial x_j} - i \sum_{j=1}^3 \frac{\partial A_j}{\partial x_j} \alpha_j u(x) \\ - i \sum_{j,k=1}^3 \frac{\partial A_j}{\partial x_k} \alpha_k \alpha_j u(x) + \left(\sum_{j=1}^3 A_j(x)^2 + 2\lambda q(x) \right. \\ \left. - q(x)^2 \right) u(x) = (\lambda^2 - 1) u(x). \end{aligned}$$

Then we can show $u = 0$ by the method in Ikebe-Saito [14], Remark on the proof of Lemma 2.5.

Remark 2. When the potential $Q(x)$ is a spherically symmetric scalar function, spectral problems for Dirac operators frequently reduces, by separation of variables, to investigate 2×2 differential operators

$$h_k = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \frac{d}{dr} + \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} - \frac{k}{r} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + q(r) \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix},$$

$$0 < r < \infty \quad (k = \pm 1, \pm 2, \pm 3, \dots),$$

where $r = |x|$ and $q(r) = Q(x)$ (see, e.g., Dirac [15]). The operator h_k is also studied by many authors (e.g., Titchmarsh [16], Weidmann [17]). Weidmann [16] shows that every selfadjoint realization A_k of h_k has the essential spectrum $(-1, 1)^c$, and that the spectrum of A_k is absolutely continuous in $[-1, 1]^c$, when $q(r) = \frac{c}{r}$ (c is an arbitrary real number).

References

- [1] K. Jörgens, Perturbations of the Dirac operator, Conference on the theory of ordinary and partial differential equations, Lecture notes of math., 280 (1972), 87-102 (Springer).
- [2] M. Arai, On essential selfadjointness of Dirac operators, the present notes.
- [3] M. S. Birman, On the spectrum of singular boundary problems, Math. Sb., 55 (1961), 126-174 (Amer. Math. Soc. Trans., 53).
- [4] E. C. Titchmarsh, On the completeness problem for the eigenfunction formulae of relativistic quantum mechanics, Proc. Royal Soc. A, 262 (1961), 489-502.
- [5] R. T. Prosser, Relativistic potential scattering, J. Math. Physics, 4(1963), 1048-1054.
- [6] S. N. Roze, On the character of spectrum of the Dirac operator, Theoretical and Math. Physics, 2 (1970), 377-382 (in Russian).
- [7] W. D. Evans, On the unique self-adjoint extension of the Dirac operator and the existence of the Green matrix, Proc. London Math. Soc., 20 (1970), 537-557.
- [8] M. Thompson, Eigenfunction expansions and the associated scattering

- theory for potential perturbations of the Dirac equation, Quart. J. Math. Oxford, 23 (1972), 17-55.
- [9] K. Mochizuki, On the perturbation of the continuous spectrum of the Dirac operator, Proc. Japan Acad., 40 (1964), 707-712.
- [10] K. -J. Eckardt, On the existence of wave operators for Dirac operators, Manuscripta math. 11 (1974), 359-371.
- [11] K. -J. Eckardt, Scattering theory for Dirac operators, Math. Z., 139 (1974), 105-131.
- [12] O. Yamada, On the principle of limiting absorption for the Dirac operator, Publ. RIMS, Kyoto Univ., 8 (1972/3), 557-577.
- [13] O. Yamada, Eigenfunction expansions and scattering theory for Dirac operators (to be published).
- [14] T. Ikebe and Y. Saito, Limiting absorption method and absolute continuity for the Schrödinger operator, J. Math. Kyoto Univ., 12 (1972), 513-542.
- [15] P. A. M. Dirac, Principle of quantum mechanics, Oxford, 1958.
- [16] E. C. Titchmarsh, On the nature of the spectrum in problems of quantum mechanics, Quart. J. Math. Oxford(2), 12 (1961), 227-240.
- [17] J. Weidmann, Oszillationsmethoden für Systeme gewöhnlicher Differentialgleichungen, Math. Z., 119 (1971), 349-373.