

Uniqueness of prime models in definably complete locally o-minimal theories

小峰 航 (Wataru Komine)

筑波大学 理工情報生命学術院 数理物質科学研究群
(Degree Programs in Pure and Applied Sciences,
Graduate School of Science and Technology,
University of Tsukuba)

Abstract

It is well-known that every o-minimal theory has a prime model and it is unique up to isomorphism. We give sufficient conditions for uniqueness of prime models in definably complete locally o-minimal theories.

1 Introduction

We are interested in definably complete locally o-minimal structures and its theories. Pillay and Steinhorn showed the following:

Theorem 1 ([1]). *Every o-minimal theory has prime model over any sets, that are unique up to isomorphism.*

In this note, we give a condition for uniqueness of prime models in definably complete locally o-minimal theories. This condition is related to form of definable sets.

This work was supported by JST, the establishment of university fellowships towards the creation of science technology innovation, Grant Number JPMJFS2106.

2 Preliminaries

First, we define o-minimality, definable completeness and local o-minimality. Let a structure $\mathcal{M} = (M, <, \dots)$ be expansion of a linear ordered set without endpoints.

Definition 2. $\mathcal{M}$ is o-minimal structure if every definable subset of M is a finite union of points and open intervals. $\mathcal{M}$ is locally o-minimal structure if for any points $a \in M$ and definable subset A of M , there is an open interval I containing a such that $A \cap I$ is a finite union of points and open intervals. $\mathcal{M}$ is definably complete if every definable subset of M has supremum and infimum in $M \cup \{-\infty, +\infty\}$.

Example 3. Dense linear ordered sets without endpoints, real closed fields and $(\mathbb{R}, 0, 1, +, -, \cdot, \exp(-))$ are o-minimal. $(\mathbb{R}, <, 0, +, \sin(-))$ is not o-minimal structure since discrete infinite subset $\mathbb{Z} = \{x \in \mathbb{R} \mid \sin(x\pi) = 0\}$ is definable. But it is definably complete locally o-minimal structure.

In definably complete locally o-minimal structure, an analogy of monotonicity theorem holds as following (it is called strong local monotonicity theorem).

Theorem 4 (strong local monotonicity theorem, [2]). *Let I be a A -interval and $f : I \rightarrow M$ be a A -definable function. Then there exists a mutually disjoint A -definable partition $I = X_d \cup X_c \cup X_+ \cup X_-$ satisfying the following conditions:*

- (1) *the A -definable set X_d is discrete and closed;*
- (2) *the A -definable set X_c is open and f is locally constant on X_c ;*
- (3) *the A -definable set X_+ is open and f is locally strictly increasing and continuous on X_+ ;*
- (4) *the A -definable set X_- is open and f is locally strictly decreasing and continuous on X_- .*

In this note, we assume a language $\mathcal{L}$ contains $<$ and an interpretation of $<$ is linear ordering without endpoints in $\mathcal{L}$ -structures.

Definition 5. $\mathcal{L}$ -theory T is o-minimal (resp. definably complete locally o-minimal) if every model of T is o-minimal (resp. definably complete locally o-minimal) structure.

Remark 6. If $\mathcal{M}$ is o-minimal, its complete theory $\text{Th}(\mathcal{M})$ is o-minimal by uniform finiteness theorem. On the other hand, definable completeness and local o-minimality are represented by $\mathcal{L}$ -sentences. Therefore, if $\mathcal{M}$ is definably complete locally o-minimal, its complete theory $\text{Th}(\mathcal{M})$ is definably complete locally o-minimal.

In this note, we assume $\mathcal{L}$ -theory T is complete and work in monster model $\mathbb{M}$. On existence and uniqueness of prime models, the following fact is important.

Fact 7 ([1][4]). *Let a complete theory T be given.*

- (1) *Let $A \subset \mathcal{M} \models T$. Any model $\mathcal{M}$ that is constructible over A also is prime over A .*
- (2) *Suppose that for any subset A of a model $\mathcal{M}$ of T and any formula φ having parameters A , there is a complete formula with parameters from A which, relative to $\text{Th}(\mathcal{M}, a)_{a \in A}$, implies φ . Then, for any $A \subset \mathcal{M}$, there is a model $\mathcal{M}$ of T that is constructible over A . (it is enough to consider when φ has just one free variable)*
- (3) *Let $A \subset \mathcal{M} \models T$. Then any two models that are constructible over A are isomorphic over A .*

3 Main theorem

In this section, let T be a definably complete locally o-minimal theory. By using strong local monotonicity theorem (Theorem 4), We can show the following theorem with a similar argument to o-minimal case. (cf.[1])

Theorem 8. *If $\text{dcl}(\emptyset)$ is nonempty, then T has prime model over emptyset and it is unique up to isomorphism. Moreover, T has prime model over nonempty A and it is unique up to isomorphism.*

We showed the following theorem that is related to complete formulas.

Theorem 9. *We assume that T has a complete formula. Then, T has a prime model over $\emptyset$ and it is unique up to isomorphism.*

The following is the main theorem.

Theorem 10. *We assume that there is a prime model $\mathcal{M}$ of T such that it has a discrete definable subset X of M without parameters which is isomorphic to $n \times \mathbb{Z}$ as ordered sets for some positive integer n . Then prime models of T are isomorphic.*

sketch of proof. For convenience, we assume $n = 2$.

We take $\mathcal{L}$ -formula $\varphi(v)$ which define X . Since X is isomorphic to $\mathbb{Z} \sqcup \mathbb{Z}$ as ordered sets, we get decomposition $X = X_0 \sqcup X_1$ where X_0 corresponds to first part of $\mathbb{Z} \sqcup \mathbb{Z}$ and X_1 corresponds to second part of $\mathbb{Z} \sqcup \mathbb{Z}$.

We take a element $a \in X_0$ from first part. Let $\mathcal{N}$ be a constructible model over a and $X' = \varphi(\mathcal{N})$. Then we get decomposition $X' = X'_0 \sqcup X'_1$ where each X'_i isomorphic to $\mathbb{Z}$. Since $\mathcal{M}$ is prime, we get an elementary embedding $j : \mathcal{M} \hookrightarrow \mathcal{N}$. So we get decomposition $X' = j(X_0) \sqcup j(X_1)$. Then $j(X_0) = X'_0$ and $j(X_1) = X'_1$. In $\mathcal{N}$, $S^m(a) = j(a)$ for some integer m where S is successor function in X' . Since $j(a)$ is definable over a , $\mathcal{N}$ is prime over $j(a)$. Then $\mathcal{M}$ is prime over a . By uniqueness of prime models over nonemptyset, $\mathcal{M}$ and $\mathcal{N}$ are isomorphic. $\square$

References

- [1] Pillay Anand, and Charles Steinhorn, Definable sets in ordered structures. I., Transactions of the American Mathematical Society 295.2 (1986): 565-592.
- [2] Fujita Masato, Kawakami Tomohiro, Komine Wataru, Tameness of definably complete locally o-minimal structures and definable bounded multiplication, Mathematical Logic Quarterly, Volume 68, Issue 4, November 2022, Pages 496-515.
- [3] Lou van den Dries, Tame Topology and O-minimal Structures, Cambridge University Press (1998)
- [4] David Marker, Model Theory: An Introduction, Springer (2002)