

JORDAN-HÖLDER TYPE THEOREM IN NORMAL INTERMEDIATE SUBFACTOR LATTICES FOR DEPTH TWO INCLUSIONS OF AFD II_1 FACTORS

TAMOTSU TERUYA (照屋 保)

ABSTRACT. Let $N \subset M$ be a depth 2 inclusion of AFD II_1 factors with finite Jones index. Let K and L be normal intermediate subfactors of $N \subset M$. If $K \cap L = N$ and M is generated by K and L , then we can represent M, K, L, N as $M = P \otimes R, K = Q \otimes R, L = P \otimes S$, and $N = Q \otimes S$ for some inclusions $P \supset Q$ and $R \supset S$. Using this characterization, we shall prove Jordan-Hölder type theorem in normal intermediate subfactor lattices for depth 2 inclusions of AFD II_1 factors.

1. INTRODUCTION

Let $N \subset M$ be an irreducible inclusion of type II_1 factors with finite index. In [9], the author introduced the notion of normality for intermediate subfactors of $N \subset M$ as follows:

Definition 1.1. Let K be an intermediate subfactor of the inclusion $N \subset M$. Let $N \subset M \subset M_1 \subset M_2$ be the Jones tower for $N \subset M$ and K_1 the basic extension for $K \subset M$. Then K is a *normal intermediate subfactor* of the inclusion $N \subset M$ if $e_K \in \mathcal{Z}(N' \cap M_1)$ and $e_{K_1} \in \mathcal{Z}(M' \cap M_2)$, where e_K and e_{K_1} are the Jones projections for $K \subset M$ and $K_1 \subset M_1$, respectively.

TAMOTSU TERUYA

With the above notation, if the depth of $N \subset M$ is 2, then $N' \cap M_1$ and $M' \cap M_2$ are a dual pair of Hopf C^* -algebras. and $K' \cap K_1$ is a $*$ -subalgebra and a left coideal of $N' \cap M_1$ (see [1]). Then K is a normal intermediate subfactor of $N \subset M$ if and only if $K' \cap K_1$ is a subHopf algebra and the left and right adjoint action of $N' \cap M_1$ leave $K' \cap K_1$ invariant (see [3]).

Watatani[10] studied intermediate subfactor lattices $\mathcal{L}(N \subset M)$ and relations between modular identity and commuting and co-commuting (nondegenerate) square conditions. The author[9] proved if the depth of $N \subset M$ is 2, then the set $\mathcal{N}(N \subset M)$ of all normal intermediate subfactors of $N \subset M$ is a sublattice of $\mathcal{L}(N \subset M)$ and a modular lattice.

Let $N \subset M$ be an irreducible, depth 2 inclusion of AFD II_1 factors with finite index. Our purpose is to show Jordan-Hölder type theorem in normal intermediate subfactor lattices for $N \subset M$. To be more precise, we prove that if $M = A_0 \supset A_1 \supset A_2 \supset \cdots \supset A_n = N$ and $M = B_0 \supset B_1 \supset B_2 \supset \cdots \supset B_m = N$ are maximal chains of $\mathcal{N}(N \subset M)$, then $m = n$ and the inclusions $A_{i-1} \supset A_i$ are isomorphic to the inclusions $B_{j-1} \supset B_j$ in some order. To show this, we characterize tensor products of depth 2 inclusions of AFD II_1 factors with finite index as follows: Let $N \subset M$ be an irreducible, depth 2 inclusion of AFD II_1 factors with finite index. Let K and L be normal intermediate subfactors for $N \subset M$. If $K \cap L = N$ and M is generated by K and L , then we can represent M, K, L, N as $M = P \otimes R, K = Q \otimes R, L = P \otimes S$ and $N = Q \otimes S$ for some inclusions $P \supset Q$ and $R \supset S$.

JORDAN-HÖLDER TYPE THEOREM

2. A CHARACTERIZATION OF TENSOR PRODUCTS OF DEPTH 2 INCLUSIONS

Let $N \subset M$ be an irreducible, depth 2 inclusion of II_1 factors with $[M : N] < \infty$ and $\mathcal{N}(N \subset M)$ the all normal intermediate subfactors of $N \subset M$. Suppose that $K, L \in \mathcal{N}(N \subset M)$ and M is generated by K and L , and $N = K \cap L$. Then

$$\begin{array}{ccc} K & \subset & M \\ \cup & & \cup \\ N & \subset & L \end{array}$$

is commuting and co-commuting (nondegenerate) square (see [6, 8]). Let $K_1 = \langle K, e_K^M \rangle$ and $L_1 = \langle L, e_L^M \rangle$ be the basic extension with the Jones projections e_K^M and e_L^M for $K \subset M$ and $L \subset M$, respectively. Then it is well known that

$$\begin{array}{ccc} M & \subset & K_1 \\ \cup & & \cup \\ L & \subset & \langle L, e_K^M \rangle \end{array} \quad \text{and} \quad \begin{array}{ccc} M & \subset & L_1 \\ \cup & & \cup \\ K & \subset & \langle K, e_L^M \rangle \end{array}$$

are also nondegenerate commuting squares.

Lemma 2.1. *With the above notation, $L \subset K_1$ and $K \subset L_1$ are irreducible, depth 2 inclusions. Moreover, M and $\langle L, e_K^M \rangle$ are normal intermediate subfactors of $L \subset K_1$ and, M and $\langle K, e_L^M \rangle$ are normal intermediate subfactors of $K \subset L_1$*

TAMOTSU TERUYA

Proof. Since $L \subset M$ and $M \subset K_1$ are depth 2 inclusion by [9], the depth of $L \subset K_1$ is 2 by [7]. Similarly, $K \subset L_1$ is a depth 2 inclusion. It is easy to see that $L \subset K_1$ and $K \subset L_1$ are irreducible inclusions. $\square$

Lemma 2.2. *With the above notation, we have*

$$K' \cap K_1 = \langle K, e_L^M \rangle' \cap M_1 = N' \cap \langle L, e_K^M \rangle$$

$$L' \cap L_1 = \langle L, e_K^M \rangle' \cap M_1 = N' \cap \langle K, e_L^M \rangle.$$

Proof. By Lemma 2.1 and [8], we have $[M : K] = [L : N] = [L_1 : \langle K, e_L^M \rangle]$. Therefore we have

$$\dim_{\mathbb{C}}(K' \cap K_1) = \dim_{\mathbb{C}}(\langle K, e_L^M \rangle' \cap M_1) = \dim_{\mathbb{C}}(N' \cap \langle L, e_K^M \rangle).$$

Let x be an element of $K' \cap K_1$. Since e_L^M is an element of the center of $N' \cap M_1$ and $K' \cap K_1 \subset N' \cap M_1$, x and e_L^M are commutative and hence $x \in \langle K, e_L^M \rangle' \cap M_1$. So we have $K' \cap K_1 \subset \langle K, e_L^M \rangle' \cap M_1$. By $\dim_{\mathbb{C}}(K' \cap K_1) = \dim_{\mathbb{C}}(\langle K, e_L^M \rangle' \cap M_1)$, we have $K' \cap K_1 = \langle K, e_L^M \rangle' \cap M_1$.

Since M_1 is the basic extension of K_1 by $\langle L, e_K^M \rangle$ with the Jones projection e_L^M , we have $\langle L, e_K^M \rangle = \{e_L^M\}' \cap K_1$. Since e_L^M is an element of the center of $N' \cap M_1 (\supset K' \cap K_1)$, if x is an element of $K' \cap K_1$, then $x \in \{e_L^M\}' \cap K_1 = N' \cap \langle L, e_K^M \rangle$. And hence $K' \cap K_1 \subset N' \cap \langle L, e_K^M \rangle$. And $K' \cap K_1 = N' \cap \langle L, e_K^M \rangle$ by $\dim_{\mathbb{C}}(K' \cap K_1) = \dim_{\mathbb{C}}(N' \cap \langle L, e_K^M \rangle)$. Similarly, we have $L' \cap L_1 = \langle L, e_K^M \rangle' \cap M_1 = N' \cap \langle K, e_L^M \rangle$. $\square$

JORDAN-HÖLDER TYPE THEOREM

Theorem 2.3. *Let $N \subset M$ be an irreducible, depth 2 inclusion of AFD II_1 factors with $[M : N] < \infty$. If K and L are normal intermediate subfactors of $N \subset M$ such that $K \cap L = N$ and M is generated by K and L , then we can represent M, N, K, L as $M = P \otimes R, N = Q \otimes S, K = Q \otimes R$ and $L = P \otimes S$*

Proof. $N \subset M$ has the generating property, i.e., there exists a tunnel $M = N_0 \supset N = N_1 \supset N_2 \supset \cdots \supset N_i \supset \cdots$ such that

$$M = \overline{\bigcup_{i=1}^{\infty} (M \cap N_i)}^{weak} \supset N = \overline{\bigcup_{i=1}^{\infty} (N \cap N_i)}^{weak}$$

(see for example [4, 5]). Let

$$A_{00} \supset A_{01} \supset A_{02} \supset \cdots$$

$$\cup \quad \cup \quad \cup$$

$$A_{10} \supset A_{11} \supset A_{12} \supset \cdots$$

$$\cup \quad \cup \quad \cup$$

$$A_{20} \supset A_{21} \supset A_{22} \supset \cdots$$

$$\cup \quad \cup \quad \cup$$

$$\vdots \quad \vdots \quad \vdots$$

TAMOTSU TERUYA

be the commuting and co-commuting squares such that the initial commuting square is

$$\begin{array}{ccc} M & \supset & L \\ \cup & & \cup \\ K & \supset & N \end{array}$$

and $A_{ii} = N_i$ for $i = 1, 2, \dots$ as in [8]. Note that for the square

$$\begin{array}{ccc} A_{kl} & \supset & A_{k,l+1} \\ \cup & & \cup \\ A_{k+1,l} & \supset & A_{k+1,l+1}, \end{array}$$

$A_{kl} \supset A_{k+1,l+1}$ is again irreducible, depth 2 and, $A_{k,l+1}$ and $A_{k+1,l}$ are normal intermediate subfactors of $A_{kl} \supset A_{k+1,l+1}$. We put

$$\begin{aligned} P &= \overline{\bigcup_{i=1}^{\infty} (A_{00} \cap A'_{i0})}^{weak} \supset Q = \overline{\bigcup_{i=1}^{\infty} (A_{10} \cap A'_{i0})}^{weak} \\ R &= \overline{\bigcup_{i=1}^{\infty} (A_{00} \cap A'_{0i})}^{weak} \supset S = \overline{\bigcup_{i=1}^{\infty} (A_{01} \cap A'_{0i})}^{weak}. \end{aligned}$$

Then we can see $M = P \otimes R$, $N = Q \otimes S$, $K = Q \otimes R$ and $L = P \otimes S$ by Lemma 2.2 and [2]. $\square$

3. JORDAN-HÖLDER TYPE THEOREM

In this section, we shall prove Jordan-Hölder type theorem for depth 2 inclusions of AFD II_1 factors.

JORDAN-HÖLDER TYPE THEOREM

Theorem 3.1. *Let $N \subset M$ be an irreducible, depth 2 inclusion of AFD II_1 factor.*

If K and L are normal intermediate subfactors of $N \subset M$, then $K \subset K \vee L$ and $K \cap L \subset L$ are conjugate.

Proof. Since the set $\mathcal{N}(N \subset M)$ of all normal intermediate subfactors of $N \subset M$ is a sublattice of $\mathcal{L}(N \subset M)$, $K \vee L$ and $K \cap L$ are elements of $\mathcal{N}(N \subset M)$. Therefore $N \subset K \vee L$ and $N \subset K \cap L$ are depth 2 inclusion by [9, Theorem 4.6]. Moreover $K \cap L$ is a normal intermediate subfactor of $N \subset K \vee L$ by [9, Proposition 3.7]. So we have $K \cap L \subset K \vee L$ is depth 2 inclusion by [9, Theorem 4.6]. By theorem 2.3, there exist inclusions $P \supset Q$ and $R \supset S$ such that $K \vee L = P \otimes R$, $K = P \otimes S$, $L = Q \otimes R$ and $K \cap L = Q \otimes S$. So we can see both $K \vee L \subset K$ and $L \supset K \cap L$ are conjugate to $R \subset S$. $\square$

Theorem 3.2. *Let $N \subset M$ be an irreducible, depth 2 inclusion of AFD II_1 factors with $[M : N] < \infty$. Let $K, \tilde{K}, L, \tilde{L}$ be normal intermediate subfactors of $N \subset M$ with $K \supset \tilde{K}$ and $L \supset \tilde{L}$. Then the pairs $\tilde{K} \vee (K \cap L) \supset \tilde{K} \vee (K \cap \tilde{L})$ and $\tilde{L} \vee (K \cap L) \supset \tilde{L} \vee (\tilde{K} \cap L)$ are conjugate.*

Proof. Since $\tilde{K} \vee (K \cap L) = (\tilde{K} \vee (K \cap \tilde{L})) \vee (K \cap L)$, the pairs $\tilde{K} \vee (K \cap L) \supset \tilde{K} \vee (K \cap \tilde{L})$ and $K \cap L \supset (K \cap L) \cap (\tilde{K} \vee (K \cap \tilde{L}))$ are conjugate by the previous theorem. Similarly, the pair $\tilde{L} \vee (K \cap L) \supset \tilde{L} \vee (\tilde{K} \cap L)$ and $K \cap L \supset (K \cap L) \cap (\tilde{L} \vee (\tilde{K} \cap L))$ are conjugate. Since $\mathcal{N}(N \subset M)$ is a modular lattice by [9], we have $(K \cap L) \cap (\tilde{K} \vee (K \cap \tilde{L})) = ((K \cap L) \cap \tilde{L}) \vee (K \cap \tilde{L}) = (K \cap \tilde{L}) \vee (K \cap \tilde{L})$. Similarly, we have

TAMOTSU TERUYA

$(K \cap L) \cap (\tilde{L} \vee (\tilde{K} \cap L)) = (K \cap \tilde{L}) \vee (K \cap \tilde{L})$. We have thus proved the theorem. $\square$

In a lattice L , a finite chain $x = x_0 \supseteq x_1 \supseteq \cdots \supseteq x_d = y$ is maximal if $x_i \not\supseteq x_{i+1}$ and $x_i \supseteq a \supseteq x_{i+1}$ implies $x = a$ or $x_{i+1} = a$ for $i = 1, 2, \dots, d-1$.

Theorem 3.3. *Let $N \subset M$ be an irreducible, depth 2 inclusion of AFD II_1 factors with $[M : N] < \infty$. If $M = A_0 \supset A_1 \supset \cdots \supset A_n = N$ and $M = B_0 \supset B_1 \supset \cdots \supset B_m$ are two maximal chain of $\mathcal{N}(N \subset M)$, then $m = n$ and the inclusions $A_{i-1} \supset A_i$ are isomorphic to the inclusions $B_{j-1} \supset B_j$ in some order.*

Proof. Put

$$A_{ij} = A_i \vee (A_{i-1} \cap B_j)$$

and

$$B_{ji} = B_j \vee (A_i \cap B_{j-1}).$$

Then $A_{i,j-1} \supset A_{ij}$ is isomorphic to $B_{j,i-1} \supset B_{ji}$ by Theorem 3.2. Since $A_0 \supset A_1 \supset \cdots \supset A_s$ is maximal chain, for any $i (i = 1, 2, \dots, s)$, there uniquely exists j such that $A_{i-1} = A_{i,j-1} \supset A_{ij} = A_i$. Then $B_{j-1} = B_{j,i-1} \not\supseteq B_{ji} = B_j$. And hence $A_{i-1} \supset A_i$ is isomorphic to $B_{j-1} \supset B_j$. $\square$

Example 3.4. Let G be a semi direct group $B \rtimes A$ of finite groups A and B . Let

$$M = P \rtimes_\gamma B \supset N = P^{(A,\gamma)} = \{x \in P \mid \gamma_a(x) = x, \forall a \in A\},$$

where γ is an outer action of G on II_1 factor P . Then the depth of $N \subset M$ is 2 (see for example [7]). Let $A_0 = A \supsetneq A_1 \supsetneq \cdots \supsetneq A_r = \{e\}$ and $B_0 = B \supsetneq B_1 \supsetneq$

JORDAN-HÖLDER TYPE THEOREM

$\cdots \supsetneq B_s = \{e\}$ be normal subgroups of G such that if H is a normal subgroup of G with $A_{i-1} \supsetneq H \supset A_i$ or $B_{j-1} \supsetneq H \supset B_j$, then $H = A_i$ or $H = B_j$. Then $M = P \rtimes_{\gamma} B_0 \supset P \rtimes_{\gamma} B_1 \supset \cdots P = P^{(A_r, \gamma)} \supset P^{(A_{r-1}, \gamma)} \supset \cdots P^{(A_0, \gamma)} = N$ is a maximal chain of $\mathcal{N}(N \subset M)$ by [9]. Therefore if $M = C_0 \supset C_1 \supset \cdots C_n = N$ a maximal chain of $\mathcal{N}(N \subset M)$, then $n = r + s$ and the inclusions $C_{k-1} \supset C_k$ are isomorphic to $R \rtimes F \supset P$ or $R \supset R^F$ for some II_1 factor and some finite group F .

REFERENCES

1. M. Izumi, R. Longo, and S. Popa. A Galois correspondence for compact groups of automorphisms of von Neumann algebras with a generalization to Kac algebras. preprint.
2. Y. Kawahigashi. Classification of paragroup actions on subfactors. *Publ. RIMS, Kyoto Univ.*, 31:481–517, 1995.
3. S. Montgomery. *Hopf Algebras and Their Actions on Rings*. CBMS series number 82, 1992.
4. A. Ocneanu. Quantum symmetry, differential geometry of finite graphs, and classification of subfactors, 1991. Univ. of Tokyo Seminary Notes.
5. S. Popa. Classification of subfactors: the reduction to commuting squares. *Invent. Math.*, 101:19–43, 1990.
6. S. Popa. Classification of amenable subfactors of type II. *Acta Math.*, 172:163–255, 1994.
7. T. Sano. Commuting co-commuting squares and finite dimensional Kac algebras. to appear in *Pacific. J. Math.*
8. T. Sano and Y. Watatani. Angles between two subfactors. *J. Operator Theory*, 32:209–242, 1994.
9. T. Teruya. Normal intermediate subfactors. *J. Math. Soc. Japan*, to appear.

TAMOTSU TERUYA

10. Y. Watatani. Lattices of intermediate subfactors. J. Funct. Anal., to appear.