

Introducing a metric on the space of fuzzy continuous mappings and the completeness of the space

Nagata Furukawa (Soka University)

We consider the space of the mappings which take their values in the set of fuzzy numbers, and introduce a metric on the space. We prove that the space constitutes a complete space under the metric.

A fuzzy number we treat in this paper is as follows.

Definition 1. A *fuzzy number* is a fuzzy set with a membership function $\mu: \mathbf{R} \rightarrow [0, 1]$ satisfying the following conditions:

- (i) there are real numbers a and b such that

$$\text{cl}\{t \in \mathbf{R} \mid \mu(t) > 0\} = [a, b],$$
- (ii) there exists a unique real number $m (a \leq m \leq b)$ such that $\mu(m) = 1$,
- (iii) $\mu(t)$ is upper semi-continuous on $[a, b]$,
- (iv) $\mu(t)$ is nondecreasing on $[a, m]$ and nonincreasing on $[m, b]$.

The set of all fuzzy numbers is denoted by $\mathbb{F}(\mathbf{R})$. Let ρ denote the Hausdorff distance among bounded closed intervals in $\mathbf{R}$. We introduce a distance on $\mathbb{F}(\mathbf{R})$ by the following:

Definition 2. For two fuzzy numbers $\tilde{a}$ and $\tilde{b}$ in $\mathbb{F}(\mathbf{R})$, the distance $d(\tilde{a}, \tilde{b})$ between $\tilde{a}$ and $\tilde{b}$ is defined by

$$d(\tilde{a}, \tilde{b}) = \sup_{\alpha \in [0, 1]} \rho(\tilde{a}_\alpha, \tilde{b}_\alpha),$$

where $\tilde{a}_\alpha$ and $\tilde{b}_\alpha$ denote the α -cuts of $\tilde{a}$ and $\tilde{b}$, respectively.

Definition 3. For $\varepsilon > 0$ and $\tilde{a} \in \mathbb{F}(\mathbf{R})$, two kinds of ε -neighborhoods of $\tilde{a}$ are defined by

$$B(\tilde{a}; \varepsilon) = \{\tilde{b} \in \mathbb{F}(\mathbf{R}) \mid d(\tilde{a}, \tilde{b}) < \varepsilon\},$$

$$\overline{B}(\tilde{a}; \varepsilon) = \{\tilde{b} \in \mathbb{F}(\mathbf{R}) \mid d(\tilde{a}, \tilde{b}) \leq \varepsilon\}.$$

Definition 4. Let $\tilde{a}$ and $\tilde{b}$ be two fuzzy numbers. Then

$$\tilde{a} \preceq \tilde{b} \text{ iff } \left(\sup \tilde{a}_\alpha \leq \sup \tilde{b}_\alpha \right) \& \left(\inf \tilde{a}_\alpha \leq \inf \tilde{b}_\alpha \right) \text{ for } \forall \alpha \in [0, 1],$$

and

$$\tilde{a} \prec \tilde{b} \text{ iff } \left(\sup \tilde{a}_\alpha < \sup \tilde{b}_\alpha \right) \& \left(\inf \tilde{a}_\alpha < \inf \tilde{b}_\alpha \right) \text{ for } \forall \alpha \in [0, 1].$$

Proposition 1. For $\varepsilon > 0$ and $\tilde{a} \in \mathbb{F}(\mathbf{R})$, it holds that

- (i) $\tilde{b} \in \overline{B}(\tilde{a}; \varepsilon) \Leftrightarrow \tilde{a} - \varepsilon \preceq \tilde{b} \preceq \tilde{a} + \varepsilon$,
- (ii) $\tilde{b} \in B(\tilde{a}; \varepsilon) \Rightarrow \tilde{a} - \varepsilon \prec \tilde{b} \prec \tilde{a} + \varepsilon$.

The condition (iv) in Definition 1 is sometimes exchanged by the following :

- (iv)' $\mu(t)$ is strictly increasing on $[a, m]$ and strictly decreasing on $[m, b]$.

Denote the set of all fuzzy sets satisfying (i), (ii), (iii) in Definition 1 and (iv)' by $\mathbb{F}'(\mathbf{R})$. For $\tilde{a} \in \mathbb{F}'(\mathbf{R})$, let

$$B'(\tilde{a}; \varepsilon) = \{\tilde{b} \in \mathbb{F}'(\mathbf{R}) \mid d(\tilde{a}, \tilde{b}) < \varepsilon\}.$$

Proposition 2. For $\varepsilon > 0$ and $\tilde{a} \in \mathbb{F}'(\mathbf{R})$, it holds that

$$\tilde{b} \in B'(\tilde{a}; \varepsilon) \Leftrightarrow \tilde{a} - \varepsilon \prec \tilde{b} \prec \tilde{a} + \varepsilon.$$

Proposition 3. For $\tilde{a} \in \mathbb{F}(\mathbf{R})$, let

$$i(\alpha) = \inf \tilde{a}_\alpha, \quad s(\alpha) = \sup \tilde{a}_\alpha, \quad \alpha \in [0, 1].$$

Then $i(\alpha)$ and $s(\alpha)$ are lower semi-continuous and upper semi-continuous on $[0, 1]$, respectively.

Proposition 4. Let X be a metric space. Let f_n ($n = 1, 2, \dots$) be a real-valued function defined on X . Suppose that the sequence $\{f_n\}$ converges uniformly to a function f defined on X . If, for each n , f_n is lower (resp. upper) semi-continuous on X , then f is lower (resp. upper) semi-continuous on X .

Theorem 1. $(\mathbb{F}(\mathbf{R}), d)$ is a complete metric space.

Definition 5. Let X be a metric space, and let $\tilde{f}$ a mapping from X to $\mathbb{F}(\mathbf{R})$. Let x be a point of X . Then, $\tilde{f}$ is said to be continuous at x , iff for every $\varepsilon > 0$, there exists a positive number $\delta = \delta(x)$ satisfying that

$$y \in S(x; \delta) \Rightarrow \tilde{f}(y) \in B(\tilde{f}(x); \varepsilon).$$

If $\tilde{f}$ is continuous at every x in X , then $\tilde{f}$ is said to be continuous on X .

Proposition 5. Every continuous mapping from a compact metric space X to $\mathbb{F}(\mathbf{R})$ is uniformly continuous on X .

Definition 6. Let X be a metric space. Denote the class of all continuous mappings from X to $\mathbb{F}(\mathbb{R})$ by $\mathcal{CF}[X]$. For two members $\tilde{f}$ and $\tilde{g}$ in $\mathcal{CF}[X]$, define the distance between $\tilde{f}$ and $\tilde{g}$ by

$$\delta(\tilde{f}, \tilde{g}) = \sup_{x \in X} d(\tilde{f}(x), \tilde{g}(x)).$$

Proposition 3. Let X be a compact metric space. Then, for every pair $(\tilde{f}, \tilde{g})$ of fuzzy mappings in $\mathcal{CF}[X]$, $\delta(\tilde{f}, \tilde{g})$ assumes a finite value and is represented by

$$\delta(\tilde{f}, \tilde{g}) = \max_{x \in X} d(\tilde{f}(x), \tilde{g}(x)).$$

Theorem 2. Let X be a compact metric space. Then $(\mathcal{CF}[X], \delta)$ is a complete metric space.

References

- [1] P. Diamond and P. Kloeden, Metric spaces of Fuzzy Sets, World Scientific, Singapore. (1994)
- [2] A. George and P. Veeramani, Some Theorems in Fuzzy Metric Spaces, The Journal of Fuzzy Mathematics, 3 (1995) 933-940.
- [3] Wang Geping, Distance Functions for Fuzzy Sets, The Journal of Fuzzy Mathematics, 3 (1995) 789-802.