

Computability problems of piecewise continuous functions

Mariko Yasugi and Yoshiki Tsujii
Faculty of Science, Kyoto Sangyo University
yasugi@cc.kyoto-su.ac.jp, tsujii@cc.kyoto-su.ac.jp

Main Subject : Interrelation between computability in the limit and computability in the uniform space ¹

1 Introduction

In studies of algorithm in analysis, one puts the basis of considerations on the computability of real numbers and the computability of continuous functions.

The notions of computability respectively of a real number (a sequence of real numbers) and of a continuous function (a sequence of continuous functions) are generally agreed to be natural and in a sense the strongest.

For a continuous function, computability means that there is a way to nicely approximate the values for computable inputs, and this notion depends on the continuity.

Very often, however, we compute values and draw a graph of a discontinuous function. We thus hope that some class of discontinuous functions can be attributed a certain kind of computability. In an attempt of computing a discontinuous function, a problem arises in the computation of the value at a jump point (a point of discontinuity). The problem is caused by the fact aht it is not generally decidable if a real number is a jump point, that is, the question " $x = a$?" is not decidable even for computable x and a .

One method of dissolving this problem was proposed in [10] by Pour-El and Richards. In their theory, a function is regarded as computable as a point in a function space. This is sufficient in order to draw a rough graph of the function, but does not supply us with information how to compute individual values.

There are many ways of characterizing computation of a discontinuous function. Here we will first report two of the approaches to this problem by Brattka, Mori, Tsujii, Washihara and Yasugi, and then claim that they are equivalent for piecewise continuous functions which jump at a computable sequence of real numbers, worked out by Tsujii and Yasugi [18]. One is to express the value

¹This work has been supported in part by JSPS Science Grant 12440031 and by Institute for Comprehensive Reserch, Kyoto Sangyo University (Yasugi Project 03).

of a function at a jump point in terms of limiting recursive functions instead of recursive functions ([16]). Another is to change the topology of the domain of a function ([12]).

References of related works and some other approaches are listed in **References**, details of which will not be mentioned here. Pour-El theory as well as its succeeding works on computability structures for Frechet spaces and metric spaces are also explained in [21].

2 Preliminaries

We first give some basic definitions. Details are seen in, for example, [10] and [21].

A sequence of rationals $\{r_n\}$ is recursive if it is represented as

$$r_n = (-1)^{\beta(n)} \frac{\gamma(n)}{\delta(n)}$$

with β, γ, δ recursive.

A real number x is *computable* (**R**-computable) if

$$\forall m \geq \alpha(p). |x - r_m| < \frac{1}{2^p}$$

for recursive α and $\{r_m\}$. We will write this relation as

$$x \simeq \langle r_m, \alpha \rangle.$$

These definitions can be extended to sequences of rationals and reals.

A real (continuous) function f is *computable* if (i) and (ii) below hold.

(i) f preserves sequential computability: If $\{x_n\}$ is computable, then $\{f(x_n)\}$ is computable.

(ii) f is continuous with a *recursive* modulus of continuity, say β :

$$\forall p \forall n \in \mathbf{N}^+ \forall k \geq \beta(n, p) \forall x, y \in [n, n+1],$$

$$|x - y| < \frac{1}{2^k} \Rightarrow |f(x) - f(y)| < \frac{1}{2^p}.$$

The definition can be extended to a computable sequence of functions.

The notions of computability of real numbers and continuous functions as above are generally agreed.

3 Points of discontinuity and various approaches

We would like to give some expression to a notion of computation of some discontinuous functions.


The problem that arises when computing the value of a discontinuous function is identifying a point of discontinuity, that is, $x = a$? is *not* decidable even for computable x and a .

There are various approaches to this problem. Our approach is a mathematical one, that is, using the mathematical language except for some elementary properties of recursive functions. Since we do not resort to any particular theory of computation, our approaches amounts to an axiomatic one.

What attracted us first was the functional approach proposed by Pour-El and Richards [10]. In their theory, a function is regarded as computable as a point in a space. Pour-El and Richards considered the computability structure on the Banach space and the Hilbert space. Succeeding them, Washihara investigated the computability structure on Frechét spaces, such as the spaces of the δ -function, the function space of bounded variation; Yoshikawa has studied the interpolation space, the Hilbert space [23] and others; Zhong has worked on the Solev space, to list a few.

In these theories, pointwise computation is not in sight. However, very often we need and indeed do pointwise computation of a function which is discontinuous here and there. How can we describe such deeds with mathematical terms? We have attempted two methods, one by the limiting recursive functions and one by changing the topology of the domain into a certain uniform topological space.

Below is a simplified map of the circumstances.



4 Topological computability

Definition 4.1 (Effective uniformity) Let X be a non-empty set. $\{V_n\}$ will denote an *effective* uniformity on X , that is, it is a map

$$V_n : X \rightarrow P(X)$$

and satisfies the following five axioms. ($\mathcal{T} = \langle X, \{V_n\} \rangle$ is then called an effective uniform topological space.)

$$A_1 \& A_2 : \cap_n V_n(x) = \{x\}.$$

There are recursive functions $\alpha_1, \alpha_2, \alpha_3$ such that

$$\forall n, m \in \mathbb{N} \forall x \in X. V_{\alpha_1(n,m)}(x) \subset V_n(x) \cap V_m(x) \quad (\text{effective } A_3).$$

$$\forall n \in \mathbb{N} \forall x, y \in X. x \in V_{\alpha_2(n)}(y) \rightarrow y \in V_n(x) \quad (\text{effective } A_4).$$

$$\forall n \in \mathbb{N} \forall x, y, z \in X. x \in V_{\alpha_3(n)}(y), y \in V_{\alpha_3(n)}(z) \rightarrow x \in V_n(z) \quad (\text{effective } A_5).$$

Definition 4.2 (Effective convergence) $\{x_k\} \subset X$ *effectively converges* to x in X if there is a recursive function γ satisfying $\forall x \forall k \geq \gamma(n)(x_k \in V_n(x))$.

The definition can be extended to effective convergence of a multiple sequence.

Definition 4.3 (Computability structure) (1) Let $\mathcal{S}$ be a family of sequences from X (Multiple sequences included).

$\mathcal{S}$ is called a *computability structure* if:

C1: (Non-emptiness) $\mathcal{S}$ is nonempty.

C2: (Re-enumeration) If $\{x_k\} \in \mathcal{S}$ and α is a recursive function, then $\{x_{\alpha(i)}\}_i \in \mathcal{S}$.

This condition can be extended to multiple sequences.

C3: (Limit) If $\{x_{lk}\}$ belongs to $\mathcal{S}$, $\{x_l\}$ is a sequence from X , and $\{x_{lk}\}$ converges to $\{x_l\}$ effectively, then $\{x_l\} \in \mathcal{S}$. ($\mathcal{S}$ is closed with respect to effective convergence.)

This condition can be extended to multiple sequences.

(2) A sequence belonging to $\mathcal{S}$ is called *computable*.

(3) x is *computable* if $\{x, x, \dots\}$ is computable.

(4) We write

$$\mathcal{C}_{\mathcal{T}} = \langle X, \{V_n\}, \alpha_1, \alpha_2, \alpha_3, \mathcal{S} \rangle.$$

Definition 4.4 (Effective approximation) $\{e_k\} \in \mathcal{S}$ is an *effective approximating set* of $\mathcal{S}$: $\forall \{x_l\}$ computable, there is a recursive function ν such that

$$\forall n \forall l (e_{\nu(n,l)} \in V_n(x_l)).$$

Definition 4.5 (Effective separability) Suppose $\{e_k\}$ is an effective approximating set and dense in X :

$$\forall n \forall x \exists k (e_k \in V_n(x))$$

Then $\mathcal{C}_{\mathcal{T}}$ is *effectively separable*, and $\{e_k\}$ is called an *effective separating set*.

Definition 4.6 (Relative effective completeness) (1) $\{x_j\} \in X$ is *effectively Cauchy* if

$$\forall n \forall j \geq \alpha(n) (x_j \in V_n(x_{\alpha(n)}))$$

for a recursive α .

This can be extended to multiple sequences.

(2) $\mathcal{E}_T = \langle X, \{V_n\}, \alpha_1, \alpha_2, \alpha_3, \mathcal{S} \rangle$ is *relatively effectively complete* (with respect to $\mathcal{S}$) if every computable and effectively Cauchy multiple sequence $\{x_{mj}\}$ effectively converges to a sequence $\{x_m\} \in X$.

Definition 4.7 (Relative computability) (1) $f : X \rightarrow \mathbf{R}$ is *relatively computable* (with respect to $\mathcal{S}$) if:

- (i) f preserves sequential computability:
 $\{x_m\}$ computable $\rightarrow \{f(x_m)\}$ computable
 - (ii) For any $\{x_m\} \in \mathcal{S}$ there exists a recursive function $\gamma(m, p)$ such that $y \in V_{\gamma(m, p)}(x_m)$ implies $|f(y) - f(x_m)| \leq \frac{1}{2^p}$.
- (2) can be extended to a sequence of functions.

Fact: f is relatively computable in the sense of (1) if and only if the sequence $\{f, f, f, \dots\}$ is relatively computable in the sense of (2).

Definition 4.8 (Computable function) (1) $f : X \rightarrow \mathbf{R}$ is *computable* if:

- (i) f preserves sequential computability.
- (ii) f is relatively computable, and there exist an effective approximating set, say $\{e_k\} \in \mathcal{S}$, and a recursive function $\gamma_0(k, p)$ for which

$$y \in V_{\gamma_0(k, p)}(e_k) \text{ implies } |f(y) - f(e_k)| \leq \frac{1}{2^p}$$

and

$$\bigcup_{k=1}^{\infty} V_{\gamma_0(k, p)}(e_k) = X$$

for p .

- (2) Can be extended to a sequence of functions.

Definition 4.9 (Uniform computability) f is *uniformly computable* if f preserves sequential computability and there is a recursive modulus of uniform continuity for f .

5 Computation in the limit

A limiting recursive function was first proposed by Gold [4]. It is defined to be the limit of a recursive function (if the limit exists).

Definition 5.1 (Limiting recursion: Gold) Let $r, s \geq 0$ and $g, \phi_1, \dots, \phi_r$ be recursive. Then the partial function h as defined below is called *limiting recursive*.

$$h(p_1, \dots, p_s) = \lim_n g(\tilde{\phi}_1(n), \dots, \tilde{\phi}_r(n), p_1, \dots, p_s, n)$$

where $\tilde{\phi}(n)$ is a code for

$$\langle \phi(0, p_1, \dots, p_s), \dots, \phi(n, p_1, \dots, p_s) \rangle,$$

Example

$$h(p_1, \dots, p_s) = \lim_n \phi(n, p_1, \dots, p_s).$$

Definition 5.2 (Computation by limit) A function $f : \mathbf{R} \rightarrow \mathbf{R}$ is said to be computed by limit if

$$x \simeq \langle r_p, \alpha \rangle \mapsto f(x) \simeq \langle s_p, \beta \rangle,$$

where $\{r_p\}, \alpha$ and $\{s_p\}$ are recursive, while β is limiting recursive.

6 Equivalence

We will set the following assumption.

[Assumption] $\{a_k\}_{k \in \mathbf{Z}}$, an $\mathbf{R}$ -computable sequence, is called a basic sequence if it holds

$$a_k < a_{k+1}, \cup_k [a_k, a_{k+1}] = \mathbf{R}.$$

Assume $a_k \simeq \langle v_{kp}, \gamma \rangle$, with $\{v_{kp}\}$ and γ recursive.

We will consider the computation of real functions relative to the basic sequence $\{a_k\}$.

Definition 6.1 ($\mathcal{A}$ -space) $A_k = \{a_k\}$, $J_k = (a_k, a_{k+1})$, $J = \cup_k J_k$,

$$A = \{a_k : k \in \mathbf{Z}\} = \cup_k A_k, \mathbf{A}_{\mathbf{R}} = A \cup J$$

(As a set, $\mathbf{A}_{\mathbf{R}} = \mathbf{R}$)

$$n = 1, 2, 3, \dots, x \in \mathbf{A}_{\mathbf{R}}$$

$$U_n(x) := \{x\} = \{a_k\} \text{ if } x \in A_k;$$

$$U_n(x) := \{y : y \in J_k, |x - y| < \frac{1}{2^n}\} \text{ if } x \in J_k$$

$$\mathcal{A} = \langle \mathbf{A}_{\mathbf{R}}, \{U_n\} \rangle$$

Corollary 1 $\{U_n\}$ is an effective uniformity on $\mathbf{A}_{\mathbf{R}}$

Definition 6.2 ($\mathcal{A}$ -computability)

$$\mathbf{A}_Q = J \cap Q$$

$k \in \mathbf{Z} \mapsto \iota_k$ (ι -symbol): a "symbolic name" for a_k

$$\mathbf{A}_Q^* = \mathbf{A}_Q \cup \cup_{k \in \mathbf{Z}} \{\iota_k\}$$

$\{q_{\mu n}\} \subset \mathbf{A}_Q^*$ is an $\mathcal{A}$ -sequence:

for each μ ,

$$\exists k \in \mathbf{Z}, q_{\mu n} = \iota_k \text{ for all } \mu \text{ or } \{q_{\mu n}\} \subset J_k$$

An $\mathcal{A}$ -recursive sequence: a recursive $\mathcal{A}$ -sequence

$$\{x_m\} \subset \mathbf{A}_{\mathbf{R}}$$

$$\{x_m\} \simeq_{\mathcal{A}} \langle q_{mn}, \alpha_A(m, p) \rangle \quad (*):$$




Figure 1: discrete sequence




Figure 2: approximation

$$\begin{aligned}
 &\{q_{mn}\} \subset J_k \text{ if } x_m \in J_k, \\
 &\{q_{mn}\} = \{\iota_k\} \text{ if } x_m \in A_k \\
 &\forall p \forall l \geq \alpha_A(m, p) (|x_m - q_{ml}|_A < \frac{1}{2^p}) \quad (**) \\
 &\text{if } x_m \in J_k (|a - b|_A = |a - b|, a, b \in J_k) \\
 &\{q_{mn}\} \text{ effectively } \mathcal{A}\text{-approximates } \{x_m\} \text{ with modulus of convergence } \alpha_A: \\
 &\{x_{im}\} \subset \mathbf{A}_R, \\
 &\{x_{im}\} \simeq_{\mathcal{A}} \langle q_{imn}, \alpha_A(i, m, p) \rangle \quad (*)
 \end{aligned}$$

$\{x_m\} \subset \mathbf{A}_R$ is $\mathcal{A}$ -computable if it is effectively approximated by a recursive $\mathcal{A}$ -sequence $\{q_{mn}\} \subset \mathcal{A}_Q^*$ and a recursive function $\alpha_A(m, p)$.

This can be extended to any multiple sequences.

A double sequence $\{x_{im}\} \subset \mathbf{A}_R$ is called $\mathcal{A}$ -computable if $\{q_{imn}\} \subset \mathbf{A}_Q^*$, $\alpha_A(i, m, n)$ are recursive

x is $\mathcal{A}$ -computable: $\{x, x, x, \dots\}$ is $\mathcal{A}$ -computable.

Proposition 1 (R- and $\mathcal{A}$ -computability) For a single real number x , x is $\mathbf{R}$ -computable if and only if x is $\mathcal{A}$ -computable.

Definition 6.3 (Limiting recursive sequence)

$$A(m, p) : |v_{k_m+1\gamma}(k_m+1, p) - r_{m\alpha(m, p)}| \leq \frac{4}{2^p}$$

$$B(m, p) : |v_{k_m+1\gamma}(k_m+1, p) - r_{m\alpha(m, p)}| > \frac{4}{2^p}$$

$$N_{mp} = 0 \quad \text{if } A(m, p)$$

$$N_{mp} = 1 \quad \text{if } B(m, p)$$

$$l_m := \lim_p \{N_{mp}\}_p: \text{limiting recursive.}$$

Theorem 1 (Relations between two notions of computability). (1) Suppose $x_m \simeq \langle r_{mn}, \alpha \rangle$ is an $\mathbf{R}$ -computable sequence.

Then we can construct an $\mathcal{A}$ -computable double sequence of real numbers, say $\{z_{mp}\}$, which converges to $\{x_m\}$ with a modulus of convergence ν which is "recursive in $\{l_m\}$."

(2) Suppose $\{x_m\}$ is an $\mathcal{A}$ -computable sequence of real numbers with $x_m \simeq_{\mathcal{A}} \langle q_{mn}, \alpha_A \rangle$. Then $\{x_m\}$ is $\mathbf{R}$ -computable.

Definition 6.4 (Sequential computability) (1) f is $\mathcal{L}$ -sequentially computable: for any $\mathbf{R}$ -computable $\{x_m\}$ ($x_m \simeq \langle r_{mn}, \alpha \rangle$), can construct a recursive sequence of rational numbers $\{s_{mp}\}$ and a function δ which is “recursive in $\{l_m\}$ ” such that $f(x_m) \simeq \langle s_{mp}, \delta \rangle$

(2) f is $\mathcal{A}$ -sequentially computable: for any $\mathcal{A}$ -computable sequence of real numbers $\{x_m\}$, we can construct a recursive sequence of rational numbers $\{s_{mp}\}$ and a recursive function β such that $f(x_m) \simeq \langle s_{mp}, \beta \rangle$.

Theorem 2 (Equivalence of two notions of sequential computability) (1) (From $\mathcal{L}$ to $\mathcal{A}$) An $\mathcal{L}$ -sequentially computable function f is $\mathcal{A}$ -sequentially computable.

(2) (From $\mathcal{A}$ to $\mathcal{L}$) An $\mathcal{A}$ -sequentially computable function f is $\mathcal{L}$ -sequentially computable.

Definition 6.5 (Piecewise computable function) $f : \mathbf{R} \rightarrow \mathbf{R}$ is called *piecewise computable* if the following hold.

- (i) For each ($\mathbf{R}$ -)computable real number x , $f(x)$ is $\mathbf{R}$ -computable.
- (ii) There is a recursive function κ with which, for any x, y such that $a_k < x, y < a_{k+1}$ and $|x - y| < \frac{1}{2^{\kappa(k,p)}}$, $|f(x) - f(y)| \leq \frac{1}{2^p}$.

Conclusion A piecewise continuous function whose jump points form a computable sequence of real numbers can be regarded as computable if it is piecewise computable and is $\mathcal{L}$ - (hence $\mathcal{A}$ -) sequentially computable.

References

- [1] L.Blum, M.Shub and S.Smale, On a theory of computation and complexity over the real numbers: *NP-completeness, recursive functions and universal machines*, Bulletin of American Mathematical Society, vol.21, no.1 (1989), 1-46.
- [2] V.Brattka, *Computing uniform bounds*, ENTCS 66-1(2002), 12pages; <http://www.elsevier.nl/locate/entcs/volume66.html>.
- [3] V.Brattka and K.Weihrauch, *Computability on subsets of Euclidean space I: Closed and compact subsets*, Theoretical Computer Science, 219 (1999), 65-93.
- [4] E.M.Gold, *Limiting recursion*, JSL, 30-1(1965), 28-48.
- [5] M.Nakata and S.Hayashi, *A limiting first order realizability interpretation*, SCMJ Online, vol.5(2001), 421-434.
- [6] S.Hayashi and M.Nakata, *Towards Limit Computable Mathematics*, International Workshop, Type 2000, Selected Papres, LNCS, 2277, Springer, 125-144, 2002.
- [7] T.Mori, *Computabilities of Fine continuous functions*, Computability and Complexity in Analysis, LNCS 2064, Springer, 200-221, 2001.

- [8] T.Mori, *On the computability of Walsh functions*, TCS 284(2002), 419-436.
- [9] T.Mori, Y.Tsujii and M.Yasugi, *Computability structure on metric spaces*, Combinatorics, Complexity and Logic (Proceedings of DMTCS'96), Springer(1996),351-362.
- [10] M.B.Pour-El and J.I.Richards, *Computability in Analysis and Physics*, Springer-Verlag(1989).
- [11] H.Tsuiki, *Real number computation through Gray code embedding*, TCS, vol.284, no.2(2002), 467-485.
- [12] Y.Tsujii, M.Yasugi and T.Mori, *Some properties of the effectively uniform topological space*, Computability and Complexity in Analysis, LNCS 2064, (2001),Springer.336-356.
- [13] M.Washihara, *Computability and Frechet spaces*, Math. Jap., vol.42(1995), 1-13.
- [14] M.Washihara, *Computability and tempered distribution*, to appear in Math.Jap..
- [15] M.Washihara and M.Yasugi, *Computability and metrics in a Frechet space*,Math. Jap., vol.43(1996), 431-443.
- [16] M.Yasugi, V.Brattka and M.Washihara, *Computability aspects of some discontinuous functions*, Scientiae Mathematicae Japonicae (SCMJ) Online, vol.5 (2001), 405-419.
- [17] M.Yasugi,T.Mori and Y.Tsujii, *Effective Properties of Sets and Functions in Metric Spaces with Computability Structure*, Theoretical Computer Science, 219(1999), 467-486.
- [18] M. Yasugi and Y.Tsujii, *Two notions of sequential computability of a function with jumps*, ENTCS 66 No.1(2002), 11 pages;
<http://www.elsevier.nl/locate/entcs/volume66.html>.
- [19] M. Yasugi and Y. Tsujii, *Computability of a function with jumps- Effective uniformity and limiting recursion-*, to appear in the Proceedings of International Conference on Topology in Matsue (TIM 2002), Topology and its Applications (Elsevier Science); seen also at
<http://www.kyoto-su.ac.jp/yasugi/page/recent.html>
- [20] M.Yasugi, Y.Tsujii and T.Mori, *Metrization of the uniform space and effective convergence*, Math. Log. Quart. 48(2002) Suppl. 1, 123-130.
- [21] M.Yasugi and M.Washihara, *Computability structures in analysis*, Sugaku Expositions (AMS) 13(2000),no.2,215-235.

- [22] M.Yasugi and M.Washihara, *A note on Rademacher functions and computability*, Words, Languages and Combinatorics III, World Scientific, 2003, 466-475.
- [23] A.Yoshikawa, *On an ad hoc computability structure in a Hilbert space*, Proceedings of the Japan Academy, vol.79,Ser.A,No.3(2003), 65-70.