

Knots and links in spatial graphs

谷山公規
Kouki Taniyama

〒 169-8050

東京都新宿区西早稲田 1-6-1

早稲田大学教育学部数学教室

Department of Mathematics, School of Education, Waseda University, 1-6-1

Nishi-Waseda, Shinjuku-ku, Tokyo, 169-8050, Japan

e-mail address: **taniyama@waseda.jp**

Abstract We study the relations of knots and links contained in a spatial graph.

This is an survey article on the results about knots and links contained in a spatial graph. We do not intend to cover all results in this topic. We only treat some of them here.

The set of knots and links contained in a spatial graph is a naive invariant of spatial graph. However it is of course not a complete invariant in general. For example Kinoshita's theta curve in Fig. 1 is not trivial but contains only trivial knots as the trivial theta curve. See for other such examples [5], [20] and [15].




Fig. 1

Anyway we are interested in the set of knots and links contained in a spatial graph. In [6] it is shown that any given $n(n-1)/2$ knot types are realized by an embedding of θ_n at once. Here θ_n denotes the graph on two vertices and n edges joining them. For example, suppose that trefoil knot, figure eight knot and $(2, 5)$ -torus knot are given. Then there is an embedding of $\theta = \theta_3$ that contains all of them. See Fig. 2 for such an example.




Fig. 2

Now we give a precise definition. Let G be a finite graph. We consider G as a topological space as well as a combinatorial object. Let Γ be a set of subgraphs of G . Suppose that for each $H \in \Gamma$, an embedding $\phi_H : H \rightarrow R^3$ is given. Then we say that the set of embeddings $\{\phi_H | H \in \Gamma\}$ is *realizable* if there is an embedding $\varphi : G \rightarrow R^3$ such that the restriction map $\varphi|_H$ is ambient isotopic to ϕ_H for each $H \in \Gamma$. The fundamental problem is whether or not given $\{\phi_H | H \in \Gamma\}$ is realizable.

Let $f : G \rightarrow R^3$ be an embedding. Then the Wu invariant $\mathcal{L}(f)$ of f is an element of an abelian group $L(G)$ associated to G . See [16] for their definitions. Let H be a subgraph of G . Then there is a natural homomorphism $h_H : L(G) \rightarrow L(H)$. Let I_G be a subset of $L(G)$ that is defined by $I_G = \{\mathcal{L}(f) | f : G \rightarrow R^3 \text{ is an embedding}\}$. Then the following is known in [17] as a necessary condition of realizability.

Theorem 1. *Suppose that $\{\phi_H | H \in \Gamma\}$ is realizable. Then there is an element $x \in I_G$ such that $h_H(x) = \mathcal{L}(\phi_H)$ for each $H \in \Gamma$.*

From now on we only consider the case that $\Gamma = \Gamma(G)$ is the set of all cycles of G . Here a *cycle* is a subgraph of G that is homeomorphic to a circle. A cycle on n vertices is called an n -cycle. Let $\Gamma_n(G)$ be the set of all n -cycles of G . We say that a graph

G is *adaptable* if any set of embeddings $\{\phi_H | H \in \Gamma(G)\}$ is realizable. Then the result stated above is rephrased that θ_n is adaptable. In [21] it is shown that K_4 is adaptable. Here K_n denotes the complete graph on n vertices. Moreover in [22] it is shown that all proper subgraphs of K_5 are adaptable. In [22] Yasuhara established a method of realization of knots and links in a spatial graph based on band description of knots. Now we are interested in whether or not K_5 is adaptable. The answer is ‘No’. In fact we have the following theorem.

Theorem 2. *A set of embeddings $\{\phi_H | H \in \Gamma(K_5)\}$ is realizable if and only if there is an integer m such that*

$$\sum_{H \in \Gamma_5(K_5)} a_2(\phi_H(H)) - \sum_{H \in \Gamma_4(K_5)} a_2(\phi_H(H)) = \frac{m(m-1)}{2}.$$

We note that the ‘only if’ part of Theorem 2 is shown in [8] and the ‘if’ part of Theorem 2 is shown in [19]. We refer the reader to [19], [12], [13] and [11] for related results.

Now we are interested in the existence of nontrivial knots and links in a large complete graph. The following theorem in [1] is a milestone in this area.

Theorem 3. (1) *For any embedding $f : K_6 \rightarrow R^3$ the sum of the linking numbers of the links in $f(K_6)$ is an odd number.*

(2) *For any embedding $f : K_7 \rightarrow R^3$ the sum of the second coefficients of the Conway polynomials of the knots of 7-cycles in $f(K_7)$ is an odd number.*

In [9] it is shown that for any knot J there is a natural number n such that every linear embedding of K_n into R^3 contains a cycle that is ambient isotopic to J . See also [7] [10] etc. for related results.

In [3] it is shown that every embedding of K_{10} into R^3 contains a 3-component non-splittable link. In [4] it is shown that for any natural number n there is a graph G such that every embedding of G into R^3 contains an n -component nonsplittable link.

In [2] it is shown that for any natural number n there is a natural number m such that every embedding of K_m contains a 2-component link whose absolute value of the linking number is greater than or equal to n . It is also shown in [2] that for any natural number n there is a natural number m such that every embedding of K_m contains a knot whose absolute value of the second coefficient of the Conway polynomial is greater than or equal to n . In the first result m is actually given by a polynomial of n whose degree is 2. In the second result m is actually given by a polynomial of n whose degree is 1. Recently the author and Shirai showed that in the first result m can be given by a polynomial of n whose degree is 1, and in the second result m can be given by a polynomial of n whose degree is $1/2$. See [14] for more details.

Let σ_{2n+3}^n be the n -skeleton of a $(2n+3)$ -simplex. In [18] it is shown that for any embedding of σ_{2n+3}^n into the $(2n+1)$ -sphere the sum of the linking numbers of the 2-component n -links contained in the embedding is an odd number. The case $n=1$ is just Theorem 3 (1). Thus this result is a higher dimensional generalization of Theorem 3 (1).

References

- [1] J. H. Conway and C. McA. Gordon: Knots and links in spatial graphs, *J. Graph Theory*, 7, 445-453, 1983.
- [2] E. Flapan: Intrinsic knotting and linking of complete graphs, preprint.
- [3] E. Flapan, R. Naimi and J. Pommersheim: Intrinsically triple linked complete graphs, *Topology Appl.*, 115, 239-246, 2001.
- [4] E. Flapan, J. Pommersheim, J. Foisy and R. Naimi: Intrinsically n -linked graphs, *J. Knot Theory Ramifications*, 10, 1143-1154, 2001.
- [5] A. Kawauchi: Almost identical imitations of $(3,1)$ -dimensional manifold pairs, *Osaka J. Math.*, 26, 743-758, 1989.
- [6] S. Kinoshita: On θ_n -curves in R^3 and their constituent knots, in *Topology and Computer Science* edited by S. Suzuki, Kinokuniya, 211-216, 1987.
- [7] M. S. Miyauchi: Topological Ramsey theorem for complete bipartite graphs, *J. Combin. Theory, Ser. B*, 62, 164-179, 1994.
- [8] T. Motohashi and K. Taniyama: Delta unknotting operation and vertex homotopy of graphs in R^3 , *Proceedings of Knots '96 Tokyo*, (S. Suzuki ed.), World Scientific Publ. Co., 1997.
- [9] S. Negami: Ramsey theorems for knots, links and spatial graphs, *Trans. Amer. Math. Soc.*, 324, 527-541, 1991.

- [10] S. Negami: Ramsey-type theorems for spatial graphs and good drawings, *J. Combin. Theory, Ser. B*, 72, 53-62, 1998.
- [11] R. Nikkuni and K. Onda: A characterization of knots in a spatial graph II, to appear in *J. Knot Theory Ramifications*.
- [12] Y. Ohyaama and K. Taniyama: Vassiliev invariants of knots in a spatial graph, *Pacific J. Math.*, 200, 191-205, 2001.
- [13] K. Onda: A characterization of knots in a spatial graph, *J. Knot Theory Ramifications*, 9, 1069-1084, 2000.
- [14] M. Shirai and K. Taniyama: A large complete graph in a space contains a link with large link invariant, in preparation.
- [15] T. Soma: Spatial-graph isotopy for trivalent graphs and minimally knotted embeddings, *Topology Appl.*, 73, 23-41, 1996.
- [16] K. Taniyama: Homology classification of spatial embeddings of a graph, *Topology Appl.*, 65, 205-228, 1995.
- [17] K. Taniyama: Knotted subgraphs in a spatial graph, *Kobe J. Math.*, 14, 207-212, 1997.
- [18] K. Taniyama: Higher dimensional links in a simplicial complex embedded in a sphere, *Pacific J. Math.*, 194, 465-467, 2000.
- [19] K. Taniyama and A. Yasuhara: Realization of knots and links in a spatial graph, *Topology Appl.*, 112, 87-109, 2001.
- [20] Ying Qing Wu: Minimally knotted embeddings of planar graphs, *Math. Z.*, 214, 653-658, 1993.
- [21] M. Yamamoto: Knots in spatial embeddings of the complete graph on four vertices, *Topology Appl.*, 36, 291-298, 1990.
- [22] A. Yasuhara: Delta-unknotted operation and adaptability of certain graphs, *Proceedings of Knots '96 Tokyo*, (S. Suzuki ed.), World Scientific Publ. Co., 1997.