

Properties of Zip4 accumulation during zinc deficiency and its usefulness to evaluate zinc status: A study of the effects of zinc deficiency during lactation

Ayako Hashimoto¹, Miki Nakagawa², Natsuki Tsujimura², Shiho Miyazaki¹, Kumiko Kizu³, Tomoko Goto², Yusuke Komatsu², Ayu Matsunaga⁴, Hitoshi Shirakawa², Hiroshi Narita⁴, Taiho Kambe^{1*} and Michio Komai²

AUTHOR CONTRIBUTIONS

A.H., T.K., and M.K. conceived of and designed the research study; A.H., M.N., N.T., S.M., K.K., T.G., Y.K., A.M., H.S., H.N., T.K., and M.K. performed experiments; A.H., M.N., N.T., S.M., T.G., T.K., and M.K. analyzed data; A.H., M.N., N.T., T.G., T.K., and M.K. interpreted the results of experiments; A.H., M.N., N.T., T.G., H.S., T.K., and M.K. prepared figures; A.H., T.K., and M.K. drafted the manuscript.

¹The Division of Integrated Life Science, Graduate School of Biostudies, Kyoto University, Kyoto 606-8502, Japan

²Department of Science of Food Function and Health, Graduate School of Agricultural Science, Tohoku University, Sendai 981-8555, Japan

³Department of Life and Living, Osaka Seikei College, Osaka 533-0007, Japan

⁴Department of Food Science, Kyoto Women's University, Kyoto 605-8501, Japan

Running head: Properties of Zip4 accumulation during zinc deficiency

23 *Address for reprint requests and other correspondence: Taiho Kambe, Division of Life Science, Graduate
24 School of Biostudies, Kyoto University, Kyoto 606-8502, Japan. Tel.: +81-75-753-6273; Fax:
25 +81-75-753-6274; E-mail: kambe1@kais.kyoto-u.ac.jp
26

ABSTRACT

Systemic and cellular zinc homeostasis is elaborately controlled by ZIP and ZnT zinc transporters. Therefore, detailed characterization of their expression properties is of importance. Of these transporter proteins, Zip4 functions as the primarily important transporter to control systemic zinc homeostasis because of its indispensable function of zinc absorption in the small intestine. In this study, we closely investigated Zip4 protein accumulation in the rat small intestine in response to zinc status using an anti-Zip4 monoclonal antibody that we generated, and contrasted this with the zinc-responsive activity of the membrane-bound alkaline phosphatase (ALP). We found that Zip4 accumulation is more rapid in response to zinc deficiency than previously thought. Accumulation increased in the jejunum as early as 1 day following a zinc-deficient diet. In the small intestine, Zip4 protein expression was higher in the jejunum than in the duodenum and was accompanied by reduction of ALP activity, suggesting that the jejunum can become zinc deficient more easily. Furthermore, by monitoring Zip4 accumulation levels and ALP activity in the duodenum and jejunum, we reasserted that zinc deficiency during lactation may transiently alter plasma glucose levels in the offspring in a sex-specific manner, without affecting homeostatic control of zinc metabolism. This confirms that zinc nutrition during lactation is extremely important for the health of the offspring. These results reveal that rapid Zip4 accumulation provides a significant conceptual advance in understanding of the molecular basis of systemic zinc homeostatic control, and that properties of Zip4 protein accumulation are useful to evaluate zinc status closely.

Keywords: zinc deficiency; Zip4 processing; small intestine; alkaline phosphatase; glucose homeostasis.

INTRODUCTION

Zinc plays a great variety of roles in diverse biological processes (35, 61), and therefore is crucial as an essential micronutrient for human health. Zinc deficiency causes a broad range of defects including diarrhea, alopecia, skin lesions, immune system and neuronal dysfunction, and taste disorders (13, 21, 35, 55), and is a clinical risk factor for a number of diseases in older people (20, 50). Because zinc deficiency also impairs optimal growth and development in infants, extremely low levels of zinc in breast milk, which is rarely caused by genetic diseases, results in severe health consequences in breast-fed infants (5, 26, 41, 47, 49). Moreover, maternal zinc deficiency during lactation, which results in low secretion of zinc (7), is shown to affect physiological responses in offspring in animal studies, such as effects on glucose metabolism (28), immune competence (56), and arterial blood pressure and renal functions (60). However, molecular investigations have not yet been fully performed.

Cellular and systemic zinc homeostasis is elaborately regulated by two zinc transporter families, ZIP and ZnT (6, 15, 36, 43). Among these, Zip4 is the most important transporter for the regulation of systemic zinc homeostasis because it plays an essential role in dietary zinc absorption in the small intestine (17). Additionally, it was identified as the gene responsible for acrodermatitis enteropathica (1, 33, 40, 63). Zip4 protein expression is dynamically regulated in response to zinc deficiency; Zip4 protein accumulates on the apical membrane by escaping from endocytosis and degradation, and undergoes processing by removal of the long extracellular amino-terminal region via proteolytic cleavage during prolonged zinc deficiency (9, 31, 37, 48, 64). Molecular and physiological properties of Zip4 protein expression and function have been characterized (2-4, 10, 11, 17, 45, 46, 62), but further investigation is needed for complete understanding. In the small intestine, membrane-bound alkaline phosphatase (ALP) activity is significantly decreased during

zinc-deficient conditions (32), and, thus, increased Zip4 expression and decreased membrane-bound ALP activity are considered to accurately reflect zinc status.

Zip4 accumulation in the small intestine during zinc deficiency has not yet been extensively investigated. In the present study, we examined it using an anti-Zip4 monoclonal antibody that we recently generated (22). The antibody clearly showed more rapid accumulation of Zip4 protein in the jejunum in response to a zinc-deficient diet than previously thought. Furthermore, in intimate monitoring of Zip4 expression levels detected by this antibody and the membrane-bound ALP activity, we examined the effects of zinc deficiency during lactation on glucose tolerance in the offspring, which has been reported to be affected (28). Evaluation of zinc status based on this perspective has not yet been conducted, and, therefore, would give novel insights into zinc physiology.

MATERIALS AND METHODS

Rat care, use, and dietary zinc manipulation. Four-week-old male Sprague-Dawley (SD) rats were purchased from Japan SLC (Hamamatsu, Japan), and acclimated for 3 d (fed a zinc-sufficient [Zn-Suf] diet). After 3 d of acclimation, rats were divided into three groups and fed one of the following diets for 4 d: zinc-deficient (Zn-Def), low-zinc (Zn-Low), and Zn-Suf diets, which contained 2.2, 4.1, and 33.7 mg zinc/kg, respectively, as described previously (38). In experiments of the effects of zinc-deficient dams on offspring, pregnant SD rats (16 d of pregnancy, d1=vaginal plug) were purchased from Japan SLC, and fed a MB-3 diet (Funabashi Farms, Funabashi, Japan) until parturition. On the day of parturition (considered lactation day 0), dams were divided into two groups and fed a Zn-Suf or Zn-Low diet until weaning day (3 weeks). After 3 weeks, nursing neonatal pups were weaned and fed a Zn-Suf diet and drinking water with free access. In all experiments, rats were housed in individual stainless steel cages under controlled

conditions (temperature, $23 \pm 2^{\circ}\text{C}$; humidity, $50 \pm 10\%$; 12-h light–dark cycle, 08:00 am to 08:00 pm), and were decapitated after being food deprived for 5 h. The blood and tissues, including small intestines, were collected at the time points specified in each experiment from the dams and pups, quickly frozen in liquid nitrogen, and used for experiments, such as membrane proteins preparation, as described below. Zinc concentrations were measured in plasma. Food intakes and body weights were measured each day. The experimental plan for the present study was approved by the Animal Research-Animal Care Committee of Tohoku University and Kyoto University. The entire experiment was carried out in accordance with the guidelines issued by this committee and Japanese governmental legislation (2005). The same committee supervised the care and use of the rats used in this study.

Generation of an anti-ZnT4 monoclonal antibody. Fused proteins consisting of the amino-terminal cytosolic region of human ZnT4 (112 amino acid residues from Met¹ to Ala¹¹²) and maltose binding protein were used as a ZnT4 antigen. The same peptide fused to glutathione S transferase protein was used as a solid-phase antigen for hybridoma screening as described previously (26, 34). The hybridoma that produces the anti-ZnT4 antibody was subcloned by the limiting dilution method. An ascites was generated by injection of 1×10^7 hybridoma cells into pristine-primed mice. Membrane proteins (20 μg) prepared from DT40 cells stably expressing carboxyl-terminally HA-tagged human ZnT4 (14) were used for examining the specificity of the antibody.

Preparation of membrane proteins from the small intestines and mammary glands. The small intestines, which were dissected and divided 0.5–2.5 cm distal from the pylorus for the duodenum and 5.5–8.5 cm for the jejunum in some experiments, were collected from male rats fed a Zn-Def, Zn-Low, or

Zn-Suf diet or lactating dams fed a Zn-Suf or Zn-Low diet and pups fostered by the dams on the indicated days. The mammary gland was simultaneously harvested from the dams. Tissue samples including the duodenum, jejunum, or mammary gland were homogenized using a homogenizer (Disperser T 10 basic, IKA, Tokyo, Japan) or multi-beads shocker (Yasui Kikai, Osaka, Japan), and membranes were recovered from the post-nuclear supernatant by centrifugation ($24,000 \times g$ for 30 min at 4°C). The membrane pellet was washed once with phosphate buffered saline (PBS) and suspended in alkaline phosphatase buffer (10 mM Tris-HCl, pH 7.5, 0.5 mM MgCl_2 , and 0.1% Triton X-100) and frozen at -80°C until assayed as described previously (32). The protein concentrations were determined using Protein Assay CBB Solution (Nacalai Tesque, Kyoto, Japan). Extraction of membrane proteins by Na_2CO_3 was performed as described previously (31). Briefly, membranes prepared from the small intestine of rats fed a Zn-Def diet for several days were resuspended in 0.05 mM Na_2CO_3 , and incubated on ice for 15 min before centrifugation for 30 min at $24,000 \times g$ at 4°C . The supernatant fraction (peripheral membrane proteins) and pellet fraction (integral membrane proteins) were recovered.

Immunoblot blot analyses. Membrane proteins (5 or 10 μg , if not indicated) were lysed in $6 \times$ SDS-sample buffer at 37°C for 30 min, separated by electrophoresis through 7.5% SDS polyacrylamide gels and transferred to polyvinylidene difluoride membranes (Pall Corporation, East Hills, NY). The blot was blocked with blocking solution (1% skim milk and 0.1% Tween 20 in PBS) or SuperBlock blocking buffer (Thermo Fisher Scientific, Rockford, IL) supplemented with 0.1% Tween 20 for detection of rat Zip4, and then incubated with an anti-Zip4 monoclonal antibody (1:1000 dilution) (22), anti-ZIP4 polyclonal antiserum (1:600) (64), anti-ZnT2 (1:2000) (26), anti-ZnT4 (1:1000), anti-GPR39 (Abcam, Cambridge, UK, 1:1500), and anti-GRP78 (ABR, Golden, CO, 1:1000). Horseradish peroxidase-conjugated anti-mouse or

anti-rabbit secondary antibodies (GE Healthcare, Waukesha, WI) were used at a 1:3000 dilution for detection. Immunoreactive bands were visualized using Immobilon Western Chemiluminescent HRP Substrates (Millipore, Billerica, MA) or Chemi-Lumi One L (Nacalai Tesque). The fluoroimage was obtained using a LAS1000 plus image analyzer (Fujifilm, Tokyo, Japan) and LAS500 (GE Healthcare).

Membrane-bound alkaline phosphatase assay. Membrane proteins (1–3 μ g) were assayed for alkaline phosphatase activity with 2 mg/mL *p*-nitrophenyl phosphate in 1 M diethanolamine buffer, pH 9.8, containing 0.5 mM MgCl₂ as described previously (16, 32). *p*-Nitrophenol release was measured by the absorbance at 405 nm. Calf intestine alkaline phosphatase (Promega, Madison, WI) was used as the standard (14).

Measurement of plasma glucose and insulin levels. Plasma glucose levels were measured by enzymatic colorimetric methods using the Glucose C-II test kit (Wako Pure Chemical Co., Osaka, Japan) after 5 h of starvation at 4, 7, 9, 11, 13, and 15 weeks. Plasma insulin levels were measured using a rat insulin ELISA kit (Morinaga Institute of Biological Science, Inc. Yokohama, Japan).

Statistical analyses. Values for animal tissues are expressed as means $\pm$ standard error of the mean. Statistical comparisons between each group were performed using one-way analysis of variance followed by Tukey's multiple comparisons *post hoc* test. $P < 0.05$ was considered significant. The experimental data of biochemical measurements including ALP activity are depicted as mean $\pm$ standard deviation, where statistical significance was determined by the Student's *t*-test and accepted as significant at $P < 0.05$.

RESULTS

Intestinal Zip4 expression is rapidly increased in the rat small intestine. Expression of Zip4 in the small intestine is highly induced in mice fed a zinc-deficient diet (9, 10, 32, 45, 64), but that has not been well investigated in other species, including rats. The anti-Zip4 monoclonal antibody, which we have recently reported (22), clearly detected Zip4 as two bands (~75 kDa and ~38 kDa) in the membrane fractions prepared from the small intestine of rats fed the Zn-Def diet (Fig. 1A, left panel). The monoclonal antibody was generated against the extracellular amino terminal portion (ectodomain) of Zip4 (22), and, thus, the two bands correspond to the full-length (~75 kDa) and the cleaved ectodomain (~38 kDa). This was confirmed by similar results obtained using ZIP4 polyclonal antiserum (Fig. 1A, right panel), which was directed against the large intracellular loop between Zip4 transmembrane domains (TMDs) III and IV. Thus, the antibody detected both the full-length and the processed Zip4 corresponding to Zip4 with eight TMDs without the ectodomain (31). In our previous report (31), we calculated the size of the processed Zip4 as ~37 kDa, but the band was detected as ~40 kDa in the present study because of the use of different protein size markers. The cleaved ectodomain (~38 kDa) of rat Zip4 was prepared in the membrane fractions as a peripheral membrane protein loosely associated with the membrane fractions, as in the case of mouse Zip4 (31) because it was extracted in the supernatant with Na_2CO_3 , while the full-length Zip4 and Znt4 (both integral membrane proteins) were not (Fig. 1B).

The clear detection of Zip4 by our monoclonal antibody enabled us to investigate more closely Zip4 expression regulation in adult male rats fed the Zn-Def diet. First, we examined the response of Zip4 expression to zinc-deficient diets. We divided rats in three groups fed Zn-Def, Zn-Low, or Zn-Suf diets and

monitored Zip4 expression in the small intestine by immunoblot analysis (Fig. 2A). Both the full-length form and the cleaved ectodomain of Zip4 were faintly detected after 1 d, and the cleaved ectodomain was clearly detected after 2 d. The membrane-bound ALP activity, which reflects zinc-deficient status (16, 32), decreased dose-dependently in response to the zinc contents of the diets (e.g., 3.8 mU/ μ g protein in the Zn-Suf group, 2.0 mU/ μ g protein in the Zn-Low group, and 1.2 mU/ μ g protein in the Zn-Def group after 1 d) (Fig. 2B). Moreover, rat plasma zinc concentrations were reduced by about 50% after 1 d when fed the Zn-Low or Zn-Def diet compared with the Zn-Suf diet (Fig. 2C). Thus, Zip4 protein accumulation and processing events likely occurred following the reduction of plasma zinc concentrations and membrane-bound ALP activity. However, the processing event may be sophisticatedly controlled by the degree of severity of zinc deficiency because the ectodomain of Zip4 was more enhanced in the Zn-Def group than in the Zn-Low group compared with differences in plasma zinc concentrations and ALP activity on days 1 and 2 (Fig. 2A, B and C).

We then explored region-specific differences in Zip4 expression. We dissected and divided the small intestine into two pieces, the duodenum (0.5–2.5 cm distal from the pylorus) and jejunum (5.5–8.5 cm), and examined Zip4 expression (Fig. 2D). Zip4 expression increased more intensely and rapidly in the jejunum than in the duodenum. This indicates zinc sensitivity was different between the duodenum and jejunum. These close investigations clearly indicate Zip4 expression is sensitively responsive to zinc deficiency in the rat small intestine.

The effects of zinc deficiency on lactating dams. To investigate the effects of zinc deficiency during lactation on pups, dams were fed Zn-Low or Zn-Suf diets from birth until weaning (for 3 weeks). As described in previous reports (18, 53, 58), the amount of food intake decreased in dams fed the Zn-Low diet

(Fig. 3A). Anorexia from zinc deficiency was observed by a typical cyclical 3- to 4-d pattern of decreased food intake in dams fed the Zn-Low diet. This was accompanied by decreases in body weights (Fig. 3B) and reduced plasma zinc concentrations (about 80% lower) (Fig. 3C) compared with those fed the Zn-Suf diet. Membrane proteins were prepared from the duodenum and jejunum of dams fed both Zn-Suf and Zn-Low diets at 3 weeks and analyzed by immunoblot. Zip4 of the full-length and the ectodomain, the latter of which was much higher, was detected in the duodenal and jejunal membranes prepared from the Zn-Low dams (Fig. 4A) but not in those of the Zn-Suf dams. Zip4 expression was somewhat higher in the jejunum than duodenum, although less significant compared with that in male rats fed a Zn-Def diet for a shorter period (Fig. 2D). Consistent with the result, the reduction rate of the membrane-bound ALP activity was more significant in the jejunum than in the duodenum of dams fed a Zn-Low diet, both of which were significantly reduced (Fig. 4B). Because Znt4 and the zinc receptor GPR39 are known to be highly expressed in the small intestine (19, 44, 51, 52), expression levels of both proteins were also examined. Compared with that of Zip4, the expression was not significantly affected by the Zn-Low diet (Fig. 4A). Moreover, because we have antibodies against both Znt2 and Znt4, both of which are known to be responsible for supplying zinc into the breast milk in humans (5, 26, 41, 47, 49) and in mice (24, 42), we investigated the effects of zinc deficiency on the expression of both proteins in mammary glands. The expression was not significantly changed between dams fed Zn-Low and Zn-Suf diets compared with Zip4 expression in the jejunum and duodenum (Fig. 4C). However, the membrane-bound ALP activity in the mammary glands was significantly reduced in dams fed a Zn-Low diet (Fig. 4D).

The effects of zinc deficiency during the lactating period on pre-weanling pups. We next examined the effects of Zn-Low diet feeding of dams on their pups. The body weights of pups fostered by dams fed a

Zn-Low diet were reduced at 1 week after birth and more significantly reduced at weaning (3 weeks after delivery) compared with those fed a Zn-Suf diet (Fig. 5A). At weaning (3 weeks), plasma zinc concentrations were about 60% lower in pups fostered by dams fed a Zn-Low diet than those fed a Zn-Suf diet (Fig. 5B), indicating that the pups were severely zinc deficient. This was clearly confirmed by the fact that Zip4 expression was increased in both the duodenum and jejunum of pups fostered with dams fed a Zn-Low diet (Fig. 5C). Moreover, membrane-bound ALP activity was severely decreased in both tissues (Fig. 5D). Under our experimental conditions, the effects of Zn-Low diet feeding of dams were obvious on Zip4 expression and ALP activity in pups. Znt4 and GPR39 expression levels were not significantly changed between the two groups (Fig. 5C), as with the dams (Fig. 4A).

The effects of zinc deficiency during lactation on plasma glucose levels in offspring. The weaned pups fostered by dams fed Zn-Suf or Zn-Low diets were fed a Zn-Suf diet. Because the amount of food intake and the body weight gain became almost the same within 1 week, we isolated the duodenum and jejunum from pups in both groups at 4 weeks and examined the same items as those measured at 3 weeks. Plasma zinc concentrations were not significantly different between both groups (Fig. 6A), and Zip4 expression was not detected at the same exposure time as used in Fig. 5C in both the duodenum and jejunum of pups fostered by dams fed a Zn-Low diet until weaning (Fig. 6B). Consistent with this, membrane-bound ALP activity in both tissues was almost equivalent between the two groups (Fig. 6C). These results confirmed that pups fostered by dams fed a Zn-Low diet until weaning were not apparently zinc deficient and maintain normal homeostatic control of zinc metabolism. Znt4 and GPR39 expression levels were not significantly changed between the two groups (Fig. 6B), consistent with observations in the pups at weaning (3 weeks).

Last, we investigated the effects of zinc deficiency during lactation on plasma glucose

homeostasis in the offspring because maternal zinc deficiency during this period is shown to induce long-term changes related to abnormal glucose tolerance in the offspring (28). Plasma glucose levels were higher in male offspring fostered by dams fed a Zn-Low diet at 11 and 13 weeks (Fig. 7A); however, plasma insulin levels were not significantly changed (Fig. 7B). We did not observe similar changes in plasma glucose levels in the female offspring fostered by dams fed a Zn-Low diet (Fig. 7C). Taken together, these results suggest that zinc deficiency during lactation may transiently alter glucose homeostasis in the offspring in a sex-specific manner, even after the status and homeostatic control of zinc become normal.

DISCUSSION

Given that many zinc nutrition studies are performed using rats, clarification of zinc absorption processes in the rat small intestine, including Zip4 expression regulation, is important. However, there are few examinations of rat Zip4 expression. Moreover, the fundamental issue relating to rat Zip4 is that several discrepancies have been identified between rat and mouse Zip4 detection by immunoblot analysis, despite the high sequence identities (over 89%; similarity, 93%). Specifically, Zip4 protein accumulation during zinc deficiency was drastically increased or not, and the immunoreactive bands corresponding to Zip4 were detected at either 70–75 kDa or 35–40 kDa (25, 27, 31, 36, 45, 64). In this study, we revealed that Zip4 was detected in several bands corresponding to the full-length, the processed or the cleaved ectodomain of Zip4. Furthermore, Zip4 accumulation was dramatic in the cleaved ectodomain form (implying that the processed Zip4 was also increased) but not so significant in the full-length form during zinc deficiency (Fig. 2A and D). Thus, previous discrepancies in Zip4 detection likely resulted from differences in the detected immunoreactive bands, which are altered by zinc deficiency duration. The inducibility of Zip4 accumulation is different in the regions of the small intestine (e.g., Zip4 expression is higher in the jejunum than

duodenum), which may be an alternative reason why the discrepancies occurred in different immunoblots. Our results are important in clarifying these points and, moreover, revealed that Zip4 expression regulation in rats and mice is consistent. The rapid up-regulation of Zip4 expression in the small intestine during zinc-deficient conditions corroborates the important role of Zip4 in the systemic control of zinc homeostasis.

Another interesting aspect in this study on Zip4 expression regulation is that the processing of Zip4 clearly occurred by 2 d at the latest after feeding with the Zn-Def and Zn-Low diets. The cleaved ectodomain was faintly detected after 1 d in the jejunum (Fig. 2D, right panel). This is in contrast to the previous thought that Zip4 processing needs a prolonged amount of time during zinc-deficient conditions, which was derived from an *in vitro* study using Hepa cells or transfected cells (31). The appearance timing of the cleaved ectodomain of Zip4 preceded the first significant suppression of food intake, which began at 3 d after feeding the same Zn-Low diet in our experimental conditions (53) (Fig. 3A). An orexigenic signal by dietary zinc, which could be derived from the intestine to the brain via vagus nerve signal transduction, occurs during oral but not intraperitoneal administration (53). Thus, it is an intriguing hypothesis that the processing of Zip4 (or the processed or the cleaved ectodomain of Zip4) in enterocytes may be involved in the suppression of food intake, as well as controlling zinc transport activity as discussed below.

The association of the cleaved Zip4 ectodomain with the cellular membrane after processing, which was proposed in an *in vitro* study (31), was confirmed in the rat small intestine (see Fig. 1B). The processing event in Zip4 or the cleaved ectodomain of Zip4 itself can be considered physiologically relevant because acrodermatitis enteropathica-causing mutations inhibit the processing of Zip4 (31). Considering these and the abundance of histidine and cysteine residues in the ectodomain (15, 35), the ectodomain may capture dietary zinc and control its transfer to the pore formed by eight TMDs of Zip4 (3). Alternatively, the ectodomain itself may control pore opening or closing through its releasing (or inhibiting its release), as

previously speculated (59). Processing may be an important, conserved regulatory mechanism in some ZIP transporters because similar ectodomain cleavage has been found in ZIP6 and ZIP10 (12, 23), both of which possess the long extracellular amino-terminal portion abundant in histidine and cysteine residues like ZIP4 (15, 35). Interestingly, the ectodomain of the transporter involved in copper absorption is proteolytically cleaved, which is thought to control copper uptake (54). Thus, the processing event in metal transporters may be one of the important regulatory mechanisms to control metal uptake and absorption rates from the extracellular milieu.

In vertebrates, several ALP isozymes (tissue non-specific, intestine and germ cell specific, and placenta specific in humans) are expressed. Each of these is probably zinc dependent, although zinc dependence of the germ cell-specific ALP has not been molecularly examined (29, 30). In this study, we used membrane-bound ALP activity as a marker reflecting zinc status in examined tissues as in our previous report (36). Membrane-bound ALP activity in the small intestine was dependent on dietary zinc contents, which was inversely associated with Zip4 expression levels (see Fig. 2A and B). Along with more enhanced Zip4 expression (both the full length and the cleaved ectodomain) in the jejunum than the duodenum (Figs. 2D, 4A and 5C), the reduction rate of the membrane-bound ALP activity, which is considered by comparing that in Zn-Low with that in Zn-Suf, was more significant in the jejunum than the duodenum (Figs. 4B and 5D). These results indicate that the jejunum likely becomes more zinc deficient than the duodenum during zinc deficiency, and may suggest that a distal part of the intestine from the pylorus may become more easily zinc deficient. This hypothesis should probably be extensively explored because one of representative features of zinc deficiency is diarrhea. Moreover, membrane-bound ALP activity in mammary glands was significantly decreased in the dams fed a Zn-Low diet compared with those fed a Zn-Suf diet. Znt2 and Znt4 protein expression levels were not significantly changed between both groups (see Fig. 4C and 4D),

indicating both Znt2 and Znt4 protein expression are less sensitive to zinc deficiency in mammary glands. Thus, membrane-bound ALP activity would be useful to evaluate apparent zinc status and to investigate zinc-responsive protein expression including ZIP and ZnT transporters in tissues.

Lastly, the quite important aspect of this study is that we reasserted the effects of zinc deficiency during lactation on plasma glucose homeostasis in male offspring, even if zinc status and homeostatic control were normally regulated. Similar results, including the sex-specific effects of blood glucose levels, were previously reported by Jou et al. (28); however, some discrepancies are present between our results and previous reports. This may be because dams were fed a zinc-deficient diet (7 mg/kg) and control diet (25 mg/kg) for 6 weeks from 3 weeks preconception to 21 d post-parturition in the previous study (28), while dams were fed Zn-Low (4.1 mg/kg) and Zn-Suf (33.7 mg/kg) diets in this study. Importantly, the molecular mechanism of how plasma glucose levels are affected by zinc deficiency during lactation has not yet been clarified, although epigenetic changes may occur on the promoter of molecules involved in glucose homeostasis during zinc deficiency, like in the case of the epigenetic alteration in the MT2 promoter region during zinc deficiency *in utero* (39). These points should be more intensely investigated in a future study. About 80% of pregnant and lactating women are estimated to be at risk for marginal zinc deficiency (8, 57). Thus, zinc levels during this period should receive much attention from viewpoints of optimal growth and development, as well as future good health in offspring.

PERSPECTIVES AND SIGNIFICANCE

Zip4 accumulation in the small intestine rapidly occurred during zinc deficiency. The processing of Zip4 also rapidly occurred during zinc deficiency. These properties of Zip4 may be useful as a biomarker reflecting zinc deficient levels in the body, as well as for evaluating bioavailable zinc levels. In fact, the

potential efficacy of detecting Zip4 expression as a marker was described in the study of zinc deficiency during lactation, contrasting this with the zinc-responsive activity of ALP. Moreover, the rapid processing of Zip4 during zinc deficiency suggests its physiological importance in the regulation to control (or activate) its zinc transport function. The result would lead to new directions in the research field of zinc transporters because the processing is found in other ZIP transporters. Zinc deficiency puts human health at risk and is still a worldwide problem (65, 66). Thus, comprehensive understanding of the expression properties of ZIP4 is of critical importance, given its primary importance to control systemic zinc homeostasis. The present study may provide useful information to establish new strategies to prevent zinc deficiency.

ACKNOWLEDGMENTS

We thank Dr. Glen K. Andrews (University of Kansas Medical Center) for providing anti-ZIP4 polyclonal antiserum. We are also grateful to Kana Morimoto for her technical assistance.

GRANTS

This work was supported by Grants-in-Aid for Challenging Exploratory Research and Scientific Research (B) (KAKENHI, Grant Nos. 26660086 and 15H04501 to T.K.), and Scientific Research (A) (KAKENHI, Grant No. 23248020 to M.K.), from the Japan Society for the Promotion of Science and by the Fuji Foundation for Protein Research, the Central Miso Research Institute, the Skylark Food Science Institute, the Iijima Memorial Foundation for the Promotion of Food Science and Technology, the Japan Food Chemical Research Foundation, the Asahi Group Foundation, the Mitsui Sumitomo Insurance Welfare

Foundation, and the Yamazaki Spice Promotion Foundation (to T.K.), and the Food Science Institute Foundation (to M.K.). A.H. is a Research Fellow (DC2) of the Japan Society for the Promotion of Science.

DISCLOSURES

No conflicts of interest, financial or otherwise, are declared by the authors.

REFERENCES

1. **Andrews GK.** Regulation and function of Zip4, the acrodermatitis enteropathica gene. *Biochem Soc Trans* 36: 1242-1246, 2008.
2. **Antala S, and Dempski RE.** The human ZIP4 transporter has two distinct binding affinities and mediates transport of multiple transition metals. *Biochemistry* 51: 963-973, 2012.
3. **Antala S, Ovchinnikov S, Kamisetty H, Baker D, and Dempski RE.** Computation and Functional Studies Provide a Model for the Structure of the Zinc Transporter hZIP4. *J Biol Chem* 290: 17796-17805, 2015.
4. **Bafaro EM, Antala S, Nguyen TV, Dzul SP, Doyon B, Stemmler TL, and Dempski RE.** The large intracellular loop of hZIP4 is an intrinsically disordered zinc binding domain. *Metallomics* 7: 1319-1330, 2015.
5. **Chowanadisai W, Lonnerdal B, and Kelleher SL.** Identification of a mutation in SLC30A2 (ZnT-2) in women with low milk zinc concentration that results in transient neonatal zinc deficiency. *J Biol Chem* 281: 39699-39707, 2006.

- 378 6. **Cousins RJ, Liuzzi JP, and Lichten LA.** Mammalian zinc transport, trafficking, and signals. *J*
379 *Biol Chem* 281: 24085-24089, 2006.
- 380 7. **Dempsey C, McCormick NH, Croxford TP, Seo YA, Grider A, and Kelleher SL.** Marginal
381 maternal zinc deficiency in lactating mice reduces secretory capacity and alters milk composition. *J Nutr*
382 142: 655-660, 2012.
- 383 8. **Donangelo CM, and King JC.** Maternal zinc intakes and homeostatic adjustments during
384 pregnancy and lactation. *Nutrients* 4: 782-798, 2012.
- 385 9. **Dufner-Beattie J, Kuo YM, Gitschier J, and Andrews GK.** The adaptive response to dietary
386 zinc in mice involves the differential cellular localization and zinc regulation of the zinc transporters ZIP4
387 and ZIP5. *J Biol Chem* 279: 49082-49090, 2004.
- 388 10. **Dufner-Beattie J, Wang F, Kuo YM, Gitschier J, Eide D, and Andrews GK.** The
389 acrodermatitis enteropathica gene ZIP4 encodes a tissue-specific, zinc-regulated zinc transporter in mice. *J*
390 *Biol Chem* 278: 33474-33481, 2003.
- 391 11. **Dufner-Beattie J, Weaver BP, Geiser J, Bilgen M, Larson M, Xu W, and Andrews GK.** The
392 mouse acrodermatitis enteropathica gene Slc39a4 (Zip4) is essential for early development and
393 heterozygosity causes hypersensitivity to zinc deficiency. *Hum Mol Genet* 16: 1391-1399, 2007.
- 394 12. **Ehsani S, Salehzadeh A, Huo H, Reginold W, Pocanschi CL, Ren H, Wang H, So K, Sato C,**
395 **Mehrabian M, Strome R, Trimble WS, Hazrati LN, Rogaeva E, Westaway D, Carlson GA, and**
396 **Schmitt-Ulms G.** LIV-1 ZIP Ectodomain Shedding in Prion-Infected Mice Resembles Cellular Response to
397 Transition Metal Starvation. *J Mol Biol* 422: 556-574, 2012.
- 398 13. **Fraker PJ, and King LE.** Reprogramming of the immune system during zinc deficiency. *Annu*
399 *Rev Nutr* 24: 277-298, 2004.

- 400 14. **Fujimoto S, Isumura N, Tsuji T, Anan Y, Tsuji N, Ogra Y, Kimura T, Miyamae Y,**
401 **Masuda S, Nagao M, and Kambe T.** Cooperative Functions of ZnT1, Metallothionein and ZnT4 in the
402 Cytoplasm Are Required for Full Activation of TNAP in the Early Secretory Pathway. *PLoS One* 8: e77445,
403 2013.
- 404 15. **Fukada T, and Kambe T.** Molecular and genetic features of zinc transporters in physiology and
405 pathogenesis. *Metallomics* 3: 662-674, 2011.
- 406 16. **Fukunaka A, Kurokawa Y, Teranishi F, Sekler I, Oda K, Ackland ML, Faundez V,**
407 **Hiromura M, Masuda S, Nagao M, Enomoto S, and Kambe T.** Tissue Nonspecific Alkaline Phosphatase
408 Is Activated via a Two-step Mechanism by Zinc Transport Complexes in the Early Secretory Pathway. *J*
409 *Biol Chem* 286: 16363-16373, 2011.
- 410 17. **Geiser J, Venken KJ, De Lisle RC, and Andrews GK.** A Mouse Model of Acrodermatitis
411 Enteropathica: Loss of Intestine Zinc Transporter ZIP4 (Slc39a4) Disrupts the Stem Cell Niche and Intestine
412 Integrity. *PLoS Genet* 8: e1002766, 2012.
- 413 18. **Goto T, Shirakawa H, Furukawa Y, and Komai M.** Decreased expression of carbonic
414 anhydrase isozyme II, rather than of isozyme VI, in submandibular glands in long-term zinc-deficient rats.
415 *Br J Nutr* 99: 248-253, 2008.
- 416 19. **Guthrie GJ, Aydemir TB, Troche C, Martin AB, Chang SM, and Cousins RJ.** Influence of
417 ZIP14 (slc39A14) on intestinal zinc processing and barrier function. *Am J Physiol Gastrointest Liver*
418 *Physiol* 308: G171-178, 2015.
- 419 20. **Haase H, Mocchegiani E, and Rink L.** Correlation between zinc status and immune function in
420 the elderly. *Biogerontology* 7: 421-428, 2006.
- 421 21. **Hambidge M.** Human zinc deficiency. *J Nutr* 130: 1344S-1349S, 2000.

- 422 22. **Hashimoto A, Ohkura K, Takahashi M, Kizu K, Narita H, Enomoto S, Miyamae Y, Masuda**
423 **S, Nagao M, Irie K, Ohigashi H, Andrews GK, and Kambe T.** Soybean extracts increase cell surface
424 ZIP4 abundance and cellular zinc levels: a potential novel strategy to enhance zinc absorption by ZIP4
425 targeting. *Biochem J* 472: 183-193, 2015.
- 426 23. **Hogstrand C, Kille P, Ackland ML, Hiscox S, and Taylor KM.** A mechanism for
427 epithelial-mesenchymal transition and anoikis resistance in breast cancer triggered by zinc channel ZIP6 and
428 STAT3 (signal transducer and activator of transcription 3). *Biochem J* 455: 229-237, 2013.
- 429 24. **Huang L, and Gitschier J.** A novel gene involved in zinc transport is deficient in the lethal milk
430 mouse. *Nat Genet* 17: 292-297., 1997.
- 431 25. **Huang ZL, Dufner-Beattie J, and Andrews GK.** Expression and regulation of SLC39A family
432 zinc transporters in the developing mouse intestine. *Dev Biol* 295: 571-579, 2006.
- 433 26. **Itsumura N, Inamo Y, Okazaki F, Teranishi F, Narita H, Kambe T, and Kodama H.**
434 Compound Heterozygous Mutations in SLC30A2/ZnT2 Results in Low Milk Zinc Concentrations: A Novel
435 Mechanism for Zinc Deficiency in a Breast-Fed Infant. *PLoS One* 8: e64045, 2013.
- 436 27. **Jou MY, Hall AG, Philipps AF, Kelleher SL, and Lonnerdal B.** Tissue-specific alterations in
437 zinc transporter expression in intestine and liver reflect a threshold for homeostatic compensation during
438 dietary zinc deficiency in weanling rats. *J Nutr* 139: 835-841, 2009.
- 439 28. **Jou MY, Philipps AF, and Lonnerdal B.** Maternal zinc deficiency in rats affects growth and
440 glucose metabolism in the offspring by inducing insulin resistance postnatally. *J Nutr* 140: 1621-1627, 2010.
- 441 29. **Kambe T.** Methods to Evaluate Zinc Transport into and out of the Secretory and
442 Endosomal-Lysosomal Compartments in DT40 Cells. *Methods Enzymol* 534: 77-92, 2014.

- 443 30. **Kambe T.** An overview of a wide range of functions of ZnT and Zip zinc transporters in the
444 secretory pathway. *Biosci Biotechnol Biochem* 75: 1036-1043, 2011.
- 445 31. **Kambe T, and Andrews GK.** Novel proteolytic processing of the ectodomain of the zinc
446 transporter ZIP4 (SLC39A4) during zinc deficiency is inhibited by acrodermatitis enteropathica mutations.
447 *Mol Cell Biol* 29: 129-139, 2009.
- 448 32. **Kambe T, Geiser J, Lahner B, Salt DE, and Andrews GK.** Slc39a1 to 3 (subfamily II) Zip
449 genes in mice have unique cell-specific functions during adaptation to zinc deficiency. *Am J Physiol Regul*
450 *Integr Comp Physiol* 294: R1474-1481, 2008.
- 451 33. **Kambe T, Hashimoto A, and Fujimoto S.** Current understanding of ZIP and ZnT zinc
452 transporters in human health and diseases. *Cell Mol Life Sci* 71: 3281-3295, 2014.
- 453 34. **Kambe T, Narita H, Yamaguchi-Iwai Y, Hirose J, Amano T, Sugiura N, Sasaki R, Mori K,**
454 **Iwanaga T, and Nagao M.** Cloning and characterization of a novel mammalian zinc transporter, zinc
455 transporter 5, abundantly expressed in pancreatic beta cells. *J Biol Chem* 277: 19049-19055., 2002.
- 456 35. **Kambe T, Tsuji T, Hashimoto A, and Itsumura N.** The Physiological, Biochemical, and
457 Molecular Roles of Zinc Transporters in Zinc Homeostasis and Metabolism. *Physiol Rev* 95: 749-784, 2015.
- 458 36. **Kambe T, Weaver BP, and Andrews GK.** The genetics of essential metal homeostasis during
459 development. *Genesis* 46: 214-228, 2008.
- 460 37. **Kim BE, Wang F, Dufner-Beattie J, Andrews GK, Eide DJ, and Petris MJ.** Zn²⁺-stimulated
461 endocytosis of the mZIP4 zinc transporter regulates its location at the plasma membrane. *J Biol Chem* 279:
462 4523-4530, 2004.

- 463 38. **Komai M, Goto T, Suzuki H, Takeda T, and Furukawa Y.** Zinc deficiency and taste
464 dysfunction; contribution of carbonic anhydrase, a zinc-metalloenzyme, to normal taste sensation. *Biofactors*
465 12: 65-70, 2000.
- 466 39. **Kurita H, Ohsako S, Hashimoto S, Yoshinaga J, and Tohyama C.** Prenatal zinc
467 deficiency-dependent epigenetic alterations of mouse metallothionein-2 gene. *J Nutr Biochem* 24: 256-266,
468 2013.
- 469 40. **Kury S, Dreno B, Bezieau S, Giraudet S, Kharfi M, Kamoun R, and Moisan JP.**
470 Identification of SLC39A4, a gene involved in acrodermatitis enteropathica. *Nat Genet* 31: 239-240., 2002.
- 471 41. **Lasry I, Seo YA, Ityel H, Shalva N, Pode-Shakked B, Glaser F, Berman B, Berezovsky I,**
472 **Goncearenco A, Klar A, Levy J, Anikster Y, Kelleher SL, and Assaraf YG.** A Dominant Negative
473 Heterozygous G87R Mutation in the Zinc Transporter, ZnT-2 (SLC30A2), Results in Transient Neonatal
474 Zinc Deficiency. *J Biol Chem* 287: 29348-29361, 2012.
- 475 42. **Lee S, Hennigar SR, Alam S, Nishida K, and Kelleher SL.** Essential Role for Zinc Transporter
476 2 (ZnT2)-mediated Zinc Transport in Mammary Gland Development and Function during Lactation. *J Biol*
477 *Chem* 290: 13064-13078, 2015.
- 478 43. **Lichten LA, and Cousins RJ.** Mammalian zinc transporters: nutritional and physiologic
479 regulation. *Annu Rev Nutr* 29: 153-176, 2009.
- 480 44. **Liuzzi JP, Blanchard RK, and Cousins RJ.** Differential regulation of zinc transporter 1, 2, and
481 4 mRNA expression by dietary zinc in rats. *J Nutr* 131: 46-52., 2001.
- 482 45. **Liuzzi JP, Bobo JA, Lichten LA, Samuelson DA, and Cousins RJ.** Responsive transporter
483 genes within the murine intestinal-pancreatic axis form a basis of zinc homeostasis. *Proc Natl Acad Sci U S*
484 *A* 101: 14355-14360, 2004.

- 485 46. **Liuzzi JP, Guo L, Chang SM, and Cousins RJ.** Kruppel-like factor 4 regulates adaptive
486 expression of the zinc transporter Zip4 in mouse small intestine. *Am J Physiol Gastrointest Liver Physiol*
487 296: G517-523, 2009.
- 488 47. **Lova Navarro M, Vera Casano A, Benito Lopez C, Fernandez Ballesteros MD, Godoy Diaz**
489 **DJ, Crespo Erchiga A, and Romero Brufau S.** Transient neonatal zinc deficiency due to a new autosomal
490 dominant mutation in gene SLC30A2 (ZnT-2). *Pediatr Dermatol* 31: 251-252, 2014.
- 491 48. **Mao X, Kim BE, Wang F, Eide DJ, and Petris MJ.** A histidine-rich cluster mediates the
492 ubiquitination and degradation of the human zinc transporter, hZIP4, and protects against zinc cytotoxicity. *J*
493 *Biol Chem* 282: 6992-7000, 2007.
- 494 49. **Miletta MC, Bieri A, Kernland K, Schoni MH, Petkovic V, Fluck CE, Eble A, and Mullis**
495 **PE.** Transient Neonatal Zinc Deficiency Caused by a Heterozygous G87R Mutation in the Zinc Transporter
496 ZnT-2 (SLC30A2) Gene in the Mother Highlighting the Importance of Zn (2+) for Normal Growth and
497 Development. *Int J Endocrinol* 2013: 259189, 2013.
- 498 50. **Mocchegiani E, Romeo J, Malavolta M, Costarelli L, Giacconi R, Diaz LE, and Marcos A.**
499 Zinc: dietary intake and impact of supplementation on immune function in elderly. *Age (Dordr)* 35: 839-860,
500 2013.
- 501 51. **Moechars D, Depoortere I, Moreaux B, de Smet B, Goris I, Hoskens L, Daneels G, Kass S,**
502 **Ver Donck L, Peeters T, and Coulie B.** Altered gastrointestinal and metabolic function in the
503 GPR39-obestatin receptor-knockout mouse. *Gastroenterology* 131: 1131-1141, 2006.
- 504 52. **Murgia C, Vespignani I, Cerase J, Nobili F, and Perozzi G.** Cloning, expression, and vesicular
505 localization of zinc transporter Dri 27/ZnT4 in intestinal tissue and cells. *Am J Physiol* 277: G1231-1239.,
506 1999.

- 507 53. **Ohinata K, Takemoto M, Kawanago M, Fushimi S, Shirakawa H, Goto T, Asakawa A, and**
508 **Komai M.** Orally administered zinc increases food intake via vagal stimulation in rats. *J Nutr* 139: 611-616,
509 2009.
- 510 54. **Ohrvik H, Nose Y, Wood LK, Kim BE, Gleber SC, Ralle M, and Thiele DJ.** Ctr2 regulates
511 biogenesis of a cleaved form of mammalian Ctr1 metal transporter lacking the copper- and cisplatin-binding
512 ecto-domain. *Proc Natl Acad Sci U S A* 110: E4279-4288, 2013.
- 513 55. **Prasad AS.** Discovery of human zinc deficiency and studies in an experimental human model.
514 *Am J Clin Nutr* 53: 403-412, 1991.
- 515 56. **Raqib R, Hossain MB, Kelleher SL, Stephensen CB, and Lonnerdal B.** Zinc supplementation
516 of pregnant rats with adequate zinc nutriture suppresses immune functions in their offspring. *J Nutr* 137:
517 1037-1042, 2007.
- 518 57. **Scheplyagina LA.** Impact of the mother's zinc deficiency on the woman's and newborn's health
519 status. *J Trace Elem Med Biol* 19: 29-35, 2005.
- 520 58. **Shay NF, and Mangian HF.** Neurobiology of zinc-influenced eating behavior. *J Nutr* 130:
521 1493S-1499S, 2000.
- 522 59. **Taylor KM, Morgan HE, Smart K, Zahari NM, Pumford S, Ellis IO, Robertson JF, and**
523 **Nicholson RI.** The emerging role of the LIV-1 subfamily of zinc transporters in breast cancer. *Mol Med* 13:
524 396-406, 2007.
- 525 60. **Tomat AL, Costa Mde L, and Arranz CT.** Zinc restriction during different periods of life:
526 influence in renal and cardiovascular diseases. *Nutrition* 27: 392-398, 2011.
- 527 61. **Vallee BL, and Falchuk KH.** The biochemical basis of zinc physiology. *Physiol Rev* 73: 79-118,
528 1993.

- 529 62. **Wang F, Kim BE, Dufner-Beattie J, Petris MJ, Andrews G, and Eide DJ.** Acrodermatitis
530 enteropathica mutations affect transport activity, localization and zinc-responsive trafficking of the mouse
531 ZIP4 zinc transporter. *Hum Mol Genet* 13: 563-571, 2004.
- 532 63. **Wang K, Zhou B, Kuo YM, Zemansky J, and Gitschier J.** A novel member of a zinc
533 transporter family is defective in acrodermatitis enteropathica. *Am J Hum Genet* 71: 66-73., 2002.
- 534 64. **Weaver BP, Dufner-Beattie J, Kambe T, and Andrews GK.** Novel zinc-responsive
535 post-transcriptional mechanisms reciprocally regulate expression of the mouse Slc39a4 and Slc39a5 zinc
536 transporters (Zip4 and Zip5). *Biol Chem* 388: 1301-1312, 2007.
- 537 65. **Wessells KR, and Brown KH.** Estimating the global prevalence of zinc deficiency: results based
538 on zinc availability in national food supplies and the prevalence of stunting. *PLoS One* 7: e50568, 2012.
- 539 66. **Wuehler SE, Pearson JM, and Brown KH.** Use of national food balance data to estimate the
540 adequacy of zinc in national food supplies: methodology and regional estimates. *Public Health Nutr* 8:
541 812-819, 2005.
- 542
- 543

FIGURE LEGENDS

Figure 1. Anti-Zip4 monoclonal antibody detects rat Zip4. *A*, The anti-Zip4 monoclonal antibody detected Zip4 in the rat small intestine. Membrane proteins prepared from the small intestine of rats fed a Zn-Low (L) or Zn-Suf (S) diet for 3 weeks were subject to immunoblot analysis. Zip4 protein was detected as both the full-length (~75 kDa) and the cleaved ectodomain (Ecto) (~38 kDa) only in membranes prepared from rats fed a Zn-Low diet by the anti-Zip4 monoclonal antibody (*left* panel). Note that Zip4 protein immunoreactivity detected by the anti-ZIP4 polyclonal antiserum (*right* panel) is similar, but the antiserum detects both the full-length (~75 kDa) and the processed Zip4 (corresponding to the remaining carboxyl-terminal region with eight TMDs lacking the ectodomain, ~40 kDa). The same membrane proteins were run on separate blots, and the blotted membrane was used for detection by either anti-Zip4 monoclonal antibody or anti-ZIP4 polyclonal antiserum. GRP78 was used as a loading control. *B*, The cleaved ectodomain of Zip4 was detected on membrane fractions after processing. Membrane proteins prepared from the small intestine of rats fed a Zn-Def diet were extracted in Na₂CO₃, and the pellet (Mem) and extract (Sup) were analyzed by immunoblot analysis. Note that the cleaved ectodomain of Zip4 was detected in Sup, but the full-length Zip4 and Znt4, both of which are integral membrane proteins, were detected in Mem. The Sup and Mem fractions extracted by PBS from the same membrane proteins were used as a control. The noncontiguous boxed lanes in the PBS and Na₂CO₃ panels were derived from the same blot. The specificity of the anti-ZnT4 monoclonal antibody was confirmed by verifying that the anti-ZnT4 antibody and commercial anti-HA antibodies (HA) detected the same bands in the immunoblot analysis using membrane proteins (20 µg) prepared from DT40 cells stably expressing HA-tagged human ZnT4 (+) and those not expressing (-) (*right* panels). The same blotted membrane was used for detection by each antibody after stripping. Calnexin was used as a loading control. In *A* and *B*, the representative results are displayed.

566

567 Figure 2. Rapid accumulation of Zip4 protein in the rat small intestine. A, Zip4 is expressed in the rat small
 568 intestine in response to zinc deficiency. Membrane proteins prepared from the small intestine of rats fed a
 569 Zn-Def (D), Zn-Low (L), or Zn-Suf (S) diet for up to 4 d were subject to immunoblot analysis. Znt4
 570 expression was not affected by dietary zinc levels, which was used as a loading control in D. CBB staining
 571 is also displayed as a loading control. B, ALP activity in the membrane proteins used in A. Data are
 572 expressed as means $\pm$ SD ($n = 5$, $*P < 0.05$, $**P < 0.01$). C, Plasma zinc concentrations at 1–4 d. Values are
 573 means $\pm$ SEM ($n = 5$, $**P < 0.01$). D, Accumulation of Zip4 in response to zinc deficiency is higher in the
 574 jejunum than in the duodenum. Membrane proteins prepared from the small intestine of rats fed a Zn-Def
 575 (D) or Zn-Suf (S) diet for up to 4 days were subject to immunoblot analysis. Znt4 was used as a loading
 576 control. In A and D, the lanes of day 0 are shown as “S” because rats were fed a Zn-Suf diet before being
 577 divided into three groups. The representative results are displayed.

578

579 Figure 3. Effects of dietary zinc deficiency on lactating dams. The dams were fed a Zn-Suf or Zn-Low diet
 580 from parturition (day 0 = parturition) until weaning (for 3 weeks). A, The amount of daily food intake for 3
 581 weeks. B, Body weights at 16 d gestation and each week after parturition. Values are means $\pm$ SEM ($n = 5$,
 582 $*P < 0.05$). C, Plasma zinc concentrations at 1 d after weaning their pups. Suf, Zn-Suf group; Low, Zn-Low
 583 group. Values are means $\pm$ SEM ($n = 5$, $**P < 0.01$).

584

585 Figure 4. Effects of dietary zinc deficiency on zinc transporter and zinc receptor expression on lactating
 586 dams. A, Immunoblot analysis of Zip4, Znt4, and GPR39 in the duodenum and jejunum. Membrane proteins
 587 were prepared from the duodenum or jejunum of lactating dams fed a Zn-Suf (S) or Zn-Low (L) diet for 3

weeks (at 1 d after weaning their pups) as shown in Figure 2. Two representative results of five are displayed. The noncontiguous boxed lanes were derived from the same blot in the duodenum and the jejunum. GRP78 was used as a loading control. *B*, ALP activity in the membrane proteins used in *A*. Data are expressed as means $\pm$ SD ($n = 5$, $**P < 0.01$) in the duodenum (*left*) or jejunum (*right*). *C*, Immunoblot analyses of Znt4 and Znt2 in membrane proteins prepared from mammary glands from the same rats used in *A*. Two representative results of five are displayed. *D*, ALP activity of the membrane proteins used in *C*. Data are expressed as means $\pm$ SD ($n = 5$, $*P < 0.05$). In *A* and *C*, GRP78 was used as a loading control.

Figure 5. Effects of zinc deficiency during lactation on pre-weanling pups. *A*, Change in body weights for 3 weeks. The weights of pups fostered by dams fed a Zn-Suf or Zn-Low diet were measured each week. *B*, Plasma zinc concentrations at the day of weaning (at 3 weeks). Suf, Zn-Suf group; Low, Zn-Low group. In *A* and *B*, values are means $\pm$ SEM ($n = 5$, $*P < 0.05$). *C*, Immunoblot analysis of Zip4, Znt4, and GPR39 in the duodenum or jejunum of the pups. Membrane proteins were prepared from the duodenum or jejunum of the pups. The noncontiguous boxed lanes were derived from the same blot in the duodenum and the jejunum. GRP78 was used as a loading control. Two representative results of five are displayed. *D*, ALP activity in the membrane proteins used in *C*. Data are expressed as means $\pm$ SD ($n = 5$, $**P < 0.01$) in the duodenum (*left*) or jejunum (*right*).

Figure 6. Effects of zinc deficiency during lactation on pups after weaning. *A*, Plasma zinc concentrations at 1 week after weaning (at 4 weeks). Values are means $\pm$ SEM ($n = 5$). *B*, Immunoblot analyses of Zip4, Znt4, and GPR39 in the duodenum or jejunum of the pups. Membrane proteins were prepared from the duodenum or jejunum of the pups of two groups. The noncontiguous boxed lanes were derived from the same blot in


610 the duodenum and the jejunum. GRP78 was used as a loading control. Two representative results of five are
611 displayed. C, ALP activity in the membrane proteins used in B. Data are expressed as means $\pm$ SD ($n = 5$) in
612 the duodenum (*left*) or jejunum (*right*).

613

614 Figure 7. Effects of zinc deficiency during lactation on plasma glucose homeostasis in the offspring. A,
615 Change in plasma glucose levels in male offspring of two groups from 4 weeks until 15 weeks ($n = 5$,
616 $*P < 0.05$). B, Plasma insulin levels in the same male offspring at 11 and 13 weeks. C, Change in plasma
617 glucose levels in female offspring in the same period as in A ($n = 5$).

618

A



B




Fig. 1






Fig. 2

A



B



C




Fig. 3




Fig. 4




Fig. 5






Fig. 6

A



B



C




Fig. 7