

Cohomology of conical symplectic resolutions

Kohei Hatano

10/22 2020

A set of navigation icons typically found in Beamer presentations, including symbols for back, forward, search, and other slide controls.

Kohei Hatano

Cohomology of conical symplectic resolutions

10/22 2020

1 / 7

Definition

Definition

A smooth symplectic variety (X, ω) with an action of $\mathbb{S} = \mathbb{C}^*$ is a conical if X is the following conditions,

- 1 $\phi : X \rightarrow X_0 = \operatorname{Spec} \mathbb{C}[X]$ is projective birational.
- 2 $s^* \omega = s^n \omega$ for $s \in \mathbb{C}^*$.
- 3 $\mathbb{C}^*$ acts on $\mathbb{C}[X]$ with only non-negative weights and $\mathbb{C}[X]^{\mathbb{C}^*} = \mathbb{C}$

Example

Let X be a crepant resolution of $\mathbb{C}^2/G$, where a group G is a finite subgroup of $SL_2(\mathbb{C})$. Then the Hilbert scheme on n -points of X is a conical symplectic resolution.



Kohei Hatano

Cohomology of conical symplectic resolutions

10/22 2020

2 / 7

Symplectic dual

Consider a conical symplectic resolution X with a Hamiltonian action of $\mathbb{T} = \mathbb{C}^*$, commuting $\mathbb{S}$, such that $X^{\mathbb{T}}$ is finite.

Let $\mathbb{W}$ be a Weyl group of the group of Hamiltonian symplectic morphisms of X and W be a Namikawa Weyl group of X .

Definition

A symplectic duality from X to $X^!$ consists of

- 1 There exists a order reversing bijection from the index set of $X^{\mathbb{T}}$ to the index set of $(X^!)^{\mathbb{T}}$.
- 2 $W \simeq \mathbb{W}^!$ and $W^! \simeq \mathbb{W}$
- 3 etc

Navigation icons: back, forward, search, etc.

Examples

- The Hilbert scheme on n -points of $\mathbb{C}^2$ is self dual.
- An affine A type quiver variety and other affine A type quiver variety are symplectic dual.
- A hypertoric variety and other hypertoric variety are symplectic dual.

Navigation icons: back, forward, search, etc.

Conjecture

Conjecture (Hikita,2015)

If $X \rightarrow X_0$ and $X' \rightarrow X'_0$ are symplectic dual, then we have the following graded algebra isomorphism:

$$H^*(X) \simeq C[X'_0{}^{\mathbb{T}}]$$

$$H^*(X') \simeq C[X_0{}^{\mathbb{T}}]$$

Theorem (Hikita,2015)

In the case of the Hilbert scheme on n -points of $\mathbb{C}^2$ and a hypertoric variety, type A S3-variety, this conjecture is true.

Navigation icons: back, forward, search, etc.

Case of the framed moduli space

Let $M(r, n)$ be the framed moduli space of torsion free sheaves on $\mathbb{P}^2$ with rank r and $c_2 = n$. Then $M(r, n)$ is a conical symplectic resolution .

Theorem (Nakajima-Yoshioka,2003)

we have the following equations:

$$\sum_d \dim H^d(M(r, n)) t^d = \sum_{(Y_1, \dots, Y_r)} t^{2(m - (\sum_j jl(Y_j)))}$$

where $(Y_1, \dots, Y_r)$ is r -tuples Young diagram running over $\sum |Y_j| = n$, $l(Y_j)$ is a length of Young diagram Y_j

Navigation icons: back, forward, search, etc.

Main result

Let X_r be a crepant resolution of $\mathbb{C}^2/(\mathbb{Z}/r\mathbb{Z})$.

Theorem (Ben-Licata-Proudfoot-Webster, 2014)

$\mathrm{Hilb}^n(X_r)$ and $M(r, n)$ are symplectic dual.

Theorem (Hatano)

There exists a graded vector space isomorphism

$$H^*(M(r, n)) \simeq \mathbb{C}[(\mathrm{Hilb} X_r)_0]^{\mathbb{C}^*}.$$