

Biomimetic Oxidation of β -O-4 Lignin Model Compound with *tert*-Butylhydroperoxide and Fe(III)-Octacarboxyphthalocyanine*¹

Yoshifumi TANIHARA*², Takefumi HATTORI*²,
Hirofusa SHIRAI*³ and Mikio SHIMADA*²

(Received June 1, 1994)

Keywords: Biomimetic oxidation, phthalocyanine, catalyst, ligninase model, cleavage reaction.

Lignin-peroxidase mimetic oxidation of a lignin model compound was first reported by Shimada et al.¹⁾ with synthetic and natural porphyrin iron-complexes; the one-electron oxidation mechanism for the enzymatic reaction has been proposed²⁾. A hemin catalyst mimicked the lignin peroxidase (Lip) which catalyzes the C α -C β bond cleavage of β -1 model substrate (accompanied by oxygenation and deuterium retention)^{2,3)}, and the aromatic ring cleavage of veratryl (3, 4-dimethoxybenzyl) alcohol, yielding γ - and δ -lactone products^{4,5)}.

Recently, several laboratories have been investigating similar biomimetic oxidations of lignin models with more sophisticated water-soluble synthetic porphyrins⁶⁻¹¹⁾. It is particularly interesting to note that the selective oxidative cleavage of 1, 2-dimethoxyarenes to muconic diesters with relatively high yield was catalyzed by Fe(III)-sulfonated-tetrakis (pentafluorophenyl) porphyrin¹¹⁾.

We report here the first example of C α -C β bond cleavage and ring cleavage of β -O-4 model lignin model compound catalyzed by a water soluble Fe(III)-octacarboxyphthalocyanine (OcPc) complex¹²⁾, although the oxidation of monomeric lignin model compounds with various water-soluble metallophthalocyanines has been previously reported¹³⁾.

The reaction mixture (2 ml of water adjusted to pH 3) containing the β -O-4 model substrate (1, 10 μ mol), Fe-OcPc (1 μ mol), and *tert*-butylhydroperoxide (t-BuOOH), was incubated at room temperature for 30 min. The control system, lacking the Fe-OcPc catalyst, was incubated in the same way. After the incubation oxidation products were extracted with ethyl acetate and analyzed with GC-MS after acetylation of the dried

*¹ Part of this work was presented in the 36th Lignin symposium, held in Sapporo, Japan, in October, 1991.

*² Laboratory of Biochemical Control in this Institute.

*³ Department of Functional Polymer Science, Faculty of Textile Science and Technology, Shinshu University, Ueda-city, 386 Japan.

extracts; the products (II–VII) were identified by comparison of their spectra with those of the corresponding authentic compounds.




Fig. 1. LiP-mimetic Fe-octacarboxyphthalocyanine catalyzes oxidation of β -O-4 lignin model compound (I).

Fig. 1 shows that Fe-OcPc catalyzes the C α -C β cleavage of I (yielding the products III and IV), the β -O-4 ether cleavage, (yielding the product V), and the cleavage of B-ring of I (yielding products VI and VII). The control reaction system did not yield these products under the reaction conditions used, which indicates that t-BuOOH did not cause such cleavage reactions. It is noteworthy that VII was identified for the first time from the products formed from the β -O-4 lignin substructure model. Therefore, this finding prompted us to examine a possible formation of VII even in the Lip system. Eventually, we confirmed that VII was also produced from the Lip-catalyzed oxidation of I. The alternative experiment also demonstrated that the Fe-OcPc system is capable of catalyzing the ring cleavage of veratryl alcohol, yielding γ - and δ -lactones as the ring-opened products that were also produced from the Lip system⁴⁾.

These results clearly indicate that Fe-OcPc system like the metalloporphyrin systems simulates the activity of Lip in the cleavage patterns of both dimeric β -O-4¹⁴⁾, and monomeric lignin model substrates¹⁵⁾. Furthermore, it is noteworthy that Fe-OcPc also catalyzed the C α -C β bond cleavage of β -O-4 substrate (II), yielding IV and 4-O-ethylvanillic acid¹⁶⁾ because II is not utilized as substrate by the LiP system¹⁷⁾. Since the control reaction system did not produce these cleaved products, the possibility of participation of Vayer-Villiger type reaction was eliminated.

References

- 1) M. SHIMADA, T. HABE, T. UMEZAWA, T. OKAMOTO and T. HIGUCHI: *Biochem. Biophys. Res. Commun.*, **122**, 1247–52 (1984).
- 2) T. HABE, M. SHIMADA, T. OKAMOTO, B. PANIJPAN and T. HIGUCHI: *J. Chem. Soc. Chem. Soc. Chem. Commun.*, **1985**, 3123–52 (1985).
- 3) M. SHIMADA, T. HABE, T. OKAMOTO, B. PANIJPAN and T. HIGUCHI: *Holzforsch.*, **41**, 277–285 (1987).
- 4) M. SHIMADA, T. HATTORI, T. UMEZAWA, T. HIGUCHI and K. UZURA: *FEBS Lett.*, **221**, 327–331 (1987).
- 5) T. HATTORI, M. SHIMADA, M.S.A. LEISOLA, A. FIECHTER and T. HIGUCHI: *Agri. Biol. Chem.*, **52**, 879–880 (1988).
- 6) D. DOLPHIN, T. NAKANO, T.E. MAIONE, T.K. KIRK and R.L. FARRELL: “Lignin, Enzymic and Microbial Degradation” (Ed. E. Odijer) pp. 157–162 (1987).
- 7) P.S. SKERKER, R.L. FARRELL, DOLPHIN, F. CUI and T. WIJSEKERA: “Biotechnology in Pulp and Paper Manufacture”, Butterworth-Heinemann (Eds. T.K. Kirk and H.m. Chang), pp. 203–210 (1990).
- 8) F. CUI, D. DOLPHIN, T. WIJSEKERA, R. FARRELL and P. SKERKER: *ibid.*, pp. 481–491 (1990).
- 9) G. LABAT and B. MEUNIER: *J. Org. Chem.*, **54**, 5008–11 (1989).
- 10) H.C. TUNG and D.T. SAWYER: *FEBS Lett.*, **311**, 165–168 (1992).
- 11) I. ARTAUD, H. GRENBERG and D. MANSUY: *J. Chem. Soc. Chem. Commun.*, **1992**, 1036–38.
- 12) H. SHIRAI, A. MARUYAMA, J. TAKANO, K. KOBAYASHI and N. HOJO: *Makromol. Chem.*, **181**, 565 (1980).
- 13) Z. WEIMING and W.T. FORD: *J. Mol. Catal.*, **78**, 367–378 (1993).
- 14) T. UMEZAWA: *Wood Research*, No. **75**, 21–79 (1988).
- 15) T. HATTORI: *ibid.*, No. **78**, 15–73 (1992).
- 16) Y. TANIHARA, T. HATTORI, H. SHIRAI, and M. SHIMADA: The 36th Lignin Symposium (Sapporo), pp. 103–106 (1991).
- 17) T.K. KIRK, M. TIEN, P.J. KERSTEN, M.D. MOZUCH and B. KALYANARAMAN: *Biochem. J.*, **236**, 279–287 (1986).