

Biological Sciences: Evolution

Paleontological and developmental evidence resolve the homology and dual embryonic origin of a mammalian skull bone, the interparietal

Daisuke Koyabu^{a,b,1}, Wolfgang Maier^c, Marcelo R. Sánchez-Villagra^{a,1}

^aPalaeontological Institute and Museum, University of Zürich, Karl Schmid-Strasse 4, 8006 Zürich, Switzerland; ^bThe Kyoto University Museum, Yoshida Honmachi, 606-8501 Kyoto, Japan; ^cUniversity of Tübingen, Auf der Morgenstelle 28, 72076 Tübingen, Germany

¹To whom correspondence may be addressed.

Daisuke Koyabu: Palaeontological Institute and Museum, University of Zürich, Karl Schmid-Strasse 4, 8006 Zürich, Switzerland, TEL 044 634 2447, koyabu@pim.uzh.ch
or

Marcelo R. Sánchez-Villagra: Palaeontological Institute and Museum, University of Zürich, Karl Schmid-Strasse 4, 8006 Zürich, Switzerland, TEL 044 634 2342, m.sanchez@pim.uzh.ch

Author contributions: D.K., W.M., and M.R.S.-V. designed research and collected data; D.K. analyzed data; and D.K. and M.R.S.-V. wrote the paper. The authors declare no conflict of interest.

key words: morphological evolution, embryology, synapsids, neural crest, mesoderm

1 **Abstract**

2 The homologies of mammalian skull elements are now fairly well established, except for the
3 controversial interparietal bone. A previous experimental study reported an intriguing mixed
4 origin of the interparietal: the medial portion being derived from the neural crest cells,
5 whereas the lateral portion from the mesoderm. The evolutionary history of such mixed origin
6 remains unresolved, and contradictory reports on the presence or absence and developmental
7 patterns of the interparietal among mammals have complicated the question on its homology.
8 Here we provide a novel perspective on the evolutionary identity of the interparietal, based on
9 a comprehensive study across more than 300 extinct and extant taxa, integrating
10 embryological and paleontological data. Although the interparietal has been regarded as being
11 lost in various lineages, our investigation on embryos demonstrates its presence in all extant
12 mammalian “orders.” The generally accepted paradigm has regarded the interparietal as
13 consisting of two elements that are homologized to the postparietals of basal amniotes. The
14 tabular bones have been postulated as being lost during the rise of modern mammals.
15 However, our results demonstrate that the interparietal consists not of two but of four
16 elements. We propose that the tabulars of basal amniotes are conserved as the lateral
17 interparietal elements, which quickly fuse to the medial elements at the embryonic stage, and
18 that the postparietals are homologous to the medial elements. Hence, the dual developmental
19 origin of the mammalian interparietal can be explained as the evolutionary consequence of the
20 fusion between the crest-derived “postparietals” and the mesoderm-derived “tabulars.

1 \body

2 The evolutionary identity or homology of most mammalian skull elements is now
3 well established (1-3), although many contradictory statements still exist on the interparietal,
4 a dermal skull roof element situated between the parietal and supraoccipital (e.g., 4, 5-8).
5 Several authors have suggested a homology of the paired mammalian interparietals to the
6 paired postparietals in more basal reptiles (6, 9). However, inconsistent patterns reported
7 among mammals (7, 10, 11), and even humans (12), and a unique mixed embryonic origin (1)
8 make the question on homology of the interparietal an unresolved issue of vertebrate anatomy.

9 An experimental embryological study in the mouse by Jiang et al. (13)
10 demonstrated a striking dual developmental origin of the interparietal: the median portion is
11 derived from the neural crest cells, whereas the lateral portion is derived from the mesoderm.
12 This report has triggered a renewed interest among developmental biologists in the evolution
13 of the interparietal in amniotes (1, 14, 15). Although integrating fossil record and comparative
14 anatomical data with newly gained experimental findings should provide relevant information
15 for the current discussions on the origin of skull roof bones, it has been hampered by
16 terminological inconsistencies, even within the stem mammal (synapsid) lineage, and
17 unsolved homologies across major living groups (4, 6, 7, 9, 11, 16). In humans, the
18 interparietal develops from two pairs of ossification centers, one medial pair and one lateral
19 pair, eventually fusing seamlessly to the supraoccipital (17). In contrast to the human pattern,
20 it is generally regarded for mammals that the interparietal arises in development as a pair of
21 bones that soon fuse at the midline to form a single bone (6, 8, 18). However, the absence of

1 the interparietal has been pointed out in marsupials (11, 18), xenarthrans (19), shrews (4),
2 seals (10), hippopotamuses (10), and pangolins (11). The classic work of de Beer (20)
3 reported that the interparietal is lacking in monotremes, bandicoots, armadillos, and pigs.

4 The paucity of appropriate embryonic material across mammalian species has
5 undoubtedly hindered evaluations as to the primordial anlage, general presence, and variation
6 of the interparietal. As exemplified in humans, the interparietal generally fuses with
7 neighboring bones early in the perinatal period (12). Because the majority of work has been
8 based on mature specimens, previous observers could have been misled by the early fusion of
9 the interparietal with other bones. Here, with the unique availability of wide taxonomic and
10 ontogenetic sampling across all extant mammalian orders and major non-mammalian fossil
11 taxa, we address the issues of variation, patterns, and homology of the interparietal. Our
12 results question the generally accepted homology hypothesis, and we suggest an alternative
13 model that explains the intriguing dual tissue origin of the interparietal. Our integrative study
14 provides a bridge between paleontology and developmental biology and a synthetic
15 understanding of the dermal skull roof of vertebrates.

16 17 **Results**

18 Below we present an overview of our findings. A full description of our detailed
19 survey is given in the supplementary information (SI Appendix). All 51 non-mammalian
20 synapsid species examined possessed the postparietal, which has been posited as homologous
21 to the interparietal (9), either unpaired or paired (Fig. S3 and Table S1). The Jurassic

1 docodont mammal *Haldanodon exspectatus* reportedly possesses the interparietal (21), as do
2 the extant monotremes including the platypus *Ornithorhynchus* (22) and the long-beaked
3 echidna *Zaglossus* (23). A small medial membrane bone, a remnant of the interparietal, is
4 found in the short-beaked echidna *Tachyglossus aculeatus* (24). Marsupials reportedly lack
5 the interparietal (11), but we confirmed its presence in various species (*Trichosurus*,
6 *Monodelphis*, *Didelphis*, *Macropus*, and *Sminthopsis*). Contrary to the reported absence of the
7 interparietal in *Monodelphis* (25), in the examined embryonic series of the gray short-tailed
8 opossum *M. domestica*, both the supraoccipital and interparietal are present. Here, the
9 interparietal was found to be unpaired, paired, tripartite, and quadripartite, depending on the
10 stage (Fig. S4).

11 We recorded the presence of dermal interparietal bone in four xenarthran species
12 including the long-nosed armadillo *Dasypus hybridus* (Fig. 1A and Table S1). Our
13 investigation confirmed the presence of the interparietal across afrotherians (Figs. S5A-B, S6,
14 and Table S1). We identified unpaired, paired, and tripartite interparietals in the tree hyrax
15 *Dendrohyrax arboreus*. The interparietal in the rock hyrax *Procavia capensis* was found to be
16 unpaired, paired, tripartite, and quadripartite. In some individuals, the lateral elements fuse
17 earlier to the supraoccipital rather than to the medial elements (Fig. S5B). Tripartite
18 interparietal elements were described for the cape golden mole *Chrysochloris asiatica* (2). In
19 sirenians (*Dugong dugon*, *Trichechus manatus*, and *T. senegalensis*) the interparietal showed
20 a maximum of four elements. Among extinct South American notoungulates, *Toxodon*

1 *burmeisteri*, *Typotherium cristatum*, *Nesodon imbricatus*, and *Pachyrucus* were confirmed to
2 possess the interparietal.

3 We observed the interparietal universally in Euarchontoglires (Figs. 2I-J, S9D-F,
4 and Table S1). In the kangaroo rat *Dipodomys*, the interparietal was essentially formed from
5 four ossification centers (26). One median and two lateral elements were found in various
6 stages of the degu *Octodon degu* (Figs. 3B-C and 4) and in the Patagonian cavy *Dolichotis*
7 *patagonum* (Figs. 3D-E). Among primates, the interparietal was widely found in perinatal
8 individuals (Fig. 2J and Table S1). In humans, the interparietal develops from four elements
9 (12, 27).

10 Paired interparietals were recorded in lipotyphlans (Figs. 1B, 2A, and Table S1).
11 There was no previously published information on the presence of the interparietal in moles.
12 Nevertheless, we found thin paired membranous interparietals, which initiate their ossification
13 from the dorsal tip of the supraoccipital and then grow gradually toward the rostral direction,
14 in both the Iberian mole *Talpa occidentalis* (Fig. S10A) and the European mole *T. europaea*.
15 Similarly, we found the developing dermal interparietal in the coast mole *Scapanus orarius*
16 (Figs. S10C-D) and the Japanese mole *Mogera wogura*. Augier (4) noted that shrews have no
17 interparietal, but we recorded this bone in various shrews (*Sorex araneus*, *S. unguiculatus*,
18 and *Crocidura attenuata*) (Fig. 1B).

19 The interparietal is present in artiodactyls, cetaceans, perissodactyls, carnivorans,
20 bats, and pangolins (Figs. 2B-H, S9A-C, and Table S1). The bottle-nosed dolphin *Tursiops*
21 *truncatus* possesses an expanded median and two lateral interparietal elements (Fig. 2G).

1 Eales (28) described the greatly expanded four-element interparietal in the narwhal *Monodon*
2 *monoceros*. The cow *Bos primigenius* exhibited quadripartite interparietal (Fig. 2H). The
3 tripartite interparietal is reported for the yak *B. grunniens* (29). Wegner (1960) documented
4 the tripartite interparietal in the jaguar *Panthera onca* and cougar *Puma concolor*. Although
5 Giebel (10) noted that seals lack the interparietal, we observed that prenatal skulls of fur seals
6 *Arctocephalus* (Fig. S5G) and *Callorhinus ursinus* (Fig. S8B) display enlarged interparietals
7 and the supraoccipital is extremely reduced in size compared with the interparietal, which
8 dominates the caudal portion of the vault. The earless seal *Phoca* is reported to exhibit the
9 lateral interparietal element (2, 25). Gregory (30) reported that pangolins lack the interparietal,
10 but we found its presence in an immature individual of the Chinese pangolin *Manis*
11 *pentadactyla* (Fig. S5E).

12

13 **Discussion**

14 **Basic structure and evolutionary patterns of the interparietal.** The presence of the
15 interparietal is almost universal among extant mammals. Previous reports have noted that
16 there is no interparietal in monotremes (18, 20), marsupials (11, 18), xenarthrans (11, 19, 20),
17 moles and shrews (4), seals (10), and pangolins (30). However, we documented the presence
18 of the interparietal in monotremes (*Ornithorhynchus* and *Tachyglossus*), marsupials (23
19 species), xenarthrans (*Bradypus*, *Dasybus*, *Tamandua*, and *Cyclopes*), moles (*Talpa*, *Mogera*,
20 and *Scapanus*), shrews (*Crocidura*, *Suncus*, and *Sorex*), seals (*Phoca*, *Arctocephalus* and
21 *Callorhinus*), and pangolins (*Manis*) (Table S1). Previous studies may have overlooked the

1 presence of the interparietal because of its early fusion to adjacent bones, either the
2 supraoccipital or the parietal. The membranous bone found rostral to the endochondral
3 supraoccipital in these species is interpreted to be the interparietal. The case of some shrews
4 provides an instructive example of the importance of investigating embryonic materials and
5 understanding the topological relationships of calvarial bones. The expansion of the
6 interparietal and the rudimentary presence of the supraoccipital, which was often found in
7 shrews, may have misled anatomists to regard the true interparietal as the supraoccipital and
8 to consider the interparietal to be lacking in shrews (for instance, Fig. 1B). Kuratani (31)
9 observed the expansion of the dermal interparietal bone and the unique case of possible lack
10 of the cartilaginous supraoccipital in the Asian house shrew *Suncus murinus*. In the common
11 shrew *Sorex araneus* (Figs. S7C-D and S11) and the long-clawed shrew *S. unguiculatus* (Fig.
12 S12), we observed that the membranous interparietal dorsally covers the cartilaginous
13 supraoccipital, forming a double-layered structure with the supraoccipital. In the Asian gray
14 shrew *Crociodura attenuata*, the presence of the dermal interparietal was obvious (Fig. 1B).

15 While the Jurassic docodont *Haldanodon* is reported to possess the interparietal
16 (21), its absence has been noted in other basal mammalian clades, including
17 morganucodontids, multituberculates, *Sinoconodon*, and *Vincelestes* (32). Yet, as shown in
18 the present study, the “absence” of the interparietal cannot be generalized in these species,
19 because the interparietal may fuse early with other bones or it may be overlooked because of
20 the poor preservation of sutures in fossils. This is especially true for primates, in whom the

1 interparietal fuses with the supraoccipital at very early embryonic stages [c.f. Hershkovitz
2 (33), who commented that callitrichid monkeys lack the interparietal].

3 Generally, the interparietal has been regarded as comprising paired elements that
4 often fuse with each other to form an unpaired bone (8, 9, 11, 18). In contrast, a maximum of
5 four ossification centers were recognized from our survey in artiodactyls (Fig. 2H), cetaceans,
6 rodents, sirenians, hyraxes, platypuses, opossums, and humans (Table S1). The four elements
7 fusing with one another become paired or unpaired, depending on the species or
8 developmental stage. The fact that the lateral elements fuse to the medial elements at the
9 embryonic stage in many taxa makes it critically difficult to identify the lateral portions of the
10 interparietal. Tripartite interparietals were found in **opossums (Fig. 3A)**, rodents (Figs. 3B-E
11 and 4), cetaceans (Fig. 2G), felids, seals, yaks, manatees, tenrecs, hyracoids, golden moles,
12 and humans. In these taxa, the median bone is an enlarged element formed by the fusion of
13 two medial elements, and the lateral elements are separated from this median bone. Similarly,
14 extinct notoungulates also exhibit tripartite interparietals.

15 The facts presented above indicate that the interparietal consists of four basic
16 elements: one medial pair and one lateral pair (Fig. 5B). In most living mammals, the lateral
17 element fuses first with the medial element, forming paired interparietals, and later in
18 ontogeny, the fusion occurs between the two composites, forming an unpaired interparietal.
19 Nevertheless, in some species (rodents, cetaceans, tenrecs, hyracoids, felids, seals, manatees,
20 yaks, golden moles, and humans), the medial elements initially fuse with each other, while the
21 lateral elements remain separate from the medial element, producing a tripartite interparietal

(Fig. 4). Such a condition is comparable to that in most non-mammalian synapsids, in whom paired tabulars are present but are separated from the median postparietal (e.g., see *Titanophoneus* in Fig. 5A).

Homology of the mammalian interparietal. Both the postparietal and the tabular are widely present in various tetrapods, including lissamphibians, diapsids, and non-mammalian synapsids [the terminology of the postparietal has been variable depending on the authors, such as “retroparietal (34),” “dermosupraoccipital (35),” “dermooccipital (16),” and “parieto-extrascapular (36)”] (Table S1). The stem condition of lissamphibians, as illustrated by some dissorophids, exhibits paired postparietals and tabulars (37). Although modern lissamphibians were previously suggested to have lost the postparietals (38), they are present in some living anurans such as the spadefoot toad *Pelobates* (39) and fire-bellied toad *Bombina* (40). Basal amniotes exhibit paired postparietals and tabulars (41) (Fig. S1). In contrast, no extant sauropsid reportedly possesses the postparietals, with the exception of the American alligator *Alligator mississippiensis* (42). Extant chelonians (turtles, tortoises, and terrapins) (37), squamates (18, 37, 43-45), and birds (46) lack the postparietals. However, the presence of paired postparietals is common among fossil diapsids (37, 41) (Figs. S1 and S2).

Here, we hypothesize that the medial interparietal ossification centers and lateral ossification centers of mammals are homologous to the postparietals and the tabulars, respectively, of ancestral forms (Fig. 5A). The spatial arrangements of these bones, the number of elements, and the intramembranous nature of these bones lend support to this

1 postulate. The topological correspondence of the posttemporal canal in non-mammalian
2 synsids and in primitive mammals also gives support to this hypothesis. In non-mammalian
3 synsids, the tabular generally forms an arch or roof over the posttemporal canal (Fig. S3B,
4 D, F) (see also ref. 47). This canal is located at the junction of the tabular, squamosal, and
5 mastoid in *Morganucodon* (48). The posttemporal canal is found in monotremes, opossums,
6 and armadillos (49). For example in opossums, this canal is located at the junction of the
7 lateral interparietal, squamosal, and mastoid as in *Morganucodon* (Fig. S4) (see also ref. 50,
8 51).

9 Frequent fusions between the tabular and postparietal in non-mammalian
10 synsids suggest the strong coupling trend of these bones. Across the observed non-
11 mammalian synsids, 33 species exhibited one median postparietal and two tabulars (e.g.,
12 *Titanophoneus* in Fig. 5A), whereas 17 species showed only the median “postparietal” (Fig.
13 S3 and Table S1). In the later ones, the tabular is not lost but is fused to the postparietal.
14 Given the presence and shape of the tabular and postparietal in juveniles of *Cistecephalus*
15 *microrhinus*, a Late Permian anodont therapsid, the tabular is fused to the true postparietal
16 to form an enlarged median “postparietal” in mature individuals of this species (52-54), which
17 resembles the single interparietal of various extant mammals (Fig. 2B-C, and J).

18 The interparietal has been regarded as an intramembranous bone formed dorsally
19 to the tectum posterius, generally developing as a pair (7). Goodrich (9) suggested that the
20 origin of the mammalian skull from a reptilian-like condition was brought about by the
21 evolutionary loss of the prefrontal, postfrontal, postorbital, and quadratojugal. In his view, the

1 postparietal pair has remained as the interparietal pair. Similarly, Gross and Hanken (1) and
2 Morris-Kay (3) have noted that the mammalian interparietal is homologous to the postparietal
3 bone of “archaic reptiles.” In contrast, our investigation demonstrates that the interparietal is
4 constituted of four rather than just two elements. Such a developmental pattern of the modern
5 mammalian interparietal from four dermal ossification centers is not fully explained, and thus
6 this generally accepted paradigm on the homology of the interparietal (6, 8, 9, 18, 20) cannot
7 be supported. Furthermore, the “interparietal” frequently documented in non-mammalian
8 synapsids (e.g., 41, 53) is inconsistent. In cases where the bone labeled “interparietal” is
9 isolated from the tabular, calling it the “postparietal” or “medial interparietal” would be
10 appropriate.

11 Although the tabular is fused to the medial interparietal in some species of
12 Dicynodontia (e.g., *Dicynodon*, *Diictodon*, and *Emydops*), it is clearly present and
13 proportionally large in many other non-mammalian synapsids (Table S1). Among extinct
14 basal mammals, the tabular is present in *Morganucodon*, but its relative size is much reduced
15 compared to those in non-mammalian synapsids (48). *Haldanodon* shows an unpaired
16 interparietal. The interparietal is not reported in multituberculates, but we assume their
17 interparietal is fused to the supraoccipital. Thus, it appears that in crown-group mammals the
18 tabular became reduced in size and fused to the medial interparietal.

19 According to Sidor (55), who studied the variation and evolution of cranial
20 elements in synapsids leading to the living mammal crown group, there are, surprisingly, no
21 fossil species that lack the tabular, whereas the extant lineage studied (*Ornithorhynchus*,

1 *Didelphis*, and *Homo*) were reported to lack this bone. Nonetheless, as in the case of *Homo*,
2 the lateral interparietal pair, which we hypothesize to be homologous to the tabulars, is
3 identifiable in embryos (Fig. S13) (12, 27). Although the “simplification trend” of the
4 mammalian skull (55) still holds, the general assumption that the tabular is lost in the modern
5 mammalian lineage (18, 21, 55) is questionable. Broom (35) and von Huene (56) noted that
6 the postparietal found in fossil amphibian-like land vertebrate *Diadectes* (Fig. 5A) should be
7 considered homologous to the postparietal of the stegocephalians. Furthermore, Jarvik (57)
8 proposed that the postparietal and tabular of basal tetrapods are homologous to the medial and
9 lateral extrascapulars, respectively, of osteolepiform fishes. If these scenarios are to be
10 supported, the medial and lateral extrascapulars of osteolepiforms should be regarded as
11 conserved in mammals as the four basic elements constituting the interparietal.

12 As noted earlier, dual origin of the interparietal has been reported in the mouse,
13 the medial portion being derived from the neural crest cells and the lateral portion from the
14 mesoderm (Fig. 5D) (13). At least four segments are necessary to account for such mixed
15 arrangements (Fig. 5C). We found (see also ref. 26) that the rodent interparietal originates
16 from four ossification centers. Therefore, the two medial interparietal elements may
17 correspond to the neural crest cell-derived portion and the lateral elements to the mesoderm-
18 derived portion. Thus, the dual origin found for the interparietal (13) could be regarded as the
19 evolutionary consequence of the fusion between the crest-derived “postparietals” and the
20 mesoderm-derived “tabulars.”

1 Gross and Hanken (1) noted that the location of the interparietal in the mouse
2 corresponds to that of the “parietal” in the avian skull, a bone for which a controversy exists
3 on its neural crest (58) or mesodermal origin (59). From embryological (i.e., dual
4 mesenchymal origin) and topographical (or anatomical) grounds, Noden and Schneider (59)
5 proposed that the bone conventionally labeled “parietal” in the chick would more
6 appropriately be termed “interparietal” and “frontal” as “frontoparietal.” In the chick, the
7 boundary between frontal and parietal overlies the otic capsule, whereas in the mouse, this
8 boundary occurs over the orbit (59). Although the developmental origin of the avian
9 “parietal” [the “interparietal” of Noden and Schneider (59)] is controversially reported to be
10 exclusively derived from the mesoderm (59) or exclusively from the neural crest cells (58), no
11 studies have been conducted to test the dual embryonic origin for the avian “parietal” (1). If
12 the avian “parietal” is validated to be derived from both the neural crest cells and the
13 mesoderm, it will indicate that the mammalian interparietal is homologous to the avian
14 “parietal.”

15 The early ontogenetic fusion of the lateral and medial interparietal elements in
16 living mammals may have been acquired in the synapsid lineage with encephalization and
17 changes in molecular signaling. There is now good evidence that calvarial suture closures are
18 associated with the expression of transforming growth factor β receptor genes (*Tgfr1* and
19 *Tgfr2*) and fibroblast growth factor receptor genes (*Fgfr1* and *Fgfr2*) (60). Mutations in these
20 molecules and the increased signaling of transforming growth factor β (TGF- β) or fibroblast
21 growth factor (FGF) provokes early suture closure in humans—the so-called craniosynostosis

1 syndrome (60, 61). However, inactivation of *Tgfb2* in the mouse leads to the absence of the
2 interparietal and parietal (62), suggesting the critical role of TGF- β signaling in the dermal
3 skull roof development (see also ref. 63). In addition, recent findings demonstrate that *Tgfb2*
4 activity is required for axon formation and neuronal migration in the developing mammalian
5 neocortex (64) and that expression of *Fgfr1* and *Fgfr2* determines brain size (65, 66).
6 Richtsmeier et al. (67) have pointed out the analogy between human craniosynostosis and the
7 simplification trend of cranial bone elements in synapsid evolution. Given the phenotypic
8 integration of brain and neurocranium growth, they suggested the possible shared genetic
9 pathways for brain development and skull tissue growth. It is quite intriguing to recognize
10 that humans, which exhibit the most aberrant case of encephalization in vertebrate evolution
11 (68) are, at the same time, the species that shows one of the most complete and accelerated
12 fusion of the interparietal among mammals. Investigators may profit from testing whether
13 encephalization and suture fusions in the skull roof are pleiotropically caused by changes in
14 TGF- β and/or FGF signaling.

15 In conclusion, the discovery of the four-element nature of the interparietal (Fig.
16 5B) provides a synthetic understanding of the dermal skull roof of mammals. The “three-
17 element interparietal” can be interpreted as consisting of one median bone, produced by the
18 fusion of the medial interparietal elements, and two lateral interparietal elements. The “two-
19 element interparietal” consists of a right and a left bone, each of which is the fusion of the
20 medial and lateral interparietal elements. The “one-element interparietal” is interpreted as the
21 fusion of all four elements (two medial and two lateral interparietal elements) into one median

bone. However, it must be noted that the lateral interparietal elements may be fused earlier to the supraoccipital or to the parietal than to the medial interparietal in some cases (e.g., hyraxes). In terms of homology (Fig. 5A), the lateral interparietal elements derived from the mesoderm and the medial elements derived from the neural crest cells appear to correspond to the tabular and postparietal elements, respectively, of basal tetrapods. If future experimental studies on species other than the mouse similarly confirm the dual origin of the interparietal, our hypothesis shall be further supported.

Lastly, it is worth noting that wormian bones are rarely found anterior to the true interparietal in hyraxes, horses, cows, and primates (Fig. S14). This anomalous dermal bone has been referred to as “os Incae” in humans (12) and “preinterparietal” in other mammals (7). Whether the preinterparietal is a simple developmental anomaly or a trait obtained *de novo* in some mammals remains uncertain. Future investigations on developmental and evolutionary patterns of the preinterparietal are needed to further understand the anatomy of the mammalian skull.

Methods

We investigated more than 3,000 skulls of 318 species in the context of studies on ossification sequences and sutural fusion in mammals (69, 70) and conducted a comprehensive review on literature describing amniote skulls (Table S1). Fetal or perinatal stages were studied to document whether the interparietal was recognizable and to record the number of interparietal ossification centers and contacts. Following de Beer (20) and Wegner

(7), we defined the interparietal as an intramembranous bone that is formed dorsally to the tectum posterius and posteriorly to the parietal. Documentation was based on macerated skulls and cleared and stained specimens as well as on microtomographic scans (Anthropological Institute, University of Zurich, μ CT80, Scano Medical, Bassersdorf, Switzerland). 3D visualization was conducted in Amira 5.3 (Visage Imaging, Richmond, Australia). A missing distinctive interparietal could be due to real loss or primordial fusion with one of the adjacent bones. Therefore, we restricted our observations to the recording of interparietals and omitted any speculation based on the “absence” of clear signs of this element.

Acknowledgements

We thank F. Abdala, A. Forasiepi, H. Higashiyama, T. Martens, K. Ota, T. Scheyer, I. Werneburg, and L. Wilson for valuable discussions; L. Rager and S. Suzuki for photographs and helpful information; C. Zollikofer and N. Morimoto for assistance with CT scans; M. Ando, K. Catania, M. Iwasa, J. Kimura, S. Ohdachi, M. Sasaki, and A. Shinohara for providing their precious specimens; and reviewers for useful suggestions. We also thank the curators of the Botanical Garden Museum of Hokkaido University, Japan Monkey Center, Kyoto University Museum, and Natural History Museum of Basel, National Museum of Nature and Science, Tokyo for access to specimens. This study was supported by the Japan Society for Promotion of Science (23-4832) to D.K. and by the Swiss National Fund (3100A0-133032/1) to M.R.S.-V.

References

1. Gross JB & Hanken J (2008) Review of fate-mapping studies of osteogenic cranial neural crest in vertebrates. *Dev Biol* 317:389-400.
2. Broom R (1916) On the structure of the skull in *Chrysochloris*. *Proc Zool Soc Lond* 86:449-458.
3. Morriss-Kay GM (2001) Derivation of the mammalian skull vault. *J Anat* 199:143-151.
4. Augier M (1931) Quelques problèmes relatifs au développement du crâne. Voûte occipitale chez l'Homme et d'autres Mammifères. *Arch Anat Histol Embryol* 13:34-66. French
5. Broili F (1917) Unpaare Elemente im Schädel von Tetrapoden. *Anat Anz* 49:561-576. German
6. von Huene F (1912) Die Herkunft des Os interparietale der Mammalia. *Anat Anz* 42:522-524. German
7. Wegner RN (1960) Das Os interparietale als integrierender Bestandteil der Säuger- und Theromorphenschädel. *Gegenbaurs Morphol Jahrb* 100:375-419. German
8. Watson DMS (1913) IV.—On some features of the structure of the therocephalian skull. *J Nat Hist* 11:65-79.
9. Goodrich ES (1931) *Studies on the Structure and Development of Vertebrates* (Dover Publications, New York).
10. Giebel C (1874) Säugetiere: Mammalia. *Dr HG Bronn's Klassen und Ordnungen des Thierreichs, wissenschaftlich dargestellt in Wort und Bild* 6:1-36. German

- 1 11. Novacek MJ (1993) in *The Skull: Patterns of Structural and Systematic Diversity*, eds
2 Hanken J & Hall BK (University of Chicago Press, Chicago), Vol 2, pp 438-545.
- 3 12. Keith A (1948) *Human Embryology and Morphology* (Edward Arnold and Co., London).
- 4 13. Jiang X, Iseki S, Maxson RE, Sucov HM, & Morriss-Kay GM (2002) Tissue origins and
5 interactions in the mammalian skull vault. *Dev Biol* 241:106-116.
- 6 14. Roybal PG, *et al.* (2010) Inactivation of *Msx1* and *Msx2* in neural crest reveals an
7 unexpected role in suppressing heterotopic bone formation in the head. *Dev Biol* 343:28-
8 39.
- 9 15. Yoshida T, Vivatbutsiri P, Morriss-Kay G, Saga Y, & Iseki S (2008) Cell lineage in
10 mammalian craniofacial mesenchyme. *Mech Develop* 125:797-808.
- 11 16. Williston SW (1925) *Osteology of the Reptiles* (Harvard University Press, Cambridge).
- 12 17. Ranke J (1913) Über das Interparietale und die Verknöcherung des Schädeldaches bei
13 Affen. *Sitzungsberichte der Bayerischen Akademie der Wissenschaften*:223-269. German
- 14 18. Jollie M (1973) *Chordate Morphology* (Reinhold, New York).
- 15 19. Kingsley JS (1925) *The Vertebrate Skeleton from the Developmental Standpoint* (P.
16 Blakiston's Son & Co., Philadelphia).
- 17 20. de Beer GR (1937) *The Development of the Vertebrate Skull* (Oxford University Press,
18 London).
- 19 21. Kielan-Jaworowska Z, Cifelli R, & Luo ZX (2004) *Mammals from the Age of Dinosaurs:*
20 *Origins, Evolution, and Structure* (Columbia University Press, New York).
- 21 22. van Bemmelen JF (1901) *Der Schädelbau der Monotremen* (G. Fischer, Jena). German

- 1 23. Landry SO (1964) The form and phylogenetic history of the mammalian interparietal
2 bone. *Am Zool* 4:400-400.
- 3 24. Kuhn HJ & Zeller U (1987) in *Mammalia Depicta: Morphogenesis of the Mammalian*
4 *Skull*, eds Kuhn HJ & Zeller U (Verlag Paul Parey, Hamburg), pp 51-70.
- 5 25. Weber M & Abel O (1928) *Die Säugetiere. Einführung in die Anatomie und Systematik*
6 *der Recenten und Fossilen Mammalia*. (Gustav Fischer Verlag, Jena). German
- 7 26. Beer JR (1965) The interparietal in kangaroo rats. *Southwest Nat* 10:145-150.
- 8 27. Matsumura G, Uchiumi T, Kida K, Ichikawa R, & Kodama G (1993) Developmental
9 studies on the interparietal part of the human occipital squama. *J Anat* 182:197-204.
- 10 28. Eales NB (1950) The skull of the foetal narwhal, *Monodon monoceros* L. *Philos Trans R*
11 *Soc Lond, B, Biol Sci* 235:1-33.
- 12 29. Starck D (1995) *Lehrbuch der Speziellen Zoologie. Wirbeltiere. Teil 5, Säugetiere*
13 (Gustav Fischer-Verlag, Jena). German
- 14 30. Gregory WK (1910) The orders of mammals. *Bull Am Mus Nat Hist* 27:1-524.
- 15 31. Kuratani S (1994) [Mysteries of occipital bones II. - Mechanism of the segmental
16 patterning]. *The Bone* 8:137-150. Japanese.
- 17 32. Wible JR (1991) Origin of Mammalia: the craniodental evidence reexamined. *J Vert*
18 *Paleontol* 11:1-28.
- 19 33. Hershkovitz P (1977) *Living New World Monkeys (Platyrrhini): with an Introduction to*
20 *Primates* (University of Chicago Press, Chicago).

- 1 34. Maggi L (1898) Placche osteodermiche interparietali degli Stegocephali e rispondenti
2 centri di ossificazione interparietali dell' uomo. *Rendic R Ist Lomb di Sc e Lett* 2:1-38.
3 Italian
- 4 35. Broom R (1914) Some points in the structure of the diadectid skull. *Bull Am Mus Nat Hist*
5 33:109-114.
- 6 36. Säve-Söderbergh G (1935) On the dermal bones of the head in labyrinthodont
7 stegocephalians and primitive reptilia: with special reference to Eotriassic stegocephalians
8 from east Greenland. *Meddel Grønland* 98:1-121.
- 9 37. Romer AS (1956) *Osteology of the Reptiles* (University of Chicago Press, Chicago).
- 10 38. Mickoleit G (2004) *Phylogenetische Systematik der Wirbeltiere* (Pfeil Verlag, München).
11 German
- 12 39. Rocek Z (1988) Origin and evolution of the frontoparietal complex in anurans. *Amphibia-*
13 *Reptilia* 9:385-403.
- 14 40. Smirnov SV (1997) Additional dermal ossifications in the anuran skull: morphological
15 novelties or archaic elements? *Russ J Herpetol* 4:17-27.
- 16 41. Parrington F (1937) V.—A note on the supratemporal and tabular bones in reptiles. *J Nat*
17 *Hist* 20:69-76.
- 18 42. Klembara J (2001) Postparietal and prehatching ontogeny of the supraoccipital in
19 *Alligator mississippiensis* (Archosauria, Crocodylia). *J Morphol* 249:147-153.
- 20 43. Gaupp E (1900) Das chondrocranium von *Lacerta agilis*. *Anat Embryol* 15:434-594.
21 German

- 1 44. Conrad JL (2008) Phylogeny and systematics of Squamata (Reptilia) based on
2 morphology. *Bull Am Mus Nat Hist* 310:1-182.
- 3 45. Müller J, *et al.* (2011) Eocene lizard from Germany reveals amphisbaenian origins.
4 *Nature* 473:364-367.
- 5 46. Zusi RL (1993) in *The Skull: Patterns of Structural and Systematic Diversity*, eds Hanken
6 J & Hall BK (University of Chicago Press, Chicago), Vol 2, pp 391-437.
- 7 47. Lucas SG & Luo Z (1993) *Adelobasileus* from the Upper Triassic of West Texas: the
8 oldest mammal. *J Vert Paleontol* 13:309-334.
- 9 48. Kermack K, Mussett F, & Rigney H (1981) The skull of *Morganucodon*. *Zool J Linn Soc*
10 71:1-158.
- 11 49. Wible JR (2008) On the cranial osteology of the *Hispaniolan solenodon*, *Solenodon*
12 *paradoxus* Brandt, 1833 (Mammalia, Lipotyphla, Solenodontidae). *Ann Carnegie Mus*
13 77:321-402.
- 14 50. Wible JR (1990) Petrosals of Late Cretaceous marsupials from North America, and a
15 cladistic analysis of the petrosal in therian mammals. *J Vert Paleontol* 10:183-205.
- 16 51. Abdala F, Flores DA, & Giannini NP (2001) Postweaning ontogeny of the skull of
17 *Didelphis albiventris*. *J Mammal* 82:190-200.
- 18 52. Brink AS (1950) XCI.—On a new species of *Cistecephalus* Owen. *J Nat Hist* 3:985-997.
- 19 53. Broom R (1932) *The Mammal-like Reptiles of South Africa and the Origin of Mammals*
20 (HF & G. Witherby, London).

- 1 54. King GM (1988) in *Encyclopedia of Paleoherpetology*, ed Wellnhofer P (Gustav Fischer,
2 Stuttgart), pp 1-174.
- 3 55. Sidor CA (2001) Simplification as a trend in synapsid cranial evolution. *Evolution*
4 55:1419-1442.
- 5 56. von Huene FR (1913) The skull elements of the Permian Tetrapoda in the American
6 Museum of Natural History, New York. *Bull Am Mus Nat Hist* 32:315-386.
- 7 57. Jarvik E (1980) *Basic Structure and Evolution of Vertebrates* (Academic Press, New
8 York).
- 9 58. Couly GF, Coltey PM, & Le Douarin NM (1993) The triple origin of skull in higher
10 vertebrates: a study in quail-chick chimeras. *Development* 117:409-429.
- 11 59. Noden DM & Schneider RA (2006) in *Neural Crest Induction and Differentiation -*
12 *Advances in Experimental Medicine and Biology*, ed Saint-Jeannet J (Landes Bioscience,
13 Georgetown), pp 1-23.
- 14 60. Rawlins J & Opperman L (2008) in *Craniofacial Sutures Development, Disease and*
15 *Treatment*, ed Rice DP (Karger, Basel), pp 178-196.
- 16 61. Loeys BL, *et al.* (2005) A syndrome of altered cardiovascular, craniofacial,
17 neurocognitive and skeletal development caused by mutations in *Tgfb β 1* and *Tgfb β 2*. *Nat*
18 *Gen* 37:275-281.
- 19 62. Spagnoli A, *et al.* (2007) TGF- β signaling is essential for joint morphogenesis. *J Cell Biol*
20 177:1105-1117.

- 1 63. Seo HS & Serra R (2009) *Tgfr2* is required for development of the skull vault. *Dev Biol*
2 334:481-490.
- 3 64. Yi JJ, Barnes AP, Hand R, Polleux F, & Ehlers MD (2010) TGF- β signaling specifies
4 axons during brain development. *Cell* 142:144-157.
- 5 65. Smith KM, Williamson TL, Schwartz ML, & Vaccarino FM (in press) Impaired motor
6 coordination and disrupted cerebellar architecture in *Fgfr1* and *Fgfr2* double knockout
7 mice. *Brain Res*.
- 8 66. Stevens HE, *et al.* (2010) *Fgfr2* is required for the development of the medial prefrontal
9 cortex and its connections with limbic circuits. *J Neurosci* 30:5590-5602.
- 10 67. Richtsmeier JT, *et al.* (2006) Phenotypic integration of neurocranium and brain. *J Exp*
11 *Zool B Mol Dev Evol* 306:360-378.
- 12 68. Lieberman D (2011) *The Evolution of the Human Head* (Belknap Press, Cambridge).
- 13 69. Koyabu D, *et al.* (2011) Heterochrony and developmental modularity of cranial
14 osteogenesis in lipotyphlan mammals. *EvoDevo* 2:21.
- 15 70. Bärmann EV & Sánchez-Villagra MR (2012) A phylogenetic study of late growth events
16 in a mammalian evolutionary radiation —the cranial sutures of terrestrial cetartiodactyl
17 mammals. *J Mammal Evol* 19:43-56.
- 18
- 19

Figure legends

Fig. 1. Dorsal view of cleared and stained embryonic specimens. (A) Specimen of southern long-nosed armadillo (*Dasypus hybridus*). Note the dermal interparietal growing from the anterior tip of the cartilaginous supraoccipital. (B) Specimen of Asian grey shrew (*Crocidura attenuata*). Note the typical reticular appearance of the interparietal seamlessly attached to the anterior tip of the supraoccipital. IP, interparietal; SO, supraoccipital; PA, parietal; FR, frontal.


Fig. 2. Microtomographic reconstructions of the embryonic skulls in various species (caudal view). (A). Amur hedgehog (*Erinaceus amurensis*). (B) Cat (*Felis cattus*). (C) Japanese deer (*Cervus nippon*). (D) Horseshoe bat (*Rhinolopus cornutus*). (E) Hardwicke's woolly bat (*Kerivoula hardwickii*). (F) Blanford's bat (*Hesperoptenus blanfordi*). (G) Bottlenose dolphin (*Tursiops truncatus*). (H) Cow (*Bos primigenius*). (I) Large Japanese field mouse (*Apodemus speciosus*). (J) Japanese macaque (*Macaca fuscata*). IP, interparietal; SO, supraoccipital; BO, basioccipital; EO, exoccipital; SQ, squamosal; PE, petrosal; PA, parietal.

Fig. 3. Lateral view of skulls of **opossum and rodents**. (A) **dorsal view of a white-eared opossum (*Didelphis albiventris*)**. (B) and (C) Degu (*Octodon degu*) skulls of different stages. (D) Perinatal skull of Patagonian cavy (*Dolichotis patagonum*). Note the lateral interparietal element (LI). (E) A close-up image of (D). IP, interparietal; LI, lateral interparietal element ; SO, supraoccipital; SQ, squamosal; PE, petrosal; PA, parietal.


Fig. 4. Dorsal view of embryonic skulls of degu (*O. degu*). (A) Note the lateral interparietal elements (highlighted with arrows) lying lateral to the median interparietal (IP). (B) A more advanced individual of degu. Note the lateral interparietal elements (highlighted with arrows). IP, interparietal; SO, supraoccipital; PA, parietal.


Fig. 5. Evolutionary and development pattern of the interparietal. (A) Cladogram and caudal view of skulls of Early Permian lissamphibian *Diadectes*, Late Permian therapsid *Titanophoneus*, and human fetus. The medial interparietal (postparietal) are given in blue, and the lateral interparietal (tabular) is shown in pink. (B) Diversity of the interparietal elements. Single, paired, or tripartite interparietal is consequently produced, depending on the fusion pattern of the four elements. (C) Suggested basic structure of the interparietal. The interparietal is suggested to be consisted of two medial elements and two lateral elements. (D) Tissue origins of the mouse skull vault [after Jiang et al. (13)]. The interparietal has dual origins, the median portion derived from the neural crest and the lateral part from the mesoderm. NA, nasal; FR, frontal; PA, parietal; IP, interparietal; SO, supraoccipital; SQ, squamosal.






A









B











SUPPLEMENTARY INFORMATION

Paleontological and developmental evidence resolve the homology and dual embryonic origin of a mammalian skull bone, the interparietal

Daisuke Koyabu ^{a,b,1}, Wolfgang Maier^c, Marcelo R. Sánchez-Villagra^{a,1}

^aPaleontological Institute and Museum, University of Zürich, Karl Schmid-Strasse 4, 8006 Zürich, Switzerland

^bKyoto University Museum, Yoshida Honmachi, 606-8501 Kyoto, Japan

^cUniversity of Tübingen,, Auf der Morgenstelle 28, 72076 Tübingen, Germany

*corresponding authors: koyabu@pim.uzh.ch; m.sanchez@pim.uzh.ch

Table of Contents

Part A. Supplementary descriptions on the interparietal of studies species.

Part B. Supplementary Figures

Figure S1. Dorsal view of the skulls of stem amniotes and sauropsids.

Figure S2. Occipital region of stem amniotes and sauropsids.

Figure S3. Occipital region of non-mammalian synapsids.

Figure S4. Occipital region of opossums.

Figure S5. Dorsal view of the skulls of selected mammals.

Figure S6. Mammalian embryonic specimens described by Parker.

Figure S7. Mammalian embryonic specimens described by Parker (continued).

Figure S8. Caudal view of skulls.

Figure S9. CT reconstructions of the embryonic skulls (lateral view).

Figure S10. Double-stained embryonic specimens of moles, hedgehogs, and shrews.

Figure S11. An embryonic skull of *Sorex araneus* (common shrew) described by de Beer.

Figure S12. Caudal view of the occipital region in a double-stained embryonic specimen of *Sorex unguiculatus* (long-clawed shrew).

Figure S13. Interparietal development in *Homo sapiens*.

Figure S14. Interparietal and preinterparietal in *Procavia capensis* (rock hyrax).

Part C. Supplementary Table

Table S1. Observed pattern of the interparietal.

Part D. Additional references of supplementary information

Part A. Supplementary descriptions on the interparietal of studies species

Fish, stem tetrapods and, lissamphibians

Sarcopterygians such as *Youngolepis praecursor* and actinopterygians including *Moythomasia nitida*, *Kansasiella eatoni* are documented to possess paired postparietal (1, 2). Already in *Ichthyostega* the postparietals are medially fused - but not in other early tetrapods (3, 4). *Acanthostega* had both postparietal and tabular (5). Many fossil batrachomorpha (sensu 6) have retained paired postparietals, including *Eryops*, *Diplocaulus*, *Phlegethontia* (*Dolichosoma*), *Branchiosaurus* (7), and *Apateon dracyiformis* (8). *Palaeogyrinus* (*Palaeoherpeton*) is documented to possess a paired postparietal by Romer and Parsons (1). Although Jarvik (3) identified a small unpaired dermal ossification at the dorsal side of the foramen magnum of *Rana*, extant lissamphibians are commonly recorded as missing postparietals (9). On the other hand, *Eocaecilia micropodia*, which is considered to represent the stem condition of lissamphibian (10), and a member of diadectomorph *Limnoscelis* exhibit a paired postparietal (11). *Anthracosaurus* shows paired postparietal (12). At least some dissorophids (*Dolesempetron*), a putative stem group of all or some lissamphibians possess paired postparietals (13, 14). Although most modern anurans are supposed to have lost the postparietal, it is present in some living groups such as *Pelobates* (15) and *Bombina* (16).

Stem-amniotes and sauropsids

Basal amniotes such as, *Paleothyris* (17), *Seymouria* (Figs.1A and 2B) (18), and *Kotlassia* (11) exhibit paired postparietals. Broom (19) documented the unpaired postparietal in *Diadectes* and *Chilonyx* and noted that, from the direction of the fibres of this medial element, it seems probable that it originated from two ossification centers.

On the other hand, no extant sauropsid seems to possess postparietal, except for *Alligator mississippiensis* (20). Here, the bone is a small dermal plate lying ventrally and posteriorly to the posterior margin of the parietal and dorsally to the trabecular bone, forming a dorsal surface of the supraoccipital portion of the neural endocranium. Chelonians (the clade uniting turtles, tortoises and terrapins) seems to lack the postparietal; there is no postparietal found in *Proganochelys*, *Caretta*, *Lepidochelys*, *Chelonia* and *Emys* (11, 21, 22). No postparietal is found in either the embryonic *Pelodiscus sinensis* (23) or *Emydura subglobosa* (24). Several works on squamates confirm the lack of the postparietal in this group: apparently *Lacerta* (25), *Varanus* (11), *Tupinambis* (21), and *Ctenosaura* (26), as well as the comprehensive study of Conrad (27) and the recent description of an Eocene “lizard” from

Messel in Germany (28). *Sphenodon* has no postparietal (11, 21), and this is confirmed in fetal stages as well (29). No placodont and in fact no other sauropterygian possess a postparietal bone (30). On the other hand, among fossil diapsids, the presence of paired postparietal is documented in *Araeoscelis* by Parrington (18), and in *Protorothyris* (Fig. S1B), *Euparkeria* (Fig. S1C), *Milleratops* (or juvenile? of *Milleretta*) (Fig. S1D), *Milleretta* (Fig. S1E) *Machaeroprotopus* (Fig. S2C), *Youngina* (Fig. S2D) by Romer (11). The anapsid reptile *Labidosaurus* (Fig. S2B) shows an unpaired postparietal (18).

Extinct and extant birds apparently lack the postparietal (31, 32). In his careful study Müller (33) reported that *Rhea* and other primitive birds have no postparietal. Jollie (21) who depicts the cranium of a freshly hatched chick shows no sign of an postparietal. There is no postparietal in *Protoavis* (34) nor in *Archaeopteryx* (35). Raath (36) who showed a detailed drawing of the occiput records no postparietal for *Syntarsus* (*Megapnosaurus*).

Non-mammalian synapsids: “pelycosaurids” and “therapsids”

Among pelycosaurs, a paraphyletic grouping of basal synapsids, *Ophiacodon uniformis* (37), *Edaphosaurus pogonias* (37-39) and for *Theropleura retroversa* (Fig. 3A) (40) reportedly has a single bone labeled as postparietal which lies just dorsal to the supraoccipital, contacting the tabulars laterally and the parietal dorsally. Parrington (18) documented the presence of the paired postparietal and paired tabular in *Dimetrodon*, *Diops*, and *Theropleura*.

Thrinaxodon liorhinus (Fig. 3B) is reported to possess an enlarged unpaired postparietal and paired tabular (41). Various therapsids such as *Diarthrognathus* (42), *Myosaurus gracilis* (43), *Rechnisaurus cristarhynchus* (44), *Ennatosaurus tecton* (45), *Ecteninion lunensis* (46), *Emydops* (47), *Kombuisia frerensis* (48), *Scylacops*, *Cynognathus*, *Ulemosaurus*, *Synostocephalus*, *Lycedops* (Fig. 3C), *Bauria* (11), *Hofmeyria* (Fig. 3D), *Hyenosaurus* (Fig. 3E) (49), *Procynosuchus* (50), *Titanophoneus*, *Aneugomphius* (51), *Lystrosaurus amphibius* (52) are reported to possess the postparietal. The monographic work of Broom on non-mammalian synapsids (53) provides the illustrations of more than 40 species, depicting the postparietal in which postparietals are labeled.

Docodonts

Haldanodon exspectatus was reported to possess a triangular median interparietal (54).

Monotremes

According to de Beer (55), Kuhn (56), Zeller (57), and Jollie (21), monotremes do not have the interparietal and rather possess a very expanded supraoccipital. On the other hand, van Bemmelen (58) reported that *Ornithorhynchus* possesses an interparietal. Here, he labeled the medial interparietal as a “medial parietal” and the lateral interparietal as a “lateral parietal”. Landry (59) commented that the interparietal appears in the embryonic *Zaglossus* (long-beaked echidna) as a large element covering the back end of the brain case. Kierdorf and Kierdorf (60) reported a small medial membrane bone of only transient occurrence observed in *Tachyglossus aculeatus* (61), which is possibly a remnant of the interparietal element in this species.

Marsupials

From our embryonic specimens, we confirmed the presence of the interparietal in *Trichosurus*, *Monodelphis*, *Macropus*, and *Sminthopsis*. Owen (62) reported that *Macropus*, *Hypsiprymnus*, *Petaurus*, *Phalanger*, *Myrmecobius*, and *Phascolomys* exhibit the interparietal. An unpaired interparietal was described in postnatal *Didelphis marsupialis* (63), and in *Monodelphis* by Clark and Smith (64). Wible (65) noted that this bone apparently fuses seamlessly with the supraoccipital along the nuchal crest. In our ontogenetic series, the interparietal was found to be unpaired, paired, tripartite, and quadripartite (Figs. S4A-D). *Myrmecobius fasciatus* (Dasyuromorphia) is reported to have an enlarged interparietal (66). As pointed out by Wible (65), since the bone seamlessly fuses with the supraoccipital, the interparietal is often erroneously labeled in adults as part of the supraoccipital bone. Abdala (67) reported the fusion of the unpaired interparietal to the supraoccipital in *Didelphis albiventris*. The interparietal of *D. albiventris* was found to be quadripartite (Figs. S4E and F). Forasiepi (68) documented the presence of the unpaired interparietal in the fossil metatherian *Arctodictis sinclairi*. Voss and Jansa (69) discussed the presence of the interparietal in some didelphids and reported this bone in stem metatherians. According to these authors, the presence of a “large undivided interparietal bone that is wedged between the parietals anteriorly and fused to the supraoccipital posteriorly” may be a didelphid synapomorphy. The search for morphological synapomorphies for didelphids makes this question worthwhile, but the hypothesis seems questionable considering the difficulty of assessing the condition in basal crown-group Marsupialia such as *Herpetotherium* (70). The different interpretations on the condition of *Pucadelphys* (71) and *Mayulestes* (72), which have been described or illustrated as possessing different conditions, may be best explained by the fact that different

stages of cranial vault sutural fusion may have been considered. Wegner (73) reported the presence of the interparietal in *Bettongia*, *Petaurus*, *Dedrolagus*, *Phalanger*, *Wombatus*, *Phascolarctos*, *Trichosurus*, *Onychogale*, and *Sarcophilus* and commented that the interparietal is a persistent bone in marsupials.

Notoungulates

Toxodon burmeisteri, *Typotherium cristatum*, *Nesodon imbricatus*, and *Pachyrucus* exhibit the median interparietal and lateral interparietals (74).

Afrotheria

Giebel (75) and Lawlor (76) documented the presence of this bone across Hyracoidea. Our own investigations confirmed its presence in *Heterohyrax brucei* and *Procavia capensis* (Fig. S5B). In addition to this, we identified unpaired, paired, and tripartite (one triangular large medial and paired lateral elements) interparietals in *Dendrohyrax abroreus*. Similarly, unpaired, paired, tripartite (one triangular large medial and paired lateral elements), and maximally quadruple (medially paired and laterally paired) interparietals were found in *P. capensis*. Wegner (73) reported that *P. capensis* shows a tripartite interparietal consisted of one large medial and two lateral elements. Parker (77, 78) documented the interparietal in *Microgale longicaudata* (Figs. S6A and B), *Tenrec ecaudatus* (Figs. S6G and H), and *Hemicentetes madagascarensis* (Figs. S6C and D). In *M. longicaudata*, he also noted the presence of the “supratemporals”, which shall be rather called tabular, lying lateral to the interparietal. In tenrecs (*Potamogale velox*, *Hemicentetes semispinosus*, and *Tenrec ecaudatus*), Schunke and Zeller (79) observed that the interparietal bone is unpaired and separated from the supraoccipital in embryos. In *Hemicentetes semispinosus*, the interparietal fuses first with the supraoccipital. The interparietal is formed considerably later in *Potamogale* compared to *Hemicentetes* and remains separated from the supraoccipital for some time (79). Leimgruber (80) also reported a notably developed interparietal bone in *Tenrec*. In *Setifer* the interparietal bone fuses shortly after its origin with the supraoccipital bone (81). Similarly, one median and two lateral interparietals were described for *Chrysochloris asiatica* (82). The African elephant is reported to possess a small interparietal which is present at birth and fuses with other bones with growth (83). Giebel (75) reported that paired interparietals are found in young individuals of the elephant. De Beer (55) noted that the interparietals fuse with the parietals in sirenians. The golden mole *Eremitalpa* is reported to possess an unpaired interparietal bone (81). According to our investigations, a

neonate *Macroscelides proboscideus* shows a large interparietal (Fig. S5A), and a neonate of *Orycteropus afer* shows a very large interparietal. Parker (77, 78) documented the interparietal in *Rhynchocyon cirnei* (Figs. S7E and F) and greatly expanded interparietal for *O. afer* (Figs. S5G and H). We found mutli-elemented interparietal bones in *Dugong dugon*, *Trichechus manatus*, and *T. senegalensis*. The interparietal is reported also in *Rytina* by Wegner (73). He also noted the presence of the tabular, an extra bone lateral to the interparietal, in *T. manatus*.

Xenarthra

Kingsley (84) noted that no xenarthrans possesses the interparietal. On the contrary, our investigation found a paired interparietal in *Cyclopes didactylus*, an unpaired interparietal in *Tamandua tetradactyla* (Fig. S5D), and an unpaired interparietal in a prenatal *Bradypus tridactylus*. Similarly, Wegner (73) reported its presence in *Bradypus tridactylus* and *C. didactylus*. Our own examination on perinatal *Bradypus tridactylus* (Fig. S5C) also confirmed the presence of the interparietal. De Beer (55) noted that *Dasypus* lacks the interparietal, but our own investigation found a dermal interparietal bone in *Dasypus hybridus* (Fig. 1A).

Euarchontoglires

The interparietal bone is commonly found in rodents and lagomorphs. This bone in Glires is relatively enlarged (85). Beer (86) conducted a survey on the presence of the interparietal in 21 *Dipodomys* species, sampling 2360 prenatal and postnatal specimens. In *Dipodomys*, the interparietal was basically formed from four ossification centers and finally united into an unpaired shield-shaped structure. In the monographic work by Tullberg (87), the interparietal can be found in figures for more than 40 genera. We found unpaired interparietals in *Pedetes capensis*, *Galea musteloides*, *Hydromys chrysogaster*, *Jaculus jaculus*, *Micromys minutus*, and *Apodemus speciosus* (Figs. 2I and S9D). In addition, three interparietal bones, one medial and two lateral, were found in various stages of *Octodon degu* (Figs. 4A-C and 5). Similarly, median interparietals and two lateral interparietals were found in two juveniles from different localities of *Dolichotis patagonum* (Figs. 3D and E). In *Lepus*, the interparietal remains fuses with the parietal in mature individuals. Our neonatal *Oryctolagus cuniculus* shows an unpaired interparietal. Starck (22) noted that, although the interparietal generally remains independent from other bones in the Lagomorpha and Rodentia, the interparietal often fuses with the parietal rather than with the supraoccipital in rodents. Our CT investigation on the newborn of *D. patagonum* confirmed such trend of

fusion with the parietal. Similarly, the interparietal fuses with the parietal in *Pteromys volans* and *Spermophilus undulatus*. We confirmed that the interparietal fuses with the parietal in *Petaurista leucogenys* and *Pteromys momonga*.

Parker (77) documented the presence of the interparietals in *Cynocephalus volans* and *Galeopterus variegatus* (Figs. S6E and F). The presence of this bone in *Galeopterus* was documented also by Giebel (75). Wortmann (88) and Zeller (89) documented the presence of the interparietal in *Tupaia belangeri*. This investigation was confirmed in our investigation on young newborn *Tupaia belangeri*.

Among primates, young individuals of *Daubentonia madagascariensis* and *Cebus* are reported to possess the interparietals (75). There are many studies reporting the existence of the interparietal in humans (e.g., 90, 91) (Fig. S13). The presence of the interparietal is reported in the newborn *Hylobates syndactylus*, and *Pan troglodytes* (75). Kingsley (84) and Eaglen (92) noted that lemurids frequently show unpaired interparietals. Wegner (73) described the enlarged interparietal in *Loris tardigradus* and *D. madagascariensis*. Our own investigation confirmed the presence of the interparietal in the fetal skull of *Pan troglodytes* (Fig. S8C) and in young *Galago senegalensis* (Fig. S5I). In addition, our microCT examination on fetal specimens clearly showed the presence of the membranous interparietal in *Macaca fascicularis* (Fig. S9E) and in *M. fuscata* (Figs. 2J and S9F). In macaques, the interparietal develops after the supraoccipital and then rapidly fuses with this bone. Starck (93) reported that he failed to find the interparietal in a late fetal stage of *Propithecus*, although he noted a clear membranous “bone growth” at the dorsal margin of the supraoccipital, presumably an interparietal, was visible. In new world primates, Hershkovitz (94) documented the frequent presence of the interparietal across Cebidae, and Chopra (95) reported its presence in *Saimiri*.

Laurasiatherians

The presence of paired interparietal was documented in *Solenodon paradoxus* (96). In juvenile *Solenodon* there is a paired interparietal, the two parts of which become fused between them and with their neighboring bones (96). From our original investigation on embryonic specimens, we confirmed that *Erinaceus europaeus* (Figs. S7A and B) and *E. amurensis* (Fig. 2A) possess paired interparietals. The presence of the interparietal is also documented in *Echinosorex* (75). There is little published information on the presence of the interparietal in moles, possibly due to their very early closure of cranial sutures. Although Goswami and Prochel (97) noted that there is no evidence of presence of the interparietal in *T.*

europaea, our investigation in embryonic specimens found thin paired membranous interparietals which initiate their ossification from the dorsal tip of the supraoccipital and then grow gradually toward the rostral direction in both *T. occidentalis* (Fig. S10A) and *T. europaea*. The suture between the interparietal and the supraoccipital closes rapidly. We also found the developing dermal interparietal at the anterior tip of the supraoccipital in an embryonic specimen of *Scapanus orarius* (Figs. S10C and D) and *Mogera wogura*. De Beer (98) documented the interparietal in the *Sorex araneus* (Figs. S11A and B). He described that the supraoccipital is partly covered by the dermal interparietal. On the other hand, Augier (99) noted that soricids have no interparietal, but the supraoccipital forms a dermal second center of ossification in the expanded tectum posterius. Kuratani (100) observed the expansion of the dermal interparietal bone and the extremely unique case of the lack of the cartilaginous supraoccipital in *Suncus murinus*. We found dermal interparietal bones in embryonic specimens of *Sorex unguiculatus* and *Crocidura attenuata*. In *Sorex unguiculatus*, the interparietal seemed to dorsally cover the supraoccipital, forming a double layered structure with the supraoccipital (Fig. S12). In *C. attenuata*, the cartilaginous supraoccipital is small but evident, and the dermal interparietal grows toward the anterior direction (Figs. 1B and S10G-H). On the other hand, in the specimens of *C. watasei* and *Suncus murinus* we studied, the dermal interparietal bone was apparently present but the cartilaginous supraoccipital was not clearly identified at least in our ontogenetic series. Recently, presence of the interparietal was commonly found in myosoricine shrews (*Surdisorex*, *Congosorex*, and *Myosorex*) (101). Parker (78) described the interparietal in *Sorex vulgaris* (*araneus*) (Figs. S7C and D).

The reconstruction of CT images of embryonic whale specimens showed that *Tursiops truncatus* possesses an expanded interparietal (Fig. 2G). We found the interparietal also in *Phocoena phocoena* the harbour porpoise. Eales (102) described the greatly expanded four-element interparietals in *Monodon monoceros*. The interparietal is also reported for other cetaceans (*Balaenoptera borealis*, *B. musculus*, *Delphinus delphinus*, and *Inia geoffrensis*) (73, 103-105). Recent investigation on embryonic *Stenella attenuata* documented the presence of the interparietal (106). The interparietal is large and square-shaped bone, stretching toward the medial plane and making up part of the dorsal roof of the braincase. However, we assume that the bones labeled as “right and left ossifications centers of the supraoccipital” (see Fig. 6C of ref. 106) are not part of the cartilaginous supraoccipital given their dermal appearance. We postulate that the “right and left ossifications centers of the supraoccipital” in Moran (106) are one of the ossification centers of the interparietal (see ref.

102), thus exhibiting three interparietals. According to Starck (107), the interparietal fuses with the supraoccipital in cetaceans.

In a late fetal cranium of *Capreolus capreolus* large and paired interparietals were visible (Fig. S5H). Similarly, we found the interparietal in a young *Tragelaphus scriptus*. Our CT examination on embryonic specimens demonstrated paired interparietals in *Cervus nippon* (Fig. 2C) and *Bos primigenius* (Fig. S9B). Quadrupartite interparietal was also found in *B. primigenius* (Fig. 2H). The interparietal and two lateral elements are depicted for a fetal *B. grunniens* by Starck (see Table 34 of ref. 108). Kierdorf and Kierdorf (60) reported the interparietal in four *Cervus* species (*C. elaphus*, *C. nippon*, *C. duvauceli* and *C. eldi*). Shapiro and Robinson (109) described an enlarged interparietal in *Odocoileus virginianus*. According to Rörig (110), the interparietal is present in *Capreolus capreolus* and in *Dama dama*. The interparietal found in *Alces alces* apparently fuse to the parietal. A pair of the interparietal is reported in *Muntiacus reevesi* (111). In many late fetal sheep *Ovis dalli* skulls, we found that the interparietal is almost always clearly distinguished. An infantile skull of *O. dalli* shows a paired interparietal. Paired interparietals are found in a prenatal *Vicugna pacos* (Fig. S8A). A two-day old calf of *Kobus ellipsiprymnus* shows clearly separated and paired interparietals. *Tragulus* is reported to possess the interparietal (73).

It is reported that the interparietal is highly common among ursids, including *Ailuropoda melanoleuca* (73). *Canis* is reported to have paired interparietals and also a median interparietal that immediately fuses with the supraoccipital (55). *Lutra lutra* reportedly possess an interparietal bone which is triangular in shape and has external sagittal crest (crista sagittalis externa) (112). *Mustela* and *Procyon lotor* (113), *P. pygmaeus* (114) are reported to have interparietals in the posterior end of the skull roof. *Vulpes vulpes* is reported to possess the interparietal (115). *Panthera leo*, *P. onca*, and *Puma concolor* are reported to show the interparietal by Wegner (116). He also noted that the interparietals are tripartite in *P. onca* and *P. concolor*. According to our microCT images of a late fetal *Felis domestica* (Figs. 2B and S9C), the interparietal is even larger than the supraoccipital. An unpaired interparietal was found in *Panthera pardus*. Although Giebel (75) noted that phocids lack the interparietal, our prenatal *Arctocephalus* (Fig. S5G) and *Callorhinus ursinus* skulls (Fig. S8B) apparently displayed enlarged dermal interparietals. *Phoca* is reported to exhibit the lateral interparietal element (82, 116). Here, we observed that supraoccipital is extremely reduced in size compared to the interparietal which dominates the caudal portion of the vault. In some neonate dog skulls, the supraoccipital sends a prominent dorsal process between the parietals; in *Speothos*, it is incompletely separated from the supraoccipital. Gregory (117) described that

Manis spp. lack the interparietal, but our examination found the interparital which fuses with the supraoccipital (Fig. S5E).

We confirmed the presence of the interparietal in our young individual of *Equus* (Fig. S5F). Giebel (75) reported that diamond shaped unpaired interparietal is often found in embryonic and young individuals. There are studies that report the existence of paired interparietal in *Tapirus* (tapir) (118-120). Kingsley (84) noted that the interparietal fuses with the parietals in *Equus* and *Tapirus*. Cuvier (121) reported the presence of the interparietal in *Rhinoceros* and *Diceros*.

We found paired interparietals in the embryonic specimens of *Hipposideros*, *Kerivoula* (Fig. 2E), *Hesperoptenus* (Fig. 2F), *Rhinolophus* (Fig. 2D and S9A). The interparietal fused with the supraoccipital and then with the parietal. Fawcett (122) showed a figure of *Miniopterus* that has large and paired interparietal. The interparietal is reported to be present in *Roussettus* (123) and *Artibeus* (124). As documented by Giannini et al. (125) among the Megachiroptera, in *Pteropus lylei* the interparietals are an unpaired bone clearly distinguishable in adults, as is also the case in *Dobsonia pannietensis*.

Part B. Supplementary Figures

Abbreviations: BO, basioccipital; EO, exoccipital; FR, frontal; IP, interparietal; MA, mastoid; MX, maxilla; NA, nasal; OO, opisthotic (paroccipital); PA, parietal; PE, petrosal; PO, postorbital; PP, postparietal; PTC, posttemporal canal; Q, quadrate; QJ, quadrojugal; SO, supraoccipital; SQ, squamosal; ST, supratemroal; TA, lateral interparietal element (tabular).




Figure S1. Dorsal views of the skulls of stem amniotes and sauropsids. (A) *Seymouria baylorensis* [redrawn from Parrington (18)], (B) *Protorothyris*, (C) *Euparkeria*, (D) *Milleretops* (or juvenile? of *Milleretta*), and (E) *Milleretta* [redrawn from Romer (11)].




Figure S2. Occipital region of stem amniotes and sauropsids. (A) *Seymouria baylorensis*, (B) *Labidosaurus*, (C) *Machaeroprotopus gregorii*, and (D) *Youngina capensis* [adopted from Parrington (18)]. The bones labeled as IP in the original reference are renamed as PP for consistencies.




Figure S3. Occipital region of non-mammalian synapsids. (A) *Theropleura* [redrawn from Parrington (18)], (B) *Thrinaxodon liorhinus* (redrawn from Broom (53)), (C) *Lycedops*, (D) *Hofmeyria*, (E) *Hyenosaurus* [redrawn from Broom (49)] , and (F) *Galesaurus planiceps* [adopted from Broom (53)]. The bones labeled as IP in the original references are renamed as PP for consistencies.




Fig. S4. Occipital region of opossums. (A) dorsal view and (B) lateral view of a 15 day *Monodelphis domestica* (gray short-tailed opossum) (PIMUZ-635D3/5), (C) dorsal view of a 15 day *M. domestica* (PIMUZ-635D2/5), (D) dorsal view of a 15 day *M. domestica* (PIMUZ-635D4/5), (E) dorsal view and (F) caudal view of a young *Didelphis albiventris* (white-eared opossum). The lateral interparietal elements (LI) are found laterally to the medial interparietals (IP) and are partly covered by the parietals in *M. domestica* (A-D). The lateral interparietal elements (LI) are found laterally to the medial interparietal elements (IP) in *D. albiventris* (E-F). The posttemporal canal (PTC) is found at the junction of the lateral interparietal (LI), squamosal (SQ), and mastoid (MA) (F).




Figure S5. Dorsal views of the skulls of selected mammals. (A) *Macroscelides proboscideus* (short-eared elephant-shrew), (B) *Proavia capensis* (rock hyrax), (C) *Bradypus tridactylus* (pale-throated sloth), (D) *Tamandua tetradactyla* (lesser anteater), (E) *Manis pentadactyla* (pangolin), (F) *Equus grevyi* (Grévy's zebra), (G) *Arctocephalus* sp. (fur seal), (H) *Capreolus capreolus* (European roe deer), (I) *Galago senegalensis* (Senegal bushbaby).




Figure S6. Mammalian embryonic specimens described by Parker (78). Note the interparietal bone labeled as “*i.p*” and the lateral interparietal element as “*s.t.*” (A) lateral and (B) dorsal view of *Microgale longicaudata* (lesser long-tailed shrew tenrec), (C) lateral and (D) dorsal view of *Hemicentetes madagascarensis* (lowland streaked tenrec), (E) lateral and (F) dorsal view of *Galeopterus variegatus* (Sunda flying lemur), (G) lateral and (H) dorsal view of *Tenrec ecaudatus* (tailless tenrec).




Figure S7. Mammalian embryonic specimens described by Parker (77, 78). Note the interparietal bone labeled as “i.p”. (A) lateral and (B) dorsal view of *Erinaceus europaeus* (European hedgehog), (C) lateral and (D) dorsal view of *Sorex araneus* (common shrew), (E) lateral and (F) dorsal view of *Rhynchocyton cirnei* (checkered elephant shrew), (G) lateral and (H) dorsal view of *Orycteropus afer* (aardvark).




Figure S8. Caudal view of the skulls. (A) *Vicugna pacos* (alpaca), (B) *Callorhinus urusinus* (northern fur seal), and (C) *Pan troglodytes* (chimpanzee).




Figure S9. CT reconstructions of the embryonic skulls (lateral view). (A) *Rhinolopus sp.* (horseshoe bat), (B) *Bos primigenius* (cow), (C) *Felis cattus* (cat), (D) *Apodemus speciosus* (large Japanese field mouse), (E) *Macaca fascicularis* (long-tailed macaque), and (F) *Macaca fuscata* (Japanese macaque).




Figure S10. Double-stained embryonic specimens of moles, hedgehogs, and shrews. (A) *Talpa occidentalis* (Iberian mole). Note the interparietal expanding anterior to the supraoccipital. (B) *Scapanus orarius* (coast mole). The interparietal is not yet ossified in this individual. (C) *S. orarius*. The interparietal has started to ossify from the anterior tip of the supraoccipital. (D) *S. orarius*. Close-up picture of (C). Note the small dermal outgrowth from the anterior tip of the supraoccipital. (E) *Mogera wogura* (Japanese mole). The interparietal has started to ossify. Note the dermal bone growing from the anterior tip of the supraoccipital. (F) *Erinaceus europaeus* (European hedgehog). Paired interparietal can be observed. (G) *Crocidura attenuata* (Asian grey shrew). Note the dermal bone (interparietal) attached to the anterior tip of the supraoccipital. (H) *C. attenuata*. Close-up picture of (G).




Figure S11. An embryonic skull of *Sorex araneus* (common shrew) described by de Beer (98). (A) dorsal and (B) ventral view. Note the interparietal (*ip*) dorsally covers the supraoccipital (*so*), forming a double layered structure.




Figure S12. Caudal view of the occipital region in a double-stained embryonic specimen of *Sorex unguiculatus* (long-clawed shrew). The interparietal (IP) is covering the supraoccipital (SO), forming two layers of bones. This covering pattern is comparable to that of *S. araneus* (Fig. S11).




Fig. S13. Interparietal development in *Homo sapiens*. (A) Occipital squama of a 4th month fetus. (B) Occipital squama in a 5th month fetal skull. Note that the interparietal develops from four ossification centers (IP and LI) [reprinted from Matsumura et al. (91) with permission, courtesy of John Wiley & Sons].




Fig. 14. Preinterparietal in *Procapra capensis* (rock hyrax). (A) the preinterparietal is not present, (B) the preinterparietal is present anterior to the interparietal (IP).

Part C. Supplementary Table

Table S1. Observed pattern of the interparietal (or postparietal).

		Observed element number	First fusion with
Fish, stem tetrapods, and lissamphibians	<i>Youngolepis praecursor</i>	4	none
	<i>Moythomsia nitida</i>	4	none
	<i>Kansasiella eatoni</i>	4	none
	<i>Ichthyostega</i>	1	none
	<i>Eryops</i>	4	none
	<i>Diplocaulus</i>	4	none
	<i>Phlegethontia (Dolichosoma)</i>	4	none
	<i>Branchiosaurus</i>	4	none
	<i>Apateon drachyiformis</i>	4	none
	<i>Palaeogyrinus (Palaeoherpeton)</i>	4	none
	<i>Eocaecilia micropodia</i>	4	none
	<i>Limnoscelis</i>	4	none
	<i>Doleserpeton</i>	4	none
	<i>Acanthostega</i>	4	none
	<i>Rana</i>	1	none
	<i>Pelobates</i>	4	none
	<i>Bombina</i>	4	none
	<i>Anthracosaurus</i>	4	none
Stem-amniotes and sauropsids	<i>Paleothyris</i>	4	none
	<i>Seymouria</i>	4	none
	<i>Kotlassia</i>	4	none
	<i>Diadectes</i>	4	none
	<i>Chilonyx</i>	4	none
	<i>Alligator mississippiensis</i>	2	none
	<i>Proganochelys</i>	none	-
	<i>Caretta</i>	none	-
	<i>Lepidochelys</i>	none	-
	<i>Chelonia</i>	none	-
	<i>Emys</i>	none	-
	<i>Pelodiscus sinensis</i>	none	-
	<i>Emydura subglobosa</i>	none	-
	<i>Lacerta</i>	none	-
	<i>Varanus</i>	none	-
	<i>Tupinambis</i>	none	-
	<i>Ctenosaura</i>	none	-
	<i>Sphenodon</i>	none	-
	<i>Placodus</i>	none	-
	<i>Araeoscelis</i>	4	none
	<i>Youngina</i>	4	none
	<i>Euparkeria</i>	1	none
	<i>Machaeroprotopus</i>	3	none
	<i>Milleretta</i>	4	none
	<i>Milleratops (or juvenile? of Milleretta)</i>	4	none
	<i>Protorothyris</i>	4	none
	<i>Labidosaurus</i>	4	none
	<i>Archaeopteryx</i>	none	-
	<i>Rhea</i>	none	-
	<i>Gallus gallus</i>	none	-
	<i>Protoavis</i>	none	-
	<i>Syntarsus</i>	none	-

The observed number of interparietal elements, the bone to which the interparietal fuses, and the presence of lateral interparietal elements (tabular) are documented. The observed number of elements includes the number of tabulars for non-mammalian vertebrates.

Table S1 (continued).

	Observed element number	First fusion with
Non-mammalian synapsids		
<i>Ophiacodon uniformis</i>	3	none
<i>Edaphosaurus pogonias</i>	3	none
<i>Theropseura retroversa</i>	3	none
<i>Dimetrodon</i>	4	none
<i>Varanosaurus</i>	4	none
<i>Diopus</i>	4	none
<i>Mesenosaurus</i>	4	none
<i>Thrinaxodon liorhinus</i>	3	none
<i>Diarthrogathus</i>	3	parietal
<i>Myosaurus gracilis</i>	3	none
<i>Rechnisaurus cristarhynchus</i>	1	none
<i>Ennatosaurus tecton</i>	4	none
<i>Ecteninion lunensis</i>	3	none
<i>Emydops</i>	3	none
<i>Kombuisia frerensis</i>	3	none
<i>Scylacops</i>	3	none
<i>Cynognathus</i>	3	none
<i>Ulemosaurus</i>	4?	none
<i>Synostocephalus</i>	3	none
<i>Lycedops</i>	3	none
<i>Bauria</i>	3	none
<i>Hofmeyria</i>	3	none
<i>Hyenosaurus</i>	3	none
<i>Procynosuchus</i>	3	none
<i>Titanophoneus</i>	3	none
<i>Galesuchus gracilis</i>	3	none
<i>Eoarcops vanderbyli</i>	3	none
<i>Scylacognathus parvus</i>	3	none
<i>Scylacops capensis</i>	3	none
<i>Gorgonops torvus</i>	3	none
<i>Lycaenops ornatus</i>	3	none
<i>Sycosaurus laticeps</i>	3	none
<i>Oudenodon bainii</i>	1	none
<i>Tropidostoma dubium</i>	1	none
<i>Dicynodon leptoscelus (Dicynodontia indet.)</i>	1	none
<i>Sintocephalus alticeps</i>	1	none
<i>Diictodon feliceps</i>	1	none
<i>Aulacocephalodon baini</i>	1	none
<i>A. laticeps</i>	1	none
<i>A. whaitisi</i>	1	none
<i>Platycephalus haughtoni</i>	1	none
<i>Emydops longiceps</i>	1	none
<i>E. trigoniceps</i>	1	none
<i>E. longus</i>	1	none
<i>Cistecephalus microrhinus</i>	1-3	none
<i>Lystrosaurus maccaigi</i>	1	none
<i>L. murrayi</i>	1	none
<i>L. curvatus</i>	1	none
<i>L. maccaigi</i>	3	none
<i>Galesaurus planiceps</i>	3	none
<i>Trirachodon kannemeyeri</i>	1	none

Table S1 (continued).

			Observed element number	First fusion with
Docodonts		<i>Haldanodon expectatus</i>	1	?
Monotremes		<i>Ornithorhynchus</i>	1	supraoccipital
		<i>Zaglossus</i>	1	supraoccipital
		<i>Tachyglossus aculeatus</i>	1	supraoccipital
Marsupials		<i>Trichosurus</i>	1-2	supraoccipital
		<i>Monodelphis</i>	1-4	supraoccipital
		<i>Macropus</i>	1-2	supraoccipital
		<i>Sminthopsis</i>	1-2	supraoccipital
		<i>Macropus</i>	1-2	supraoccipital
		<i>Hypsiprymnus</i>	1-2	supraoccipital
		<i>Petaurus</i>	1-2	supraoccipital
		<i>Phalanger</i>	1-2	supraoccipital
		<i>Myrmecobius</i>	1-2	supraoccipital
		<i>Phascolumys</i>	1-2	supraoccipital
		<i>Didelphis marsupialis</i>	1-2	supraoccipital
		<i>Myrmecobius fasciatus</i>	1-2	supraoccipital
		<i>Didelphis albiventris</i>	1-2	supraoccipital
		<i>Arctodictis sinclairei</i>	1-2	supraoccipital
		<i>Bettongia</i>	1-2	supraoccipital
		<i>Petaurus</i>	1-2	supraoccipital
		<i>Dedrolagus</i>	1-2	supraoccipital
		<i>Phalanger</i>	1-2	supraoccipital
		<i>Wombatus</i>	1-2	supraoccipital
		<i>Phascolarctos</i>	1-2	supraoccipital
		<i>Trichosurus</i>	1-2	supraoccipital
		<i>Onychogale</i>	1-2	supraoccipital
		<i>Sarcophilus</i>	1-2	supraoccipital
Notoungulates	Toxodontids	<i>Toxodon burmeisteri</i>	3	?
		<i>Nesodon imbricatus</i>	3	?
		<i>Pachyrucos</i>	3	?
	Mesotheriids	<i>Typotherium cristatum</i>	3	?
Afrotherians	Afrosoricids	<i>Tenrec ecaudatus</i>	1-2	supraoccipital
		<i>Hemicentetes madagascarensis</i>	1-2	supraoccipital
		<i>H. semispinosus</i>	1-2	supraoccipital
		<i>Microgale longicaudata</i>	3	supraoccipital
		<i>Potamogale velox</i>	1	supraoccipital
		<i>Setifer</i>	1	supraoccipital
		<i>Eremitalpa</i>	1-2	?
	Hyracoids	<i>Heterohyrax brucei</i>	1	supraoccipital
		<i>Procavia capensis</i>	1-4	supraoccipital
		<i>Dendrohyrax abroreus</i>	1-3	supraoccipital
	Proboscids	<i>Loxodonta africana</i>	1	supraoccipital
	Macroscelids	<i>Macroscelides proboscideus</i>	1	supraoccipital
	Tubulidentates	<i>Orycteropus capensis</i>	1	supraoccipital
		<i>Rhynchocyon cirnei</i>	1	supraoccipital
	Sirenians	<i>Dugong dugon</i>	1-4	parietal
		<i>Trichechus manatus</i>	1-4	parietal
		<i>T. senegalensis</i>	1-4	parietal
		<i>Rytina</i>	1	?
		<i>Chrysochloris asiatica</i>	3	?
Xenarthrans	Pilosaurs	<i>Cyclopes didactylus</i>	1-2	supraoccipital
		<i>Bradypus tridactylus</i>	1	supraoccipital
		<i>Tamandua tetradactyla</i>	1	supraoccipital
	Cingulates	<i>Dasyus hybridus</i>	1-2	supraoccipital

Table S1 (continued).

			Observed element number	First fusion with
Euarchontoglires	Rodents	<i>Dipodomys spp.</i>	1-4	?
		<i>Pedetes capensis</i>	1-2	?
		<i>Galea musteloides</i>	1-2	?
		<i>Hydromys chrysogaster</i>	1-2	?
		<i>Jaculus jaculus</i>	1-2	?
		<i>Micromys minutus</i>	1-2	?
		<i>Apodemus speciosus</i>	1-2	?
		<i>Octodon degu</i>	1-3	parietal
		<i>Dolichotis patagonicum</i>	1-3	parietal
		<i>Pteromys volans</i>	1	parietal
		<i>Pteromys momonga</i>	1	parietal
		<i>Spermophilus undulatus</i>	1	parietal
		<i>Petaurista leucogenys</i>	1	parietal
	Lagomorphs	<i>Lepus</i>	1-2	parietal
		<i>Oryctolagus cuniculus</i>	1	parietal
	Dermopterans	<i>Cynocephalus volans</i>	1	supraoccipital
		<i>Galeopterus variegatus</i>	1	supraoccipital
	Scandentians	<i>Tupaia belangeri</i>	1	supraoccipital
	Primates	<i>Daubentonía madagascariensis</i>	1	supraoccipital
		<i>Cebus</i>	1	supraoccipital
		<i>Hylobates syndactylus</i>	1	supraoccipital
		<i>Pan troglodytes</i>	1	supraoccipital
		<i>Homo sapiens</i>	1-4	supraoccipital
		<i>Eulemur</i>	1-2	supraoccipital
		<i>Propithecus</i>	1	supraoccipital
		<i>Loris tardigradus</i>	1	supraoccipital
		<i>Galago senegalensis</i>	1	supraoccipital
		<i>Macaca fasciata</i>	1	supraoccipital
		<i>M. fascicularis</i>	1	supraoccipital
		<i>M. mulatta</i>	1	supraoccipital
		<i>Saimiri</i>	1	supraoccipital
Laurasiatherians	Lipotyphlans	<i>Solenodon paradoxus</i>	1-2	parietal
		<i>Erinaceus europaeus</i>	1-2	supraoccipital
		<i>E. amurensis</i>	1-2	supraoccipital
		<i>Echinosorex</i>	1-2	supraoccipital
		<i>T. europaea</i>	1-2	supraoccipital
		<i>T. occidentalis</i>	1-2	supraoccipital
		<i>Scapanus orarius</i>	1-2	supraoccipital
		<i>Mogera wogura</i>	1-2	supraoccipital
		<i>Surdisorex norae</i>	1	?
		<i>S. polulus</i>	1	?
		<i>S. schitteri</i>	1	?
		<i>Congosorex phillipsorum</i>	1	?
		<i>C. polli</i>	1	?
		<i>C. verheyeni</i>	1	?
		<i>Myosorex varius</i>	1	?
		<i>M. cafer</i>	1	?
		<i>M. longicaudatus</i>	1	?
		<i>M. eisentrauti</i>	1	?
		<i>M. gnoskei</i>	1	?
		<i>M. jejei</i>	1	?
		<i>M. babaulti</i>	1	?
		<i>M. bururiensis</i>	1	?
		<i>M. rumpii</i>	1	?
		<i>M. okuensis</i>	1	?
		<i>M. geata</i>	1	?
		<i>M. kihaulei</i>	1	?
		<i>M. tenuis</i>	1	?
		<i>M. zinki</i>	1	?
		<i>Sorex araneus</i>	1	supraoccipital
		<i>S. unguiculatus</i>	1	supraoccipital
		<i>Suncus murinus</i>	1	supraoccipital
		<i>Croci dura attenuata</i>	1	supraoccipital
		<i>C. watasei</i>	1	supraoccipital

Table S1 (continued).

		Observed element number	First fusion with
Cetaceans	<i>Tursiops truncatus</i>	1-3	?
	<i>Phocoena phocoena</i>	1	?
	<i>Monodon monoceros</i>	1-4	?
	<i>Balaenoptera borealis</i>	1	supraoccipital
	<i>B. musculus</i>	1	supraoccipital
	<i>Delphinus delphinus</i>	1	?
	<i>Inia geoffrensis</i>	1	?
	<i>Stenella attenuata</i>	3	supraoccipital
	<i>Capreolus capreolus</i>	1-2	parietal
	<i>Tragelaphus scriptus</i>	1-2	parietal
Artiodactyls	<i>Cervus nippon</i>	1-2	parietal
	<i>C. elaphus</i>	1-2	parietal
	<i>C. duvauceli</i>	1-2	parietal
	<i>C. eldi</i>	1-2	parietal
	<i>Odocoileus virginianus</i>	1	parietal
	<i>Dama dama</i>	1-2	parietal
	<i>Alces alces</i>	1-2	parietal
	<i>Bos primigenius</i>	1-4	parietal
	<i>Bos grunniens</i>	1-3	?
	<i>Muntiacus reevesi</i>	1-2	parietal
	<i>Ovis dalli</i>	1-2	parietal
	<i>O. dalli</i>	1-2	parietal
	<i>Lama pacos</i>	1-2	parietal
	<i>Kobus ellipsiprymnus</i>	1-2	parietal
	<i>Tragulus</i>	1-2	parietal
Carnivorans	<i>Ailuropoda melanoleuca</i>	1	supraoccipital
	<i>Ursus</i>	1	supraoccipital
	<i>Canis</i>	1-2	supraoccipital
	<i>Speothos</i>	1	supraoccipital
	<i>Lutra lutra</i>	1	supraoccipital
	<i>Mustela</i>	1	supraoccipital
	<i>Procyon lotor</i>	1	supraoccipital
	<i>P. pygmaeus</i>	1	supraoccipital
	<i>Vulpes vulpes</i>	1	supraoccipital
	<i>Panthera leo</i>	1	supraoccipital
	<i>P. onca</i>	1-3	supraoccipital
	<i>P. pardus</i>	1	supraoccipital
	<i>Puma concolor</i>	1-3	supraoccipital
	<i>Felis domestica</i>	1-2	supraoccipital
	<i>Arctocephalus</i>	1	supraoccipital
	<i>Callorhinus ursinus</i>	1	supraoccipital
	<i>Phoca</i>	1-3	supraoccipital
Pholidotans	<i>Manis</i>	1	supraoccipital
Perissodactyls	<i>Equus</i>	1-2	parietal
	<i>Tapirus</i>	1-2	parietal
	<i>Rhinoceros</i>	1	parietal
Chiropterans	<i>Diceros</i>	1	parietal
	<i>Hipposideros</i>	1-2	supraoccipital
	<i>Kerivoula</i>	1-2	supraoccipital
	<i>Hesperoptenus</i>	1-2	supraoccipital
	<i>Rhinolophus</i>	1-2	supraoccipital
	<i>Miniopterus</i>	1-2	supraoccipital
	<i>Roussettus</i>	1	supraoccipital
	<i>Artibeus</i>	1	supraoccipital
	<i>Pteropus lylei</i>	1	supraoccipital
	<i>Dobsonia pannietensis</i>	1	supraoccipital

Part D. References for the Supplementary Information

1. Romer AS & Parsons TS (1978) *The Vertebrate Body* (Saunders, Philadelphia).
2. Schultze HP (1993) in *The Skull: Patterns of Structural and Systematic Diversity*, eds Hanken J & Hall BK (University of Chicago Press, Chicago), Vol 2, pp 189-254.
3. Jarvik E (1980) *Basic Structure and Evolution of Vertebrates* (Academic Press, New York).
4. Janvier P (1996) *Early Vertebrates* (Clarendon Press, Oxford).
5. Clack JA (2007) Devonian climate change, breathing, and the origin of the tetrapod stem group. *Integr Comp Biol* 47:510-523.
6. Benton MJ (2005) *Vertebrate Paleontology* (Wiley-Blackwell, New York).
7. Goodrich ES (1931) *Studies on the Structure and Development of Vertebrates* (Dover Publications, New York).
8. Trueb L (1993) in *The Skull: Patterns of Structural and Systematic Diversity*, eds Hanken J & Hall BK (University of Chicago Press, Chicago), Vol 2, pp 255-343.
9. Mickoleit G (2004) *Phylogenetische Systematik der Wirbeltiere* (Pfeil Verlag, München). German
10. Jenkins FA, Jr., Walsh DM, & Carroll RL (2007) Anatomy of *Eocaecilia micropodia*, a limbed caecilian of the Early Jurassic. *Bull Mus Comp Zool* 158:285-365.
11. Romer AS (1956) *Osteology of the Reptiles* (University of Chicago Press, Chicago).
12. Carroll RL (1969) A Middle Pennsylvanian captorhinomorph, and the interrelationships of primitive reptiles. *J Paleontol* 43:151-170.
13. Anderson JS, Reisz RR, Scott D, Fröbisch NB, & Sumida SS (2008) A stem batrachian from the Early Permian of Texas and the origin of frogs and salamanders. *Nature* 453:515-518.
14. Marjanović D & Laurin M (2009) The origin(s) of modern amphibians: a commentary. *Evol Biol* 36:336-338.
15. Roček Z (1988) Origin and evolution of the frontoparietal complex in anurans. *Amphibia-Reptilia* 9:385-403.
16. Smirnov SV (1997) Additional dermal ossifications in the anuran skull: morphological novelties or archaic elements? *Russ J Herpetol* 4:17-27.
17. Carroll RL (1988) *Vertebrate Paleontology and Evolution* (Freeman, New York).
18. Parrington F (1937) V.—A note on the supratemporal and tabular bones in reptiles. *J Nat Hist* 20:69-76.
19. Broom R (1914) Some points in the structure of the diadectid skull. *Bull Am Mus Nat Hist* 33:109-114.
20. Klembara J (2001) Postparietal and prehatching ontogeny of the supraoccipital in *Alligator mississippiensis* (Archosauria, Crocodylia). *J Morphol* 249:147-153.
21. Jollie M (1973) *Chordate Morphology* (Reinhold, New York).
22. Starck D (1979) *Vergleichende Anatomie der Wirbeltiere* (Springer, Berlin). German
23. Sánchez-Villagra MR, et al. (2009) Skeletal development in the Chinese soft-shelled turtle *Pelodiscus sinensis* (Testudines: Trionychidae). *J Morphol* 270:1381-1399.
24. Werneburg I, Hugi J, Müller J, & Sánchez-Villagra MR (2009) Embryogenesis and ossification of *Emydura subglobosa* (Testudines, Pleurodira, Chelidae) and patterns of turtle development. *Dev Dyn* 238:2770-2786.
25. Gaupp E (1900) Das chondrocranium von *Lacerta agilis*. *Anat Embryol* 15:434-594. German
26. Bellairs AA (1969) *The Life of Reptiles* (Universe, New York).
27. Conrad JL (2008) Phylogeny and systematics of Squamata (Reptilia) based on morphology. *Bull Am Mus Nat Hist* 310:1-182.
28. Müller J, et al. (2011) Eocene lizard from Germany reveals amphisbaenian origins. *Nature* 473:364-367.
29. Werner G (1962) Das Cranium der Brückenechse, *Sphenodon punctatus* Gray, von 58 mm Gesamtlänge. *Anat Embryol* 123:323-368. German
30. Rieppel O (2000) in *Handbuch der Paläoherpetologie, Teil 12A*, eds Rieppel O, Zaher H, Tchernov E, & Polcyn MJ (Friedrich Pfeil, München), pp 1-134. German
31. Parker WK (1876) VII. On the structure and development of the bird's skull. *Trans Linn Soc Lond* 1:99-154.
32. Zusi RL (1993) in *The Skull: Patterns of Structural and Systematic Diversity*, eds Hanken J & Hall BK (University of Chicago Press, Chicago), Vol 2, pp 391-437.
33. Müller H (1963) Die Morphologie und Entwicklung des Craniums von *Rhea americana* Linné II. Viszeralskelett, Mittelohr und Osteocranium. *Zeitschrift für Wissenschaftliche Zoologie* 168:35-118. German
34. Chatterjee S (1997) *The Rise of Birds: 225 Million Years of Evolution* (Johns Hopkins University Press Baltimore, Maryland).
35. Bühler P (1985) in *The Beginnings of Birds: Proceedings of the International Archaeopteryx Conference Eichstätt*, eds Hecht MK, Ostrom JH, Viohl G, & Wellnhofer P (Freunde des Jura-Museums Eichstätt, Eichstätt), pp 135-140.
36. Raath MA, Hecht M, Ostrom J, Viohl G, & Wellnhofer P (1985) in *The Beginnings of Birds: Proceedings of the International Archaeopteryx Conference Eichstätt*, eds Hecht MK, Ostrom JH, Viohl G, & Wellnhofer P (Freunde des Jura-Museums Eichstätt, Eichstätt), pp 219-227.
37. Reisz R (1986) *Pelycosauria. Handbuch der Paläoherpetologie* (Gustav Fischer Verlag, Stuttgart). German
38. Romer AS & Price LI (1940) Review of the Pelycosauria. *Geological Society of America Special Papers* 28:1-538.
39. Watson DMS (1916) Reconstructions of the skulls of three pelycosaurs in the American Museum of Natural History. *Bull Am Mus Nat Hist* 35:637-648.
40. Broom R (1914) A further comparison of the South African dinocephalians with the American pelycosaurs. *Bull Am Mus Nat Hist* 33:135-141.
41. Parrington F (1946) LXXIV.—On the quadratojugal bone of synapsid reptiles. *J Nat Hist* 13:780-786.
42. Crompton A (1958) The cranial morphology of a new genus and species of ictidosaurian. *Proc Zool Soc Lond* 130:183-216.
43. Hammer W & Cosgriff J (1981) *Myosaurus gracilis*, an anomodont reptile from the Lower Triassic of Antarctica and South Africa. *J Paleontol* 55:410-424.

44. Cox CB (1991) The Pangaea dicynodont *Rechnisaurus* and the comparative biostratigraphy of Triassic dicynodont faunas. *Palaeontology* 34:767-784.
45. Maddin HC, Sidor CA, & Reisz RR (2008) Cranial anatomy of *Ennatosaurus tecton* (Synapsida: Caseidae) from the Middle Permian of Russia and the evolutionary relationships of Caseidae. *J Vert Paleontol* 28:160-180.
46. Martinez RN, May CL, & Forster CA (1996) A new carnivorous cynodont from the Ischigualasto Formation (Late Triassic, Argentina), with comments on eucynodont phylogeny. *J Vert Paleontol* 16:271-284.
47. Fröbisch J & Reisz RR (2008) A new species of *Emydops* (Synapsida, Anomodontia) and a discussion of dental variability and pathology in dicynodonts. *J Vert Paleontol* 28:770-787.
48. Fröbisch J (2007) The cranial anatomy of *Kombuisia frerensis* Hotton (Synapsida, Dicynodontia) and a new phylogeny of anomodont therapsids. *Zool J Linn Soc* 150:117-144.
49. Broom R (1936) On the structure of the skull in the mammal-like reptiles of the suborder Therocephalia. *Philos Trans R Soc Lond, B, Biol Sci* 226:1-42.
50. Kemp TS (1979) The primitive cynodont *Procynosuchus*: functional anatomy of the skull and relationships. *Philos Trans R Soc Lond, B, Biol Sci* 285:73-122.
51. Kemp T (1972) Whaitsiid Therocephalia and the origin of cynodonts. *Philos Trans R Soc Lond, B, Biol Sci* 264:1-54.
52. Brink AS (1951) On the genus *Lystrosaurus* Cope. *Trans Roy Soc S Afr* 33:107-120.
53. Broom R (1932) *The Mammal-like Reptiles of South Africa and the Origin of Mammals* (HF & G. Witherby, London).
54. Kielan-Jaworowska Z, Cifelli R, & Luo ZX (2004) *Mammals from the Age of Dinosaurs: Origins, Evolution, and Structure* (Columbia University Press, New York).
55. de Beer GR (1937) *The Development of the Vertebrate Skull* (Oxford University Press, London).
56. Kuhn HJ (1971) Die Entwicklung und Morphologie des Schädels von *Tachyglossus aculeatus*. *Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft* 528:1-192. German
57. Zeller U (1989) Die Entwicklung und Morphologie des Schädels von *Ornithorhynchus anatinus* (Mammalia: Prototheria: Monotremata). *Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft* 545:1-188. German
58. van Bemmelen JF (1901) *Der Schädelbau der Monotremen* (G. Fischer, Jena). German
59. Landry SO (1964) The form and phylogenetic history of the mammalian interparietal bone. *Am Zool* 4:400-400.
60. Kierdorf U & Kierdorf H (1998) Studies on the parietal region of the cervid skull III. On the occurrence of an interparietal in *Cervus*. *Anat Histol Embryol* 27:193-196.
61. Kuhn HJ & Zeller U (1987) in *Mammalia Depicta: Morphogenesis of the Mammalian Skull*, eds Kuhn HJ & Zeller U (Verlag Paul Parey, Hamburg), pp 51-70.
62. Owen R (1868) *On the Anatomy of Vertebrates. Vol 2. Birds and Mammals* (Longmans, Green, and Company, London).
63. Toeplitz C (1920) Bau und Entwicklung des Knorpelschädels von *Didelphys marsupialis*. *Zoologica* 27:1-83. German
64. Clark CT & Smith KK (1993) Cranial osteogenesis in *Monodelphis domestica* (Didelphidae) and *Macropus eugenii* (Macropodidae). *J Morphol* 215:119-149.
65. Wible JR (2003) On the cranial osteology of the short-tailed opossum *Monodelphis brevicaudata* (Didelphidae, Marsupialia). *Ann Carnegie Mus* 72:137-202.
66. Cooper CE (2011) *Myrmecobius fasciatus* (Dasyuromorphia: Myrmecobiidae). *Mammalian Species* 43:129-140.
67. Abdala F, Flores DA, & Giannini NP (2001) Postweaning ontogeny of the skull of *Didelphis albiventris*. *J Mammal* 82:190-200.
68. Forasiepi AM (2009) Osteology of *Arctodictis sinclairi* (Mammalia, Metatheria, Sparassodonta) and phylogeny of Cenozoic metatherian carnivores from South America. *Monogr Mus Argentino Cienc Nat*:1-174.
69. Voss RS & Jansa SA (2009) Phylogenetic relationships and classification of didelphid marsupials, an extant radiation of New World metatherian mammals. *Bull Am Mus Nat Hist*:1-177.
70. Sánchez-Villagra M, et al. (2007) Exceptionally preserved North American Paleogene metatherians: adaptations and discovery of a major gap in the opossum fossil record. *Biol Lett* 3:318.
71. Marshall L & de Muizon C (1995) Part II. The skull. *Pucadelphys andinus* (Marsupialia, Mammalia) from the Early Paleocene of Bolivia. *Memoires du Museum National d'Histoire Naturelle* 165:21-90.
72. de Muizon C (1998) *Mayulestes ferox*, a borhyaenoid (Metatheria, Mammalia) from the early Palaeocene of Bolivia. Phylogenetic and palaeobiologic implications. *Geodiversitas* 20:19-142.
73. Wegner RN (1960) Das Os interparietale als integrierender Bestandteil der Säuger- und Theromorphenschädel. *Gegenbaurs Morphol Jahrb* 100:375-419. German
74. Roth S (1903) Los ungulados sudamericanos. *Anales del Museo de La Plata* 5:1-36. Spanish
75. Giebel C (1874) Säugetiere: Mammalia. *Dr HG Bronn's Klassen und Ordnungen des Thierreichs, wissenschaftlich dargestellt in Wort und Bild* 6:1-36. German
76. Lawlor TE (1979) *Handbook to the Orders and Families of Living Mammals* (Mad River Press, California) p 151.
77. Parker WK (1885) On the structure and development of the skull in the Mammalia. Part II. Edentata. *Philos Trans R Soc Lond* 176:1-119.
78. Parker WK (1885) On the structure and development of the skull in the Mammalia. Part III. Insectivora. *Philos Trans R Soc Lond* 176:121-275.
79. Schunke AC & Zeller U (2010) Chondrocranium and dermal bones of the lowland streaked tenrec *Hemicentetes semispinosus* (Afrosoricida, Tenrecidae) and their comparison with *Potamogale* and other insectivoran-grade placental mammals. *Vertebr Zool* 60:37-72.
80. Leimgruber C (1939) Le crâne primordial d'un embryon de *Centetes ecaudatus*. *Biomorphosis* 1:545-585. French
81. Roux GH (1947) The cranial development of certain ethiopian "insectivores" and its bearing on the mutual affinities of the group. *Acta Zool* 28:165-397.
82. Broom R (1916) On the structure of the skull in *Chrysochloris*. *Proc Zool Soc Lond* 86:449-458.
83. Van der Merwe N, Bezuidenhout A, & Seegers C (1995) The skull and mandible of the African elephant (*Loxodonta africana*). *Onderstepoort J Vet Res* 62:245-260.

84. Kingsley JS (1925) *The Vertebrate Skeleton from the Developmental Standpoint* (P. Blakiston's Son & Co., Philadelphia).
85. Pucek Z (1962) The occurrence of wormian bones (ossicula wormiana) in some mammals. *Acta Theriol* 6:33-51.
86. Beer JR (1965) The interparietal in kangaroo rats. *Southwest Nat* 10:145-150.
87. Tullberg T (1899) *Über das system der Nagethiere, eine phylogenetische studie* (Nova Acta Regiae Societatis Scientiarum Upsaliensis, Upsala). German
88. Wortman JL (1920) On some hitherto unrecognized reptilian characters in the skull of the Insectivora and other mammals. *Proc US Natl Mus* 57:1-52.
89. Zeller U (1987) in *Mammalia depicta: morphogenesis of the mammalian skull*, eds Kuhn HX & Zeller U (Verlag Paul Parey, Hamburg), pp 17-50.
90. Srivastava H (1992) Ossification of the membranous portion of the squamous part of the occipital bone in man. *J Anat* 180:219-224.
91. Matsumura G, Uchiumi T, Kida K, Ichikawa R, & Kodama G (1993) Developmental studies on the interparietal part of the human occipital squama. *J Anat* 182:197-204.
92. Eaglen RH (1983) Parallelism, parsimony, and the phylogeny of the lemuridae. *Int J Primatol* 4:249-273.
93. Starck D (1962) Das cranium von *Propithecus* spec. (Prosimiae, Lemuriformes, Indridae). *Bibl Primat* 1:163-196. German
94. Hershkovitz P (1977) *Living New World Monkeys (Platyrrhini): with an Introduction to Primates* (University of Chicago Press, Chicago).
95. Chopra SRK (1957) The cranial suture closure in monkeys. *Proc Zool Soc Lond* 128:67-112.
96. Wible JR (2008) On the cranial osteology of the Hispaniolan solenodon, *Solenodon paradoxus* Brandt, 1833 (Mammalia, Lipotyphla, Solenodontidae). *Ann Carnegie Mus* 77:321-402.
97. Goswami A & Prochel J (2007) Ontogenetic morphology and allometry of the cranium in the common european mole (*Talpa europaea*). *J Mammal* 88:667-677.
98. de Beer GR (1929) The development of the skull of the shrew. *Philos Trans R Soc Lond, B, Biol Sci* 217:411-480.
99. Augier M (1931) Quelques problèmes relatifs au développement du crâne. Voûte occipitale chez l'Homme et d'autres Mammifères. *Arch Anat Histol Embryol* 13:34-66. French
100. Kuratani S (1994) [Mysteries of occipital bones II. - Mechanism of the segmental patterning]. *The Bone* 8:137-150 Japanese.
101. Peterhans JCK, et al. (2010) African shrews endemic to the Albertine Rift: two new species of *Myosorex* (Mammalia: Soricidae) from Burundi and the Democratic Republic of Congo. *J East Afr Nat Hist* 99:103-128.
102. Eales NB (1950) The skull of the foetal narwhal, *Monodon monoceros* L. *Philos Trans R Soc Lond, B, Biol Sci* 235:1-33.
103. Ridewood WG (1923) Observations on the skull in foetal specimens of whales of the genera *Megaptera* and *Balaenoptera*. *Philos Trans R Soc Lond, B, Biol Sci* 211:209-272.
104. Schulte HW (1916) The sei whale (*Balaenoptera borealis* Lesson). Anatomy of a foetus of *Balaenoptera borealis*. *Monographs of the pacific cetacea Memoirs of American Museum of Natural History, New Series* 1:389-499.
105. Kesteven LH (1941) Some features in the anatomy and later development of the head of *Delphinus delphinus* Linné. *Rec Aust Mus* 21:59-80.
106. Moran MM, Nummela S, & Thewissen J (2011) Development of the skull of the pantropical spotted dolphin (*Stenella attenuata*). *Anat Rec* 294:1743-1756.
107. Starck D (1967) Le crane des Mammifères. *Traité de Zoologie* 16:405-549. German
108. Starck D (1995) *Lehrbuch der Speziellen Zoologie. Wirbeltiere. Teil 5, Säugetiere* (Gustav Fischer-Verlag, Jena). German
109. Shapiro R & Robinson F (1976) The os incae. *Am J Roentgenol* 127:469-471.
110. Rörig A (1904) Das wachstum des schädels von *Capreolus vulgaris*, *Cervus elaphus* und *Dama vulgaris*. *Anat Anz* 26:17-25. German
111. Chapman NG (2003) Observations on the parietal region of the skull of *Muntiacus reevesi*. *Zeitschrift für Jagdwissenschaft* 49:237-241.
112. Yilmaz S, Dinç G, & Toprak B (2000) Macro-anatomical investigations on skeletons of otter (*Lutra lutra*). III. Skeleton axiale. *Veterinarski Arhiv* 70:191-198.
113. Korth WW & Baskin JA (2009) A new species of *Leptarctus* (Carnivora, Mustelidae) from the late Clarendonian (late Miocene) of Kansas. *Ann Carnegie Mus* 78:29-44.
114. de Villa-Meza A, Avila-Flores R, Cuarón AD, & Valenzuela-Galván D (2011) *Procyon pygmaeus* (Carnivora: Procyonidae). *Mammalian Species* 43:87-93.
115. Guzmán JA, D Elía G, & Ortiz JC (2009) Variación geográfica del zorro *Lycalopex culpaeus* (Mammalia, Canidae) en Chile: implicaciones taxonómicas. *Revista de Biología Tropical* 57:421-432. Spanish
116. Weber M & Abel O (1928) *Die Säugetiere. Einführung in die Anatomie und Systematik der Recenten und Fossilen Mammalia*. (Gustav Fischer Verlag, Jena). German
117. Gregory WK (1910) The orders of mammals. *Bull Am Mus Nat Hist* 27:1-524.
118. Lundelius EL & Slaughter BH (1976) in *Athlon: essays on paleontology in honour of Loris Shano Russell*, ed Churcher CS (Royal Ontario Museum Life Science Miscellaneous Publications, Toronto, Canada), pp 226-243.
119. Padilla M, Dowler RC, & Downer CC (2010) *Tapirus pinchaque* (Perissodactyla: Tapiridae). *Mammalian Species* 42:166-182.
120. Holbrook LT (2002) The unusual development of the sagittal crest in the Brazilian tapir (*Tapirus terrestris*). *J Zool* 256:215-219.
121. Cuvier G (1836) *Recherches sur les ossements fossiles, où l'on rétablit les caractères de plusieurs animaux dont les révolutions du globe ont détruit les espèces* (E. d'Ocagne, Paris). French

122. Fawcett E (1919) The primordial cranium of *Miniopterus schreibersi* of the 17 millimetre total length stage. *J Anat* 53:315-350. German
123. Fehse O (1990) Ein Beitrag zur Morphologie des Primordialcraniums von *Rousettus aegyptiacus* (Geoffroy, 1810), (Megachiroptera, Mammalia). Diploma thesis (University of Tübingen, Tübingen). German
124. Hartmann L (1990) Das Primordialcranium von *Artibeus jamaicensis* Leach 1821 (Phyllostomidae, Microchiroptera). Diploma thesis (University of Tübingen, Tübingen). German
125. Giannini NP, Wible JR, & Simmons NB (2006) On the cranial osteology of Chiroptera. I. *Pteropus* (Megachiroptera: Pteropodidae). *Bull Am Mus Nat Hist* 295:1-134.