

FROBENIUS MAPS AND ALGEBRAIC VARIETIES

AKIYOSHI SANNAI

ABSTRACT. Many characterization of abelian varieties in positive characteristic is given by several authors. In this article, we give a characterization of abelian varieties by the language of $F_*\mathcal{O}_X$ and its Poincare bundles.

Let k be an algebraically closed field of characteristic $p > 0$. Let X be a smooth proper variety over k . When does X satisfy the following property (*)?

- (*) $F_*\mathcal{O}_X \simeq \bigoplus_j M_j$ where $F : X \rightarrow X$ is the absolute Frobenius morphism and each M_j is a line bundle.

For example, an arbitrary smooth proper toric variety satisfies this property (*) (cf. [Achinger][Thomsen]). Thus there are many varieties which satisfy (*). But every toric variety has negative Kodaira dimension. On the other hand, In [ST] we show that ordinary abelian varieties satisfy (*) and this property gives the characterization of the ordinary abelian varieties.

Theorem 0.1. *Let k be an algebraically closed field of characteristic $p > 0$. Let X be a smooth projective variety over k . Assume the following conditions.*

- *For infinitely many $e \in \mathbb{Z}_{>0}$, $F_*^e\mathcal{O}_X \simeq \bigoplus_j M_j$ where each M_j is an invertible sheaf.*
- *$\kappa(X) \geq 0$ where $\kappa(X)$ is the Kodaira dimension of X .*

Then X is an ordinary abelian variety.

By the Theorem above, it is natural to ask the same criterion holds only for $F_*\mathcal{O}_X$. But in [ES], we gave the counter example for that problem.

Theorem 0.2. *Let X be a non-abelian smooth projective surface. Then X satisfies the following conditions if and only if X is an F -split Igusa surface:*

- $F_*\mathcal{O}_X$ is isomorphic to direct sum of line bundles.
- K_X is pseudo-effective.

We also showed that if we consider $e > 1$, then the same characterization holds.

Theorem 0.3. *Let X be a smooth projective variety over k of characteristic $p > 2$ (resp. $p > 0$). X is an ordinary abelian variety if and only if the following conditions hold:*

- $F_*\mathcal{O}_X$ (resp. $F_*^2\mathcal{O}_X$) is isomorphic to a direct sum of line bundles.
- K_X is pseudo-effective.

Another natural question is how can we characterize non ordinary abelian varieties by the language of $F_*\mathcal{O}_X$. In this article we give the characterization by using Poincare bundle on albanese varieties. For details of Poincare bundles and albanese maps, see [FGAex, Section 9].

Theorem 0.4. *Let X be a smooth projective variety over k of characteristic p . There is a canonical map $\Phi : \tilde{\mathcal{P}}_X \rightarrow F_*\mathcal{O}_X$ with the following properties:*

- $\tilde{\mathcal{P}}_X \simeq p_{1*}\mathcal{P}_{X \times \ker V}$ where $\mathcal{P}$ is the Poincare bundle on $X \times \text{Pic}^0 X$ and V is the Verschiebung map on $\text{Pic}^0 X$.
- Φ is isomorphic if and only if X is an abelian variety.
- Φ is generically isomorphic if and only if X has maximal albanese dimension.

We overview the proof of Theorem 0.4. At first, we construct the map $\Phi : \tilde{\mathcal{P}}_X \rightarrow F_*\mathcal{O}_X$.

Lemma 0.5. *Let X be a smooth projective variety over k of characteristic p and $\alpha_X : X \rightarrow \text{Alb}(X)$ be the albanese map of X . Then we have*

$$\alpha_X^* \tilde{\mathcal{P}}_{\text{Alb}(X)} \simeq \tilde{\mathcal{P}}_X$$

Proof. $\alpha_X : X \rightarrow \text{Alb}(X)$ induces the following map;

$$\alpha_X^* : \text{Pic}^0 \text{Pic}^0 \text{Pic}_{red}^0(X) \simeq \text{Pic}^0(X)$$

Here we can see that the kernel of the Verschiebung maps corresponds each other. Hence, we have

$$\alpha_X^* \text{Ker } V_{\text{Pic}^0(X)} \simeq \text{Ker } V_{\text{Pic}^0 \text{Pic}^0 \text{Pic}_{red}^0(X)}$$

. By considering the universal bundles on them, we get the assersion. $\square$

Lemma 0.6. *Let X be a smooth projective variety over k of characteristic p . There is a canonical map $\Phi : \tilde{\mathcal{P}}_X \rightarrow F_*\mathcal{O}_X$.*

Proof. We use the following theorem.

Theorem 0.7 (Oda). *Let $f : X \rightarrow Y$ be an isogeny of abelian varieties over k . Set $\hat{f} : \hat{Y} \rightarrow \hat{X}$ to be the dual of f . Let $L \in \text{Pic}^0(X)$. Then, $f_*L \simeq \text{pr}_{1*}(\mathcal{P}_Y|_{Y \times \hat{f}^{-1}([L])})$*

$$f_*L \simeq \text{pr}_{1*}(\mathcal{P}_Y|_{Y \times \hat{f}^{-1}([L])})$$

where $\mathcal{P}_Y$ is the normalized Poincare line bundle of $(Y, 0)$.

By applying this theorem to $f = F$ and $L = \mathcal{O}_X$, we have $\Phi : \tilde{\mathcal{P}}_X \rightarrow F_*\mathcal{O}_X$ for any abelian varieties X . (Note that Verschiebung map is defined by the dual of relative Frobenius map.)

In general, we have the base change map $\alpha^*F_*\mathcal{O}_{\text{Alb}(X)} \rightarrow F_*\mathcal{O}_X$. Then by the discussion above and lem 0.6,

$$\alpha^*F_*\mathcal{O}_{\text{Alb}(X)} \simeq \alpha^*\tilde{\mathcal{P}}_{\text{Alb}(X)} \simeq \tilde{\mathcal{P}}_X$$

Hence, by composing them, we get the assersion. $\square$

Claim 0.8. *If Φ_X is isomorphism, then X has maximal albanese dimension, namely $\dim X = \dim \alpha(X)$*

Proof. By Stein factrizaton theorem, we have the following decomposition;

$$\alpha : X \xrightarrow{f} Y \xrightarrow{g} Z \xrightarrow{h} A = \text{Alb}(X)$$

where g, h is finite, and g is purely inseparable, h is separable, Y, Z is normal.

We first see the map

$$\Phi_Y : \tilde{\mathcal{P}}_Y \rightarrow F_*\mathcal{O}_Y$$

. Pulling back by f , we have

$$\tilde{\Phi}_Y : f^*\tilde{\mathcal{P}}_Y \xrightarrow{f^*\Phi_Y} f^*F_*\mathcal{O}_Y \rightarrow F_*\mathcal{O}_X$$

where the second map is base change map.

We also have

$$f^*\tilde{\mathcal{P}}_Y \simeq f^*g^*\tilde{\mathcal{P}}_{\text{Alb}(X)} \simeq \alpha^*\tilde{\mathcal{P}}_{\text{Alb}(X)} \simeq \tilde{\mathcal{P}}_X.$$

Hence, we see that $\tilde{\Phi}_Y = \Phi_X$. Hence, by calculating the rank of $F_*\mathcal{O}$, we have $\dim X \leq \dim Y$. This implies $\dim X = \dim Y = \dim \alpha(X)$. $\square$

Claim 0.9. *If Φ_X is isomorphism, then Φ_Y is also isomorphism.*

Proof. Since X has maximal albanese dimension, f is a birational contraction. Let U, V be the isomophic locus of f in X, Y respectively. Then U, V have codimension two in their ambient space. Here we have

$$F_*\mathcal{O}_Y|U \simeq F_*\mathcal{O}_X|V \simeq \mathcal{P}_X|V \simeq \mathcal{P}_Y|U$$

Since all the sheaf above are reflexive and all the varieties above are normal, and U, V are codimension two subset, we get $\Phi_Y : \mathcal{P}_Y \simeq F_*\mathcal{O}_Y$. $\square$

Claim 0.10. *There is a decomposition*

$$Y \rightarrow Z_1 \rightarrow \dots \rightarrow Z_i \rightarrow Z_{i+1} \dots \rightarrow Z$$

such that $h_i : Z_i \rightarrow Z_{i+1}$ is a purely inseparable extension of height 1 of normal varieties. Futhermore, if Φ_Y is an isomorphism, then Φ_{Z_i} is also an isomorphism.

Proof. The first assersion follows by taking field extensions of height 1 and normalizations in the fields. We see the second assersion. Since Z_i is normal, $F_*\mathcal{O}_{Z_i}$ is reflexive. Hence we may assume Z_{i+1} is a spectrum of a DVR. Since torsion free module over DVR is faithfully flat, h_i^* is a faithfully flat functor. Hence by the induction starting from Y , we have the second assersion. $\square$

By standard calculation of Grothendiek duality and Kunz's theorem, we have the following proposition.

Proposition 0.11. *Let X be normal projective variety over k . Assume that Φ_X is an isomorphism. Then X is smooth and $-(p-1)K_X$ is an effective divisor.*

We prove the main theorem for purelu inseparable extension of height 1.

Proposition 0.12. *Let $\alpha : Z \rightarrow A$ be the albanese map of normal variety Z . Assume that α is purely inseparable of height 1 and Φ_Z is an isomorphism. Then α is an isomorphism.*

Proof. By [Eke], any purely inseparable of height 1 morphism can be described by a quotient of 1-foliation. Namely, there is a 1-foliation $\mathcal{F} \subset \mathcal{T}_A$ such that $Z \simeq A/\mathcal{F}$ and $\omega_Z^p \simeq \alpha^*\omega_A \otimes (\det \mathcal{F})^{(1-p)}$. Since A is an abelian variety, $\mathcal{T}_A$ is trivial. This implies $\det \mathcal{F} \subset \oplus \mathcal{O}_A$. Hence $\det \mathcal{F}$ is negative line bundle, put $\det \mathcal{F} \simeq \mathcal{O}(-D)$ for an effective divisor D . Then we have

$$\omega_Z^p \simeq \mathcal{O}_Z((p-1)D).$$

Prop 0.11 and this implies $\kappa(Z) = 0$. Hence by [HP], we get the assertion. $\square$

By claim 0.10 and Prop 0.12, h is an isomorphism. Next, we show g is an isomorphism.

Claim 0.13. *g is an isomorphism.*

Proof. Put

$$K_Y = g^*K_A + R$$

where R is the ramification divisor of separable morphism g . Since $K_A = 0$ and by Prop 0.11, we have $R = 0$.

Hence $\alpha : Y \rightarrow A$ is etale in codimension one. Then, by the Zariski–Nagata purity, α is etale. By [Mumford, Section 18, Theorem], Y is also an abelian variety and we are done. $\square$

Claim 0.14. *f is an isomorphism*

Proof. We can write

$$K_X = f^*K_A + E$$

where E is an f -exceptional divisor. Since A is terminal (cf. [KM, Section 2.3]), E is effective. By Prop 0.12, we have $K_X \equiv 0$. Since $K_X \equiv 0$, we see that E is f -nef. By the negativity lemma (cf. [KM, Lemma 3.39]), we see $E = 0$. Thus, the codimension of $\text{Ex}(f)$ in X is at least two. Since A is smooth, f is an isomorphism. $\square$

Acknowledgments. The author would like to thank Professors Taro Fujisawa, Takuzo Okada, Taro Sano, for their invitation to the symposium The author was partially supported by JSPS Grant-in-Aid for Young Scientists (B) 16K17581.

REFERENCES

- [Achinger] P. Achinger, *A characterization of toric varieties in characteristic p* , to appear in Int. Math. Res. Notices.
- [Atiyah1] M. F. Atiyah, *On the Krull-Schmidt theorem with application to sheaves*, Bulletin de la S. M. F., tome **84** (1956), 307–317.
- [Atiyah2] M. F. Atiyah, *Vector bundles over an elliptic curve*, Proc. London Math. Soc, **7** (1957), 414–452.
- [Eke] T. Ekedahl, *Canonical models of surfaces of general type in positive characteristic*, Inst. Hautes Etudes Sci. Publ. Math. **67** (1988), 97 – 144.
- [ES] S. Ejiri, A. Sannai, *A characterization of ordinary abelian varieties by the Frobenius push-forward of the structure sheaf II.* to appear in Mathematical research notices.

- [FGAex] B. Fontechi, L. Göttsche, L. Illusie, S. L. Kleiman, N. Nithure, A. Vistoli, *Fundamental Algebraic Geometry: Grothendieck's FGA Explained*, Math. Surveys and Monographs, Vol. **123** (2005).
- [HP] C. D. Hacon, Z. Patakfalvi, *On the characterization of abelian varieties in characteristic $p > 0$* , arXiv:1602.01791
- [Hirokado] M. Hirokado, *Zariski surfaces as quotients of Hirzebruch surfaces by 1-foliations*, Yokohama Math. J. vol. 47 (2000), 103–120.
- [KS] Y. Kitadai, H. Sumihiro, *Canonical filtrations and stability of direct images by Frobenius morphisms*, Tohoku Math. J. **60**, 287–301 (2008).
- [KM] J. Kollár, S. Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts in Mathematics, Vol. **134**, 1998.
- [Liedtke] C. Liedtke, *The Canonical Map and Horikawa Surfaces in Positive Characteristic*, Int. Math. Res. Notices (2013) Vol. **2013**(2), 422–462.
- [MeSr] V. B. Mehta, V. Srinivas, *Varieties in positive characteristic with trivial tangent bundle*, Compositio Math., tome **64** no. 2, 191–212 (1987).
- [Mumford] D. Mumford, *Abelian varieties*, Tata Institute of Fundamental Research Studies in Mathematics, No. **5**, (1970).
- [MuSr] M. Mustață, V. Srinivas, *Ordinary varieties and the comparison between multiplier ideals and test ideals*, Nagoya Math. J. Volume **204** (2011), 125–157
- [Oda] T. Oda, *Vector bundles on an elliptic curve*, Nagoya Math. J., Vol. **43**, 41–72 (1971).
- [SS] K. Schwede, K. E. Smith, *Globally F -regular and log Fano varieties*, Adv. in Math., Vol. **224** Issue 3, 863–894, (2010).
- [ST] A. Sannai and H. Tanaka. *A characterization of ordinary abelian varieties by the Frobenius push-forward of the structure sheaf*. Math. Ann., **366** 1067 – 1087, 2016.
- [Sun] X. Sun, *Direct images of bundles under Frobenius morphism*, J. Algebra **226** (2000), no. 2, 865–874
- [Thomsen] J. F. Thomsen, *Frobenius direct images of line bundles on toric varieties*, J. Algebra **226** (2000), no. 2, 865–874

RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES, KYOTO UNIVERSITY,
KYOTO 606-8502, JAPAN

E-mail address: sannai@kurims.kyoto-u.ac.jp