

1 **Title**

2 NADPH regulates human NAD kinase, a NADP⁺-biosynthetic
3 enzyme

4
5 **Author names and affiliations**

6 Kazuto Ohashi, Shigeyuki Kawai, Mari Koshimizu, Kousaku Murata*

7
8 Laboratory of Basic and Applied Molecular Biotechnology, Division of Food and
9 Biological Science, Graduate School of Agriculture, Kyoto University, Uji, Kyoto
10 611-0011, Japan

11
12 ***Corresponding author**

13 Kousaku Murata

14 Address: Laboratory of Basic and Applied Molecular Biotechnology,
15 Division of Food and Biological Science, Graduate School of Agriculture,
16 Kyoto University, Uji, Kyoto 611-0011, Japan

17 Telephone number: +81-774-38-3766

18 Fax number: +81-774-38-3767

19 E-mail: kmurata@kais.kyoto-u.ac.jp

20

21

22

23

Abstract

NAD kinase (NADK, EC 2.7.1.23) is the sole NADP⁺-biosynthetic enzyme that catalyzes phosphorylation of NAD⁺ to yield NADP⁺ using ATP as a phosphoryl donor, and thus, plays a vital role in the cell and represents a potentially powerful antimicrobial drug target. Although methods for expression and purification of human NADK have been previously established (F. Lerner, M. Niere, A. Ludwig, M. Ziegler, Biochem. Biophys. Res. Commun. 288 (2001) 69-74), the purification procedure could be significantly improved. In this study, we improved the method for expression and purification of human NADK in *Escherichia coli* and obtained a purified homogeneous enzyme only through heat treatment and single column chromatography. Using the purified human NADK, we revealed a sigmoidal kinetic behavior toward ATP and the inhibitory effects of NADPH and NADH, but not of NADP⁺, on the catalytic activity of the enzyme. These inhibitory effects provide insight into the regulation of intracellular NADPH synthesis. Furthermore, these attributes may provide a clue to design a novel drug against *Mycobacterium tuberculosis* in which this bacterial NADK is potently inhibited by NADP⁺.

Keywords

NAD kinase; human; drug design; NADP⁺; NADPH; NADH

Abbreviations

NADK, NAD kinase; LB, Luria-Bertani; SDS-PAGE, sodium dodecyl sulfate-polyacrylamide gel electrophoresis

1 **Introduction**

2
3 NAD^+ and NADP^+ are cofactors involved in oxidative–reductive reactions [1, 2].
4 The function of NAD(H) (NAD^+ and NADH) is distinguishable from that of
5 NADP(H) (NADP^+ and NADPH). NAD(H) mainly functions in catabolic reactions,
6 while NADP(H) is involved in anabolic reactions and defense against oxidative
7 stress [3]. NAD^+ also functions as a substrate for mono- and poly-ADP-ribosylation
8 and is involved in cyclic ADP-ribose synthesis [2] and histone deacetylation [4].
9 NADP^+ functions as a substrate in the synthesis of nicotinic acid adenine
10 dinucleotide phosphate, which is a potent intracellular Ca^{2+} -mobilizing messenger [1,
11 2, 5].

12 NAD kinase (NADK, EC 2.7.1.23) is the sole NADP^+ -biosynthetic enzyme that
13 catalyzes phosphorylation of NAD^+ to yield NADP^+ using ATP as a phosphoryl
14 donor [1], and thus, plays a vital role in the cell. Accordingly, NADK has been found
15 to be essential for many microbes, including *Mycobacterium tuberculosis* [6],
16 *Bacillus subtilis* [7], *Staphylococcus aureus* [8], *Haemophilus influenzae* [8],
17 *Escherichia coli* [8, 9], *Streptococcus pneumoniae* [8], *Salmonella enterica* [9, 10],
18 and *Saccharomyces cerevisiae* [11, 12], thus representing a potentially powerful
19 antimicrobial drug target [13, 14]. Several bacterial recombinant NADKs have been
20 characterized in detail, including those from *E. coli* [15], *B. subtilis* [16], and *M.*
21 *tuberculosis* [17, 18]. Moreover, tertiary structures of NADKs from the pathogenic
22 bacteria *M. tuberculosis* [19, 20] and *Listeria monocytogenes* [21] have been
23 published. However, the tertiary structure of human NADK has not yet been
24 reported.

Lerner *et al.* [22] established methods for expression and purification of human NADK and determined some properties of this enzyme [22]. Although they provided a detailed purification procedure, the procedure still has needed several steps such as heat treatment followed by DEAE column chromatography and nickel–nitrilotriacetic acid column chromatography. Thus, an improvement in the methods for expression and purification of human NADK would facilitate further study of this vital enzyme. Further determination of the structure and catalytic properties of human NADK would provide a good foundation for designing a novel antimicrobial drug [13].

In this article, we describe improved methods to express and purify human NADK, thus enabling to obtain larger amounts of the purified homogeneous enzyme. In addition, we reveal some kinetic properties of human NADK that will provide insight into the regulation of intracellular NADPH synthesis and possibly a clue to design a novel antimicrobial drug.

Materials and methods

Strains and plasmids

As a host for plasmid amplification, *E. coli* DH5 α and JM109 were routinely cultured at 37 °C in Luria-Bertani (LB) medium containing 1% tryptone, 0.5% yeast extract, 1% NaCl, and ampicillin (100 μ g/ml) (pH 7.2). *E. coli* M15 (pREP4) and SG13009 (pREP4) were purchased from Qiagen (Hilden, Germany), and *E. coli* Rosetta-gami, Rosetta-gami(DE3), Rosetta-gami(DE3)pLysS, Rosetta-gami 2(DE3),

1 and Rosetta-gami B(DE3) were obtained from Novagen (Darmstadt, Germany). The
2 culture conditions for these *E. coli* strains are described below.

3 Human NADK cDNA was amplified by PCR using human leukocyte cDNA
4 (QUICK-Clone cDNA, Clontech, shiga, Japan) as a template and primers
5 [HsNADKBamHI-f, CGGGATCCATGGAAATGGAACAAGAAAAAATGAC (the
6 *Bam*HI site is underlined) and HsNADK-C(NoRe)-r,
7 CTAGCCCTCCTCCTCCTCCTC], and was inserted into the *Hinc*II site of pUC18,
8 resulting in pMK1966. Human NADK cDNA obtained by digestion of pMK1966
9 with *Bam*HI was inserted into the *Bam*HI site of pQE-30 (Qiagen), yielding
10 pMK2071. The primary structure encoded by the cloned cDNA was identical to that
11 of the cDNA sequence (with Phe (Glu)₉ Gly as C-terminus) deposited in the NCBI
12 database (accession no. NP_001185922), except that the former contains one
13 additional Glu residue at its C-terminus, i.e., it has Phe (Glu)₁₀ Gly.

14 A DNA fragment encoding for an N-terminal sequence consisting of 87 amino
15 acid residues of human NADK was excised from the full-length cDNA clone
16 pMK2071, resulting in pMK2784 that encodes for deleted human NADK with an
17 N-terminal sequence, ⁸⁸MHIQDPASQRL⁹⁸. This deletion was generated by inverse
18 PCR using primers (5'-CACATTCAGGACCCCGCGAG-3' and 5'-
19 CATATGTCCGTGATGGTGGTGGTGG-3'). The recombinant proteins encoded by
20 pMK2071 and pMK2784 contained a His-tag (MRGSHHHHHHGS) at their
21 N-terminal sites.

22 To express human NADK cDNA, pMK2071 was introduced into *E. coli* M15
23 (pREP4), SG13009 (pREP4), and Rosetta-gami, yielding MK2105, MK2106, and
24 MK2107, respectively. pMK2784 was introduced into *E. coli* Rosetta-gami(DE3),

Rosetta-gami(DE3)pLysS, Rosetta-gami 2(DE3), and Rosetta-gami B(DE3), yielding MK2801, MK2802, MK2803, and MK2804, respectively.

Expressions

The cells were precultured overnight in LB media supplemented with 100 µg/ml ampicillin and 25 µg/ml kanamycin (MK2105 and MK2106) or with 100 µg/ml ampicillin and 34 µg/ml chloramphenicol (MK2107, MK2801, MK2802, MK2803, and MK2804). The cells were collected, inoculated in LB media supplemented with the same antibiotics, and aerobically cultured at 37 °C until A_{600} reached 0.50–1.0. Isopropyl-β-D-thiogalactopyranoside was then added to obtain a final concentration of 0.025 or 1.0 mM in order to induce expression of human NADK. Cell cultivation was continued further at 16 or 37 °C under aerobic conditions for 4, 8, 12, or 24 h. To purify human NADK, MK2107 and MK2802 were cultivated at 37 °C for 8 h (MK2107) or 12 h (MK2802) after the addition of isopropyl-β-D-thiogalactopyranoside at 1.0 mM. To prepare the crude extract, cells were collected by centrifugation, suspended in 10 mM Tris-HCl (pH8.0), and sonicated to yield cell lysate. Before and after sonication, phenylmethylsulfonyl fluoride was added to obtain a final concentration of 1.0 mM. After centrifugation of the lysate at 20,000 ×g and 4 °C for 10 min, the supernatant was used as crude extract.

1 Assays

2

3 NADK activity of human NADK was assayed at 37 °C by the modified
4 continuous method [23]. Briefly, NADPH formation was continuously measured at
5 A_{340} in a reaction mixture (1.0 ml) containing 5.0 mM NAD^+ , 5.0 mM ATP, 5.0 mM
6 glucose-6-phosphate, 0.5 U glucose-6-phosphate dehydrogenase (Sigma, St. Louis,
7 MD, USA), 5.0 mM MgCl_2 , and 100 mM Tris-HCl (pH 8.0). In the purified enzyme
8 assay, MgCl_2 was used at 10 mM. Enzyme activity was assayed by a stop method
9 [23] to determine inhibitory effects and optimum pH and temperature conditions.
10 Briefly, the amount of NADP^+ formed was enzymatically determined using 0.5 U
11 glucose-6-phosphate dehydrogenase after the reaction was terminated by immersing
12 the test tube in boiling water for 5 min. Glucose-6-phosphate and
13 glucose-6-phosphate dehydrogenase were removed from the reaction mixture as
14 described above. When the inhibitory effects of NADP(H) and NADH were
15 investigated, control reactions without human NADK were simultaneously
16 performed in every assay.

17 NADH kinase activity was assayed in a reaction mixture (1.0 ml) containing 2.0
18 mM NADH, 5.0 mM ATP, 10.0 mM MgCl_2 , and 100 mM Tris-HCl (pH 8.0) [23].
19 One unit of enzyme activity was defined as 1.0 μmol NADP(H) produced in 1 min at
20 37 °C, and specific activity was expressed in U/mg protein. Protein concentrations of
21 the crude extract and purified human NADK from MK2107 were determined by the
22 Bradford method [24]. Protein concentrations of purified human NADK from
23 MK2802 were determined using the extinction coefficient (33,920) at A_{280} . The
24 extinction coefficients were calculated using the ProtParam tool on the ExPASy

server (<http://ca.expasy.org/tools/protparam.html>).

Purification

Centrifugation was performed at $20,000 \times g$ and 4°C for 10 min. Human NADK was expressed from MK2107 cells cultured in 2 l LB (0.5 l per 2 l Erlenmeyer flask) as described above and treated as described [22]. This was followed by Mono Q chromatography. The partially purified human NADK was applied to Mono Q 5/50 GL (0.5×5.0 cm) (GE Healthcare, Buckinghamshire, England) equilibrated with buffer A (20 mM Tris-HCl (pH 8.5), 0.10 mM NAD^+ , 0.50 mM dithiothreitol, 1.0 mM ethylenediaminetetraacetic acid, and 0.10 mM pepstatin A). After washing the column with 18 ml buffer A, elution was performed with a linear gradient of NaCl from 0 to 1.0 M in buffer A. Fractions that were obtained by an elution at 370–470 mM NaCl were combined, dialyzed against buffer A, and used as purified human NADK.

Human NADK was purified from MK2802 as stated below. Crude extract was prepared from MK2802 cells cultured in 13.5 l LB (1.5 l per 2 l Sakaguchi flask) and was treated as described above. Samples were heated at 60°C for 5 min and centrifuged. The supernatant was applied to a TALON column (1.5×22 cm) (Clontech) equilibrated with 10 mM Tris-HCl (pH8.0). The column was washed with 2 l of 10 mM Tris-HCl (pH8.0) containing 0.30 M NaCl. Human NADK was eluted with 10 mM Tris-HCl (pH8.0) containing 0.30 M NaCl and 150 mM imidazole. Fractions were combined, dialyzed against 10 mM Tris-HCl (pH 8.0), and used as purified human NADK. Alternatively, the crude extract prepared from MK2802 cells

1 was treated as mentioned above and was directly applied to the TALON column
2 without heat treatment. The column was washed with 2 l of 10 mM Tris-HCl (pH8.0)
3 containing 0.30 M NaCl and 30 mM imidazole, and eluted as above.

4 5 Other analytical methods

6
7 Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) was
8 performed using a 12.5% polyacrylamide gel [25]. Proteins in the gel were visualized
9 with Coomassie brilliant blue R-250. The molecular mass of the enzyme was
10 estimated by gel filtration chromatography as described [26]; however, 10 mM
11 Tris-HCl (pH 8.0) containing 0.15 M NaCl was used as an elution buffer. For the
12 N-terminal amino acid sequence, purified and unpurified enzymes were transferred
13 to a polyvinylidene difluoride membrane and analyzed using a Procise 492 protein
14 sequencing system (Applied Biosystems Division of Perkin-Elmer, Foster City, CA,
15 USA).

16 17 **Results and discussion**

18 19 Improved methods to express and purify human NADK

20
21 According to the method described by Lerner *et al.* [22], we expressed the
22 full-length human NADK cDNA cloned into pQE-30 (pMK2071) using an *E. coli*
23 M15 (pREP4) host cell and obtained 0.23 U/mg NADK activity in the crude extract
24 (Table 1). To improve expression, effects of the host cells (*E. coli* SG13009 (pREP4)

1 or Rosetta-gami), concentrations of isopropyl- β -D-thiogalactopyranoside (0.025 or
2 1.0 mM), and induction conditions (16 or 37 °C and 4, 8, 12, or 24 h) were examined.
3 Maximum expression (1.02 U/mg) was obtained in the crude extract when
4 expression was induced at 37 °C for 8 h with 1.0 mM
5 isopropyl- β -D-thiogalactopyranoside using *E. coli* Rosetta-gami (Table 1). The
6 expressed human NADK was treated according to the purification procedure
7 described by Lerner *et al.* [22]. However, this procedure yielded an unpurified
8 enzyme. The unpurified enzyme was further purified by Mono Q chromatography as
9 described in Materials and methods. The purified enzyme (0.39 mg; 7.21 U/mg) was
10 obtained from a 2 l culture. SDS-PAGE analysis showed that the subunit molecular
11 mass of the purified enzyme was 43 kDa, which was smaller than that predicted (49
12 kDa) from the primary structure (data not shown). Analysis of the N-terminal amino
13 acid sequence of the purified enzyme confirmed that the enzyme lacked the
14 N-terminal sequences consisting of 60, 63, and 74 amino acid residues of human
15 NADK, indicating the heterogeneity of the purified enzyme.

16 To circumvent this problem, the DNA region that encoded the N-terminal
17 sequence consisting of 87 amino acid residues of human NADK was excised from
18 the full-length human NADK cDNA as described in Materials and methods. The
19 deleted human NADK was expressed in *E. coli* Rosetta-gami(DE3), and it showed a
20 specific activity of 0.79 U/mg (Table 1). Further optimization of the expression
21 condition revealed that induction at 37 °C for 8–12 h in the presence of 1.0 mM
22 isopropyl- β -D-thiogalactopyranoside using *E. coli* Rosetta-gami(DE3)pLysS as a
23 host cell gave the maximum specific activity (4.29 U/mg). This activity was
24 improved by approximately 18.7-fold compared with our initial attempt (0.23 U/mg;

Table 1). Moreover, we found that the heat treatment, followed by TALON column chromatography was sufficient to obtain the purified enzyme (9.34 U/mg, Fig. 1a). The subunit molecular mass of the purified enzyme (43 kDa) was in good agreement with the predicted mass (41 kDa). The N-terminal amino acid sequence of the enzyme was RGSHH, which corresponded to that of the His-tag, confirming the homogeneity of the purified enzyme. In all purification experiments, the purified enzyme (15 mg) was routinely obtained from a 13.5 l culture. Thus, we improved the methods for expression and purification of human NADK (Table 2, Fig. 1a).

We also found that the enzyme could be purified only through TALON column chromatography without heat treatment (Fig. 1b). However, the purified NADK without heat treatment yielded three peaks (representing 210, 400, and 650 kDa) on gel filtration chromatography; each peak showed NADK activity. NADK in each peak was not divided into three peaks on repeated gel filtration chromatography (data not shown), thus indicating the heterogeneity in the quaternary structure of this human NADK. In contrast, the purified enzyme obtained by heat treatment, which was followed by TALON column chromatography (Fig. 1a), yielded a single peak representing 172 kDa (data not shown). Thus, the procedure using heat treatment was considered preferable for enzyme purification, and the properties of human NADK purified by this procedure were determined.

Properties of human NADK

Although the properties of purified human NADK were similar to those reported [22] (Table 3), we found that human NADK exhibits sigmoidal kinetic behavior

1 toward ATP, while saturation curve toward NAD^+ (Fig. 2). The sigmoidal kinetic
2 behavior was emphasized by a Hanes–Woolf plot (Fig. 2b). The sigmoidal kinetic
3 behavior toward ATP was also observed in NADKs from *B. subtilis* [16] and *M.*
4 *tuberculosis* [18]. With regard to NAD^+ , *M. tuberculosis* NADK shows sigmoidal
5 kinetic behavior [18], whereas *B. subtilis* NADK exhibits a saturation curve [16].

6 NADH-phosphorylating (NADH kinase) activity of recombinant human NADK
7 was qualitatively, but not quantitatively, demonstrated by Pollak *et al.* [27]. We
8 showed that human NADK exhibited NADH kinase activity (0.6 U/mg), accounting
9 for only 5.0% of NADK activity (11.9 U/mg) determined in the presence of 2.0 mM
10 NAD^+ . Some bacterial NADKs, such as those from *M. tuberculosis*, utilize inorganic
11 polyphosphate as a phosphoryl donor [17, 18]. Inorganic polyphosphate is a polymer
12 of inorganic orthophosphate residues linked by a high-energy phosphoanhydride
13 bond [28], and is available as metaphosphate, hexametaphosphate, and
14 tetrapolyphosphate. The relative activities of purified human NADK in the presence
15 of 1 mg/ml metaphosphate, 1 mg/ml hexametaphosphate, and 5 mM
16 tetrapolyphosphate were 2.3, 1.9, and 0.5%, respectively, compared with NADK
17 activity assayed in the presence of 5.0 mM ATP.

18
19 Inhibitory effects of NADPH and NADH, but not of NADP^+

20
21 Determining how human NADK is regulated by NADP^+ , NADPH, and NADH
22 is a crucial step in understanding how human NADK influences the intracellular
23 redox balance. Although NADP^+ exhibited no inhibitory effect on NADK activity of
24 human NADK, NADPH and NADH showed competitive inhibitions with K_i values

1 of 0.13 mM for NADPH and 0.34 mM for NADH (Table 4, Fig. 3, Online Resource
2 Fig. S1). This agrees with a previous report, which shows that NADK activity of
3 partially purified pigeon liver NADK is competitively inhibited by NADPH and
4 NADH, while the effect of NADP^+ on NADK activity of the enzyme was not
5 investigated [29]. Although the information regarding the effect of these nucleotides
6 on NADK activity has been limited, *M. tuberculosis* NADK has been reported to be
7 inhibited by NADH, NADP^+ , and NADPH, wherein NADP^+ is the most potent
8 inhibitor [18]. The residual NADK activity of *M. tuberculosis* NADK was 22, 43,
9 and 50% in the presence of 0.40 mM NADP^+ , NADPH, and NADH, respectively,
10 although the inhibitory mechanism has not been investigated [18]. The fact that
11 NADP^+ inhibits *M. tuberculosis* NADK but not human NADK is a clue to design a
12 novel antimicrobial drug, i.e., an analog of NADP^+ , which would possibly inhibit
13 only *M. tuberculosis* NADK, thereby being lethal to the bacterium and not the host.

14 Recently, Pollak *et al.* [27] established human cell lines overexpressing human
15 NADK (NADK(+) cells). In NADK(+) cells that showed approximately 180-fold
16 higher NADK activity than control cells, the NADPH level increased by 4- to 5-fold,
17 and not 180-fold, while the NADP^+ level was not prominently altered. The authors
18 suggested that human NADK may be inhibited by NADPH. Our data demonstrating
19 the inhibition of human NADK by NADPH convincingly support their suggestion.
20 Taken together the low NADH kinase activity of human NADK, NADPH would be
21 produced by NADP^+ -dependent dehydrogenases through reduction of NADP^+ . The
22 fact that human NADK is localized in the cytosol [27] and that NADPH, but not
23 NADP^+ , is significant for the regulation of human NADK would indicate that the
24 function of cytosolic NADP^+ -dependent dehydrogenases is important for the

1 regulation of NADP⁺ synthesis (NADK activity), although the physiological role of
2 inhibition of human NADK by NADH remains unclear.

3 NADPH is required to protect cells against oxidative stress, and cytosolic
4 NADP⁺-dependent glucose-6-phosphate dehydrogenase in human cells is induced
5 when exposed to oxidative stress [30]. When cells survive against oxidative stress,
6 intracellular NADPH is consumed and resynthesized quickly, while human NADK
7 synthesizes NADP⁺ to guarantee the intracellular requirement for NADPH. When
8 oxidative stress is removed, the abundant NADPH inhibits NADP⁺ synthesis to
9 prevent accumulation of excess NADP⁺ and NADPH, thus resulting in the
10 maintenance of homeostasis of intracellular NADP⁺ and NADPH.

12 **Acknowledgments**

13 This work was supported in part by a Grant-in-Aid for SK from the Ministry of
14 Education, Culture, Sports, Science and Technology of Japan (21780069).

16 **References**

- 18 1. Pollak N, Dolle C, Ziegler M (2007) The power to reduce: pyridine nucleotides -
19 small molecules with a multitude of functions. *Biochem J* 402 (2):205-218
- 20 2. Ziegler M (2000) New functions of a long-known molecule. Emerging roles of
21 NAD in cellular signaling. *Eur J Biochem* 267 (6):1550-1564
- 22 3. Outten CE, Culotta VC (2003) A novel NADH kinase is the mitochondrial source
23 of NADPH in *Saccharomyces cerevisiae*. *EMBO J* 22 (9):2015-2024
- 24 4. Smith JS, Brachmann CB, Celic I, Kenna MA, Muhammad S, Starai VJ, Avalos

1 JL, Escalante-Semerena JC, Grubmeyer C, Wolberger C, Boeke JD (2000) A
2 phylogenetically conserved NAD⁺-dependent protein deacetylase activity in the
3 Sir2 protein family. Proc Natl Acad Sci USA 97 (12):6658-6663

4 5. Yamasaki M, Masgrau R, Morgan AJ, Churchill GC, Patel S, Ashcroft SJ,
5 Galione A (2004) Organelle selection determines agonist-specific Ca²⁺ signals in
6 pancreatic acinar and beta cells. J Biol Chem 279 (8):7234-7240

7 6. Sassetti CM, Boyd DH, Rubin EJ (2003) Genes required for mycobacterial
8 growth defined by high density mutagenesis. Mol Microbiol 48 (1):77-84

9 7. Kobayashi K, Ehrlich SD, Albertini A, Amati G, Andersen KK, Arnaud M, Asai
10 K, Ashikaga S, Aymerich S, Bessieres P, Boland F, Brignell SC, Bron S, Bunai K,
11 Chapuis J, Christiansen LC, Danchin A, Debarbouille M, Dervyn E, Deuerling E,
12 Devine K, Devine SK, Dreesen O, Errington J, Fillinger S, Foster SJ, Fujita Y,
13 Galizzi A, Gardan R, Eschevins C, Fukushima T, Haga K, Harwood CR, Hecker
14 M, Hosoya D, Hullo MF, Kakeshita H, Karamata D, Kasahara Y, Kawamura F,
15 Koga K, Koski P, Kuwana R, Imamura D, Ishimaru M, Ishikawa S, Ishio I, Le
16 Coq D, Masson A, Mauel C, Meima R, Mellado RP, Moir A, Moriya S,
17 Nagakawa E, Nanamiya H, Nakai S, Nygaard P, Ogura M, Ohanan T, O'Reilly M,
18 O'Rourke M, Pragai Z, Pooley HM, Rapoport G, Rawlins JP, Rivas LA, Rivolta C,
19 Sadaie A, Sadaie Y, Sarvas M, Sato T, Saxild HH, Scanlan E, Schumann W,
20 Seegers JF, Sekiguchi J, Sekowska A, Seror SJ, Simon M, Stragier P, Studer R,
21 Takamatsu H, Tanaka T, Takeuchi M, Thomaides HB, Vagner V, van Dijl JM,
22 Watabe K, Wipat A, Yamamoto H, Yamamoto M, Yamamoto Y, Yamane K, Yata
23 K, Yoshida K, Yoshikawa H, Zuber U, Ogasawara N (2003) Essential *Bacillus*
24 *subtilis* genes. Proc Natl Acad Sci USA 100 (8):4678-4683

- 1 8. Zalacain M, Biswas S, Ingraham KA, Ambrad J, Bryant A, Chalker AF,
2 Iordanescu S, Fan J, Fan F, Lunsford RD, O'Dwyer K, Palmer LM, So C,
3 Sylvester D, Volker C, Warren P, McDevitt D, Brown JR, Holmes DJ, Burnham
4 MK (2003) A global approach to identify novel broad-spectrum antibacterial
5 targets among proteins of unknown function. *J Mol Microbiol Biotechnol* 6
6 (2):109-126
- 7 9. Gerdes SY, Scholle MD, D'Souza M, Bernal A, Baev MV, Farrell M, Kurnasov
8 OV, Daugherty MD, Mseeh F, Polanuyer BM, Campbell JW, Anantha S, Shatalin
9 KY, Chowdhury SA, Fonstein MY, Osterman AL (2002) From genetic
10 footprinting to antimicrobial drug targets: examples in cofactor biosynthetic
11 pathways. *J Bacteriol* 184 (16):4555-4572
- 12 10. Grose JH, Joss L, Velick SF, Roth JR (2006) Evidence that feedback inhibition of
13 NAD kinase controls responses to oxidative stress. *Proc Natl Acad Sci USA* 103
14 (20):7601-7606
- 15 11. Miyagi H, Kawai S, Murata K (2009) Two sources of mitochondrial NADPH in
16 the yeast *Saccharomyces cerevisiae*. *J Biol Chem* 284 (12):7553-7560
- 17 12. Bieganski P, Seidle HF, Wojcik M, Brenner C (2006) Synthetic lethal and
18 biochemical analyses of NAD and NADH kinases in *Saccharomyces cerevisiae*
19 establish separation of cellular functions. *J Biol Chem* 281 (32):22439-22445
- 20 13. Magni G, Di Stefano M, Orsomando G, Raffaelli N, Ruggieri S (2009) NAD(P)
21 biosynthesis enzymes as potential targets for selective drug design. *Curr Med*
22 *Chem* 16 (11):1372-1390
- 23 14. Bi J, Wang H, Xie J (2011) Comparative genomics of NAD(P) biosynthesis and
24 novel antibiotic drug targets. *J Cell Physiol* 226 (2):331-340.

1 doi:10.1002/jcp.22419

2 15. Kawai S, Mori S, Mukai T, Hashimoto W, Murata K (2001) Molecular
3 characterization of *Escherichia coli* NAD kinase. Eur J Biochem 268
4 (15):4359-4365

5 16. Garavaglia S, Galizzi A, Rizzi M (2003) Allosteric regulation of *Bacillus subtilis*
6 NAD kinase by quinolinic acid. J Bacteriol 185 (16):4844-4850

7 17. Kawai S, Mori S, Mukai T, Suzuki S, Yamada T, Hashimoto W, Murata K (2000)
8 Inorganic polyphosphate/ATP-NAD kinase of *Micrococcus flavus* and
9 *Mycobacterium tuberculosis* H37Rv. Biochem Biophys Res Commun 276
10 (1):57-63

11 18. Raffaelli N, Finaurini L, Mazzola F, Pucci L, Sorci L, Amici A, Magni G (2004)
12 Characterization of *Mycobacterium tuberculosis* NAD kinase: functional
13 analysis of the full-length enzyme by site-directed mutagenesis. Biochemistry 43
14 (23):7610-7617

15 19. Mori S, Yamasaki M, Maruyama Y, Momma K, Kawai S, Hashimoto W, Mikami
16 B, Murata K (2005) NAD-binding mode and the significance of intersubunit
17 contact revealed by the crystal structure of *Mycobacterium tuberculosis* NAD
18 kinase-NAD complex. Biochem Biophys Res Commun 327 (2):500-508

19 20. Garavaglia S, Raffaelli N, Finaurini L, Magni G, Rizzi M (2004) A novel fold
20 revealed by *Mycobacterium tuberculosis* NAD kinase, a key allosteric enzyme in
21 NADP biosynthesis. J Biol Chem 279 (39):40980-40986

22 21. Poncet-Montange G, Assairi L, Arold S, Pochet S, Labesse G (2007) NAD
23 kinases use substrate-assisted catalysis for specific recognition of NAD. J Biol
24 Chem 282 (47):33925-33934

- 1 22. Lerner F, Niere M, Ludwig A, Ziegler M (2001) Structural and functional
2 characterization of human NAD kinase. *Biochem Biophys Res Commun* 288
3 (1):69-74
- 4 23. Shi F, Kawai S, Mori S, Kono E, Murata K (2005) Identification of ATP-NADH
5 kinase isozymes and their contribution to supply of NADP(H) in *Saccharomyces*
6 *cerevisiae*. *FEBS J* 272 (13):3337-3349
- 7 24. Bradford MM (1976) A rapid and sensitive method for the quantitation of
8 microgram quantities of protein utilizing the principle of protein-dye binding.
9 *Anal Biochem* 72:248-254
- 10 25. Laemmli UK (1970) Cleavage of structural proteins during the assembly of the
11 head of bacteriophage T4. *Nature* 227 (5259):680-685
- 12 26. Ochiai A, Mori S, Kawai S, Murata K (2004) Overexpression, purification, and
13 characterization of ATP-NAD kinase of *Sphingomonas* sp. A1. *Protein Expr*
14 *Purif* 36 (1):124-130
- 15 27. Pollak N, Niere M, Ziegler M (2007) NAD kinase levels control the NADPH
16 concentration in human cells. *J Biol Chem* 282 (46):33562-33571
- 17 28. Kornberg A, Rao NN, Ault-Riche D (1999) Inorganic polyphosphate: a molecule
18 of many functions. *Annu Rev Biochem* 68:89-125
- 19 29. Apps DK (1968) Kinetic studies of pigeon liver NAD kinase. *Eur J Biochem* 5
20 (3):444-450
- 21 30. Ursini MV, Parrella A, Rosa G, Salzano S, Martini G (1997) Enhanced expression
22 of glucose-6-phosphate dehydrogenase in human cells sustaining oxidative
23 stress. *Biochem J* 323 (Pt 3):801-806
- 24

Figure legends

Fig. 1. SDS-PAGE of purified human NADK. (a) Purified human NADK (110 μ g protein). (b) Purified human NADK obtained without heat treatment (60 μ g protein). Arrowheads indicate the positions of the purified human NADK.

Fig. 2. Kinetic analysis of human NADK. (a) The effect of substrate concentration ([ATP] or [NAD⁺], mM) on the initial velocity (V, U/mg) of the reaction was assayed in the presence of 5 mM NAD⁺ and various concentrations of ATP (closed squares) or 5 mM ATP and various concentrations of NAD⁺ (open squares). Data represent the means of three measurements. (b) Hanes–Woelf plot of the data in (a).

Fig. 3. Inhibitory effects of NADH (a, b) and NADPH (c, d) on human NADK. The initial velocity (V, U/mg) of the reaction was assayed in the presence of various concentrations of NAD⁺ ([NAD⁺], mM). (a) Saturation curve for NAD⁺ determined in the presence of 0 mM (closed squares), 0.3 mM (open circles), and 0.5 mM (closed circles) NADH. (b) Lineweaver–Burk plot of the data in (a). (c) Saturation curve for NAD⁺ determined in the presence of 0 mM (closed squares) and 0.5 mM (open squares) NADPH. (d) Lineweaver–Burk plot of the data in (c).

Online Resource figure legend

Online Resource Fig. S1. Inhibitory effects of NADH (a, b) and NADPH (c, d) on human NADK. (a) Hanes–Woelf plot of the data in Fig. 3a. (b) Eadie plot of the data

1 in Fig. 3a. The plot was determined in the presence of 0 mM (closed squares), 0.3
2 mM (open circles), and 0.5 mM (closed circles) NADH. (c) Hanes–Woelf plot of the
3 data in Fig. 3c. (d) Eadie plot of the data in Fig. 3c. The plot was determined in the
4 presence of 0 mM (closed squares) and 0.5 mM (open squares) NADPH.
5

1

2

Table 1

Expression of human NADK.

Host cell	Specific activity (U/mg) ^a	Relative activity (fold)
M15 (pREP4)	0.23	1
Rosetta-gami	1.02	4.4
Rosetta-gami(DE3)	0.79	3.4
Rosetta-gami(DE3)pLysS	4.29	18.7

^aSpecific activity in crude extract.

3

4

5

6

7

8

9

10

11

12

13

14

15

1

2

Table 2

Improvement in the methods for expression and purification of human NADK.

	Previous study [22]	Present study
Specific activity of expressed human NADK in crude extract	n.r. ^a (0.23 U/mg) ^b	4.29 U/mg
Purification procedure	Three steps (Heat treatment and DEAE as well as nickel–nitrilotriacetic acid column chromatography)	Two steps (Heat treatment and TALON column chromatography)
Yield of purified human NADK	n.r. ^a	15 mg ^c
Specific activity of purified human NADK	n.r. ^a	9.34 U/mg
Homogeneity of purified human NADK	n.r. ^{a, d}	Homogeneity ^d

^an.r.; not reported.^bThe value obtained in this study (Table 1).^cPurified human NADK (15 mg) was obtained from a 13.5 l culture.

^dN-terminal amino acid sequence of the purified human NADK was not determined in the previous report [22], while homogeneity of this enzyme was confirmed by analysis of the N-terminal amino acid sequence in this study.

3

4

Table 3

Properties of human NADK.

	Previous study [22]	Present study
Subunit molecular mass	49 kDa	43 kDa
Subunit composition (Molecular mass)	Homotetramer (200 kDa)	Homotetramer (172 kDa)
K_m (mM) for NAD^+	0.54	1.07
K_m (mM) for ATP	3.3	-
$S_{0.5}$ (mM) for ATP	-	3.6
n_H	-	1.6
V_{max} (U/mg) for NAD^+	6.7	18.5
V_{max} (U/mg) for ATP	-	18.5
Optimum temperature ($^{\circ}\text{C}$)	55	50
Optimum pH	7.5	8.0

1

2

3

4

5

6

7

8

Table 4

Inhibitory effects of NADP⁺, NADPH, and NADH on human NADK^a.

Compound	Concentration (mM)	Specific activity (U/mg) ^b	Relative activity (%)
None	0	8.75 ± 0.66	100
NADP ⁺	0.5	8.92 ± 2.26	102
NADPH	0.5	5.83 ± 0.49	66
NADPH	0.3	7.08 ± 0.42	81
NADH ^c	0.5	3.54 ± 0.48	40
NADH ^c	0.3	5.03 ± 0.61	58
NADH ^c	0.1	6.30 ± 0.36	72

^aNADK activity was assayed in the presence of 1.0 mM NAD⁺ and indicated compound.

^bMeans and standard deviations of three independent assays are indicated.

^cInhibitory effect of NADH was not attributed to NADH kinase activity of human NADK since the activity assayed in the presence of 0.3 or 0.5 mM NADH was only 10% of NADK activity assayed in the presence of 1.0 mM NAD⁺ (data not shown).

1

2

3

4

5

6

7





Fig. 1

1




2

3 Fig. 2

4

5

1




2

3 Fig. 3

4

5

1



2

Supplementary Fig. S1 Ohashi et al.

3