

On polynomial solutions of the Lamé and Stokes systems

Hiroya ITO*

The University of Electro-Communications

1 Introduction

The Laplacian Δ is one of the most important differential operators in Mathematics. Solutions of the Laplace equation $\Delta u = 0$ are called harmonic functions, which play significant roles in many subjects of mathematical research fields. It is well known that harmonic polynomials in n variables are well classified. Moreover, the restriction of nonzero elements of $\mathcal{H}_m$ to the unit sphere $\mathbb{S}^{n-1}$, called spherical harmonics of degree m , become eigenfunctions of the Laplace-Beltrami operator $-\Delta_{\mathbb{S}^{n-1}}$ on $\mathbb{S}^{n-1}$ with the common eigenvalue $m(m+n-2)$, and this restriction is a linear isomorphism between $\mathcal{H}_m$ and the space of spherical harmonics of degree m . Furthermore, we can think that all the harmonic polynomials in $\mathbb{R}^n$ (or all the spherical harmonics) generate most function spaces on $\mathbb{S}^{n-1}$ (see Theorem 1 below). We shall study that such beautiful theory can be partially generalized to vector-valued elliptic systems.

Fixing n variables $x_1, \dots, x_n$ with $n \geq 2$, for each $m \in \mathbb{N}_0$, we denote by $\mathcal{P}_m$ the vector space of all the homogeneous polynomials of degree m in $x = (x_1, \dots, x_n)$, and by $\mathcal{H}_m$ its subspace consisting of those in $\mathcal{P}_m$ which are harmonic. Moreover, $\mathring{\mathcal{H}}_m$ denotes the vector space of all the functions on $\mathbb{S}^{n-1}$ obtained by restricting each element of $\mathcal{H}_m$ to $\mathbb{S}^{n-1}$; each element of $\mathring{\mathcal{H}}_m$ is called a spherical harmonic of degree m :

$$\mathcal{H}_m = \{u \in \mathcal{P}_m \mid \Delta u = 0\}, \quad \mathring{\mathcal{H}}_m = \{u|_{\mathbb{S}^{n-1}} \mid u \in \mathcal{H}_m\}.$$

Then, the dimension d_m of $\mathcal{P}_m$ is given by $d_m = \binom{m+n-1}{n-1}$ and the restriction map $\mathcal{H}_m \ni u \mapsto u|_{\mathbb{S}^{n-1}} \in \mathring{\mathcal{H}}_m$ is, due to the homogeneity of elements of $\mathcal{H}_m$, a linear isomorphism: $\mathcal{H}_m \cong \mathring{\mathcal{H}}_m$. Fundamental properties of spherical harmonics on $\mathbb{S}^{n-1}$ are described in the following theorem (see, e.g., Chapter 2 of Shimakura [4], Chapter 3 of Simon [5], Nomura [2]).

Theorem 1. *The space $\mathring{\mathcal{H}}_m$ ($m \in \mathbb{N}_0$) has the following properties.*

- (i) *The dimension of $\mathring{\mathcal{H}}_m$ is given by $\dim \mathring{\mathcal{H}}_m = d_m - d_{m-2}$, where $d_{-1} = d_{-2} = 0$.*
- (ii) *$L^2(\mathbb{S}^{n-1}) = \bigoplus_{m=0}^{\infty} \mathring{\mathcal{H}}_m$ in the sense that*

$$\mathring{\mathcal{H}}_\ell \perp \mathring{\mathcal{H}}_m \ (\ell \neq m) \text{ in } L^2(\mathbb{S}^{n-1}) \quad \text{and} \quad \overline{\text{span}(\bigcup_{m=0}^{\infty} \mathring{\mathcal{H}}_m)}^{L^2} = L^2(\mathbb{S}^{n-1}).$$

In the present paper, we consider the homogeneous equation of the Lamé system

$$\mathcal{L}\mathbf{u} := \mu \Delta \mathbf{u} + (\lambda + \mu) \nabla(\text{div } \mathbf{u}) = \mathbf{0} \quad \text{in } \mathbb{R}^n \tag{1}$$

*e-mail: ito-hiroya@uec.ac.jp

for n -vector valued functions (vector fields) $\mathbf{u}$, where λ and μ are elasticity constants. We study the structure of the restriction of polynomial solutions of (1) to $\mathbb{S}^{n-1}$ (or analogues of spherical harmonics for (1)). Here, the operator $\mathcal{L}$ of (1) appears in linear theory of isotropic elasticity and the constants λ and μ are assumed to satisfy

$$\mu(\lambda + 2\mu) > 0, \quad \gamma := \frac{\lambda + \mu}{\lambda + 3\mu} \in (-1, 1). \quad (2)$$

The symbol of $\mathcal{L}$ is

$$L(\xi) = \mu|\xi|^2 I + (\lambda + \mu)\boldsymbol{\xi} \otimes \boldsymbol{\xi}, \quad (3)$$

whose eigenvalues are given by $\mu|\xi|^2$ (multiplicity $n-1$) and $(\lambda + 2\mu)|\xi|^2$ (simple), where we write ξ in boldface in order to clarify that $\boldsymbol{\xi}$ is a column vector. Assumption (2) implies that $\mathcal{L} = L(\partial)$ is a strongly elliptic system. In a similar way we also deal with polynomial solutions of the homogeneous equations of the Stokes system.

2 Orthogonally invariant partial differential operators for vector fields

Let $P(\partial)$ be a partial differential operator with constant coefficients for scalar fields u on $\mathbb{R}^n$. It is well-known that $P(\partial)$ is invariant under the special orthogonal group $\mathrm{SO}(n)$ if and only if $P(\partial)$ is in the form $P(\partial)u = f(\Delta)u$ for some polynomial $f(t)$. How about the case $P(\partial)$ is for vector fields $\mathbf{u}$ on $\mathbb{R}^n$? The following theorem In the case $P(\partial)$ for scalar functions u , it is well-known that $P(\partial)$ is invariant under $\mathrm{SO}(n)$ if and only if it is in the form $P(\partial)u = f(\Delta)u$. The following theorem shows, in a sense, the necessity of considering the Lamé system.

Theorem 2. *A partial differential operator $P(\partial)$ with constant coefficients for vector fields $\mathbf{u}$ on $\mathbb{R}^n$ is invariant under the orthogonal group $\mathrm{O}(n)$ if and only if $P(\partial)$ is in the form*

$$P(\partial)\mathbf{u} = f(\Delta)\mathbf{u} + g(\Delta)\nabla(\mathrm{div}\mathbf{u}).$$

for some polynomials $f(t)$ and $g(t)$.

Even if we restrict $\mathrm{O}(n)$ to $\mathrm{SO}(n)$ in Theorem 2, then the conclusion is valid for $n \geq 4$, but not for $n = 2, 3$, in which $P(\partial)\mathbf{u}$ may contain additional terms, for example $h(\Delta)\mathrm{rot}\mathbf{u}$ if $n = 3$.

3 L -harmonic vector fields and L -harmonics

Denote by $\mathcal{P}_m$ the vector space of all n -vector homogeneous polynomials in $x = (x_1, \dots, x_n)$ of degree m . We define subspaces $\mathcal{H}_m$ and $\mathcal{H}_m^L$ of $\mathcal{P}_m$ by

$$\mathcal{H}_m = \{\mathbf{u} \in \mathcal{P}_m \mid \Delta\mathbf{u} = \mathbf{0}\}, \quad \mathcal{H}_m^L = \{\mathbf{u} \in \mathcal{P}_m \mid \mathcal{L}\mathbf{u} = \mathbf{0}\}.$$

Elements of $\mathcal{H}_m^L$ are called *L -harmonic polynomials* of degree m .

Vector functions on $\mathbb{S}^{n-1}$ obtained by restricting L -harmonic polynomials are called *spherical L -harmonics*. We represent the vector spaces of such vector functions (vector fields) as

$$\hat{\mathcal{H}}_m = \{\mathbf{u}|_{\mathbb{S}^{n-1}} \mid \mathbf{u} \in \mathcal{H}_m\}, \quad \hat{\mathcal{H}}_m^L = \{\mathbf{u}|_{\mathbb{S}^{n-1}} \mid \mathbf{u} \in \mathcal{H}_m^L\}.$$

Corresponding to Theorem 1 for spherical harmonics, the following theorem for spherical L -harmonics has been established through joint research with Prof. Honda and Prof. Jimbo.

Theorem 3 ([1]). *The space $\mathring{\mathcal{H}}_m^L$ ($m \in \mathbb{N}_0$) has the following properties.*

- (i) *The dimension of $\mathring{\mathcal{H}}_m^L$ is given by $\dim \mathring{\mathcal{H}}_m^L = \dim \mathcal{H}_m^L = n(d_m - d_{m-2})$, where $d_{-1} = d_{-2} = 0$.*
- (ii) *For each $m \in \mathbb{N}$, the sum $\mathring{\mathcal{H}}_0^L + \mathring{\mathcal{H}}_1^L + \cdots + \mathring{\mathcal{H}}_m^L$ is a direct sum.*
- (iii) *The linear span of $\bigcup_{m=0}^{\infty} \mathring{\mathcal{H}}_m^L$ is dense in $L^2(\mathbb{S}^{n-1})$ with the L^2 -norm.*

4 Case $n = 2$

In this section we consider the case $n = 2$. The spaces $\mathcal{H}_m$, $\mathcal{H}_m^L$, $\mathring{\mathcal{H}}_m$, $\mathring{\mathcal{H}}_m^L$ are defined not only for nonnegative integer m but also negative integer m . For example, $\mathbf{u} \in \mathcal{H}_m^L$ for $m < 0$ implies that $\mathbf{u}$ is a vector field solution of (1) in $\mathbb{R}^2 \setminus \{0\}$ which is homogeneous in $x = (x_1, x_2)$ of degree m .

Let $\mathbf{u} = (u_1, u_2)$ be a real vector field solution of $\mathcal{L}\mathbf{u} = \mathbf{0}$ in $\mathbb{R}^2 \setminus \{0\}$. Then the complex function $U(z) := u_1(x_1, x_2) + iu_2(x_1, x_2)$ ($z := x_1 + ix_2$) satisfies

$$\frac{\partial}{\partial \bar{z}} \left(\frac{\partial U}{\partial z} + \gamma \frac{\bar{\partial} U}{\partial z} \right) = 0 \quad \text{in } \mathbb{C} \setminus \{0\},$$

which is solved as

$$U = \varphi(z) - \gamma z \overline{\varphi'(z)} + \overline{\psi(z)} + 2c \log|z| - \gamma \bar{c} \left(\frac{z}{|z|} \right)^2$$

where $\varphi(z), \psi(z)$ are holomorphic functions in $\mathbb{C} \setminus \{0\}$, and $c \in \mathbb{C}$ is a constant ([3]). Using this fact, we have the following assertions.

Theorem 4. *The spaces $\mathcal{H}_m^L$ and $\mathring{\mathcal{H}}_m^L$ have the following bases.*

- (i) $\mathcal{H}_0^L = \mathbb{R}^2$. For $m \neq 0$, the space $\mathcal{H}_m^L$ has a basis

$$\left\{ \begin{bmatrix} \operatorname{Re}[z^m - \gamma m z \overline{z^{m-1}}] \\ \operatorname{Im}[z^m - \gamma m z \overline{z^{m-1}}] \end{bmatrix}, \begin{bmatrix} -\operatorname{Im}[z^m + \gamma m z \overline{z^{m-1}}] \\ \operatorname{Re}[z^m + \gamma m z \overline{z^{m-1}}] \end{bmatrix}, \begin{bmatrix} \operatorname{Re}[z^m] \\ -\operatorname{Im}[z^m] \end{bmatrix}, \begin{bmatrix} \operatorname{Im}[z^m] \\ \operatorname{Re}[z^m] \end{bmatrix} \right\}.$$

- (ii) $\mathring{\mathcal{H}}_0^L = \mathbb{R}^2$. For $m \neq 0$, the space $\mathring{\mathcal{H}}_m^L$ has a basis

$$\left\{ \begin{bmatrix} \cos m\theta - \gamma m \cos(m-2)\theta \\ \sin m\theta + \gamma m \sin(m-2)\theta \end{bmatrix}, \begin{bmatrix} -\sin m\theta + \gamma m \sin(m-2)\theta \\ \cos m\theta + \gamma m \cos(m-2)\theta \end{bmatrix}, \begin{bmatrix} \cos m\theta \\ -\sin m\theta \end{bmatrix}, \begin{bmatrix} \sin m\theta \\ \cos m\theta \end{bmatrix} \right\}.$$

Corollary 5.

- (i) $\mathring{\mathcal{H}}_0^L + \mathring{\mathcal{H}}_1^L + \cdots + \mathring{\mathcal{H}}_m^L = \mathring{\mathcal{H}}_0 + \mathring{\mathcal{H}}_1 + \cdots + \mathring{\mathcal{H}}_m$ for $m \geq 1$.
- (ii) The sum $\mathring{\mathcal{H}}_0^L + \mathring{\mathcal{H}}_{-1}^L + \cdots + \mathring{\mathcal{H}}_{-m}^L$ ($m \geq 1$) is a direct sum, and satisfies

$$\mathring{\mathcal{H}}_0 + \mathring{\mathcal{H}}_1 + \cdots + \mathring{\mathcal{H}}_{m-2} \subset \mathring{\mathcal{H}}_0^L + \mathring{\mathcal{H}}_{-1}^L + \cdots + \mathring{\mathcal{H}}_{-m}^L \subset \mathring{\mathcal{H}}_0 + \mathring{\mathcal{H}}_1 + \cdots + \mathring{\mathcal{H}}_{m+2} \quad \text{for } m \geq 2.$$

References

- [1] N. Honda, H. Ito and S. Jimbo, Spherical vector functions associated with the Lamé and Stokes systems, *preprint*.
- [2] T. Nomura, Spherical Harmonics and Group Representations (*Japanese*), NipponHyoronsha (2018).

- [3] N. I. Muskhelishvili, Some Basic Problems of the Mathematical Theory of Elasticity, 2nd ed., Noordhoff N.V., 1953.
- [4] N. Shimakura, Partial Differential Operators of Elliptic Type, AMS, Providence, RI, 1992.
- [5] B. Simon, Harmonic analysis, A Comprehensive Course in Analysis, Part 3. AMS, Providence, RI, 2015.