

Drug monitoring for mycophenolic acid in graft-versus-host disease prophylaxis in cord blood transplantation

Hiroyuki Muranushi M.D¹, Junya Kanda M.D, Ph.D¹, Yasuyuki Arai M.D, Ph.D^{1, 2}, Takero Shindo M.D, Ph.D¹, Masakatsu Hishizawa M.D, Ph.D¹, Takashi Yamamoto Ph.D³, Tadakazu Kondo M.D, Ph.D¹, Kohei Yamashita M.D, Ph.D¹, Kazuo Matsubara Ph.D³, Akifumi Takaori-Kondo M.D, Ph.D¹

1. Department of Hematology and Oncology, Graduate School of Medicine, Kyoto University, Kyoto, Japan

2. Department of Transfusion Medicine and Cell Therapy, Kyoto University Hospital, Kyoto, Japan

3. Department of Clinical Pharmacology and Therapeutics, Kyoto University Hospital, Kyoto, Japan

Corresponding author:

Tadakazu Kondo, M.D, Ph.D.

Department of Hematology and Oncology, Graduate School of Medicine, Kyoto University, Kyoto, Japan

20 54, Shogoinkawahara-Cho, Sakyo-Ku, Kyoto City, 606-8507, Japan

21 Tel: +81-75-751-4964

22 Fax: +81-75-751-4963

23 E-mail: tadakazu@kuhp.kyoto-u.ac.jp

24 ‘The authors confirm that the Principal Investigator for this paper is Hiroyuki Muranushi
25 and that he had direct clinical responsibility for patients.

26 Running head: Drug monitoring for mycophenolic acid in CBT

27 Key words: Drug monitoring, mycophenolic acid, cord blood transplantation

28 Word count: 3073, Table count: 3, Figure count: 4, Supplemental figure count: 1

29 What is already known about this subject: Low level of mycophenolic acid after allogeneic
30 hematopoietic stem cell transplantation is associated with high incidence of acute
31 graft-versus-host disease.

32 What this study adds: Low level of mycophenolic acid after cord blood transplantation is
33 associated with high incidence of acute graft-versus-host disease and human herpesvirus 6
34 reactivation, whereas high level of mycophenolic acid is associated with high incidence of
35 sepsis before engraftment.

36

37

38

Abstract

Aims: We performed the retrospective analysis to clarify the significance of drug monitoring for mycophenolic acid (MPA), the active form of mycophenolate mofetil (MMF), in prophylaxis for graft-versus-host disease (GVHD) in cord blood transplantation (CBT).

Methods: We retrospectively analyzed the data of 46 patients who underwent first CBT and received GVHD prophylaxis with tacrolimus plus MMF. MPA levels were measured on days 7 and 21, and 24-hour areas under the curve (AUC_{0-24}) were estimated.

Results: The engraftment and 3-year overall survival rates of all patients were 94% and 78%, respectively. The cumulative incidence of sepsis before engraftment was higher in patients with AUC_{0-24} on day 7 of $> 60 \mu\text{g h/mL}$ than in other patients (33% vs. 6%, $p = 0.02$). The cumulative incidence of grade II–IV acute GVHD was higher in patients with AUC_{0-24} on day 21 of $\leq 30 \mu\text{g h/mL}$ than in other patients (80% vs. 50%, $p = 0.04$). The cumulative incidence of human herpesvirus 6 reactivation was higher in patients with AUC_{0-24} on day 21 of $\leq 48 \mu\text{g h/mL}$ (median) than in other patients (50% vs. 19%, $p = 0.03$).

Conclusions: Blood level of MPA was associated with the risks of acute GVHD and infection. A prospective trial evaluating the benefit of personalized MMF dosing using MPA levels is needed.

57

58 **Introduction**

59 Graft-versus-host disease (GVHD) is a major cause of morbidity and mortality after
60 allogeneic hematopoietic stem cell transplantation (allo-SCT). The combination of
61 calcineurin inhibitors (CNIs) with short-course methotrexate (MTX) after allo-SCT has been
62 standard GVHD prophylaxis for more than 20 years.[1, 2] Following the successful use of
63 mycophenolate mofetil (MMF) in the United States and Europe, the use of MMF with CNI in
64 cord blood transplantation (CBT) has recently increased in Japan to promote neutrophil
65 recovery.[3, 4] Wide interpatient variability in blood levels of mycophenolic acid (MPA), the
66 active form of MMF, has been reported even after administering the same dose of MMF.[5, 6]
67 In kidney transplantation, therapeutic monitoring of MPA is reported to be important, with
68 recommended target ranges for areas under the curve (AUCs) of 30–60 $\mu\text{g h/mL}$. AUCs <30
69 $\mu\text{g h/mL}$ are associated with transplanted kidney rejection, whereas AUCs >60 $\mu\text{g h/mL}$ are
70 associated with infection and myelosuppression.[6, 7] Similarly, there are some reports on
71 drug monitoring for MPA in bone marrow or peripheral blood stem cell transplantation.[5,
72 8-11] However, significance of drug monitoring for MPA in CBT was not fully
73 elucidated.[12] In 2015, we reported that a low blood AUC of MPA on day 21 was associated
74 with a high incidence of acute GVHD in the analysis of 24 patients who received CBT.[13]
75 Based on this previous result, we started to increase the dose of MMF in patients with low

MPA AUCs on day 7. In 2018, the number of patients increased sufficiently to perform univariate and multivariate analyses, and the observation times became long enough to evaluate the 3-year overall survival (OS). This retrospective analysis was performed to clarify the association between MPA AUCs and CBT outcomes.

Methods

Data collection

We retrospectively analyzed the data of 46 patients who underwent their first CBT and received GVHD prophylaxis with tacrolimus plus MMF in our institute between October 2011 and July 2016. Tacrolimus was continuously and intravenously administered from day -1. Blood levels of tacrolimus were measured at least three times per week and were maintained at 12–15 ng/mL. After neutrophil engraftment, tacrolimus was orally administered, with trough levels maintained at 5–10 ng/mL. The patients received 10 mg/kg of oral MMF every 8 hours from days -1 to 30 as inpatient treatment. Blood samples were collected immediately before and 1, 2, and 4 hours after the morning administration of MMF on days 7 and 21. Total MPA levels in the plasma were measured using the enzyme multiplied immunoassay technique (EMIT)[14, 15] or the Roche Total Mycophenolic Acid® assay[16] (C_0 , C_1 , C_2 , and C_4 , respectively), and C_8 (implying the trough level of the next administration) was assumed to be equal to the C_0 values based on the results of previous

studies.[9, 11, 17, 18] The 8-hour area under the curve (AUC_{0-8}) was determined using the linear trapezoidal method, and the 24-hour area under the curve (AUC_{0-24}) was calculated as three times the AUC_{0-8} . This method is frequently used in the setting of MMF administration three times daily (every 8 hours)[9, 18] because MPA levels peak within 2 hours from administration and decrease linearly after 4 hours.[17, 19] After 2014, the dose of MMF for patients with MPA AUC_{0-24} of $<30 \mu\text{g h/mL}$ on day 7 was increased. The dose of those patients from days 8 to 30 was calculated as follows: [the dose of MMF before day 7] $\times$ $40/[the AUC_{0-24} \text{ on day 7}]$. We analyzed the effect of MPA AUC_{0-24} on transplant outcomes. The cut-off values of AUC_{0-24} were set to 30 and $60 \mu\text{g h/mL}$, which were the same as the values used for kidney transplantation.[6] We also set the cut-off value as the median. To evaluate human herpesvirus 6 (HHV-6) and cytomegalovirus (CMV) reactivation, peripheral blood was collected at least once weekly until 90 days after transplantation. The HHV-6 DNA copy number was measured using real-time polymerase chain reaction methods, as described previously.[20] The present study was approved by the institutional ethics committee of the Graduate School of Medicine, Kyoto University. All patients provided consent to participate in the present study.

Definition

OS was defined as the time from transplantation to death, and patients who remained alive

at the time of the last follow-up were censored. Complete haematological remission (CR) was defined based on morphology for all the patients, except for those with lymphoma; for lymphoma patients, it was based on positron emission tomography-computed tomographic (PET-CT) imaging.[21, 22] The relapse date of patients who did not achieve CR before and after stem cell transplantation (SCT) was defined as day 1. The patients were divided into two groups according to the conditioning regimen: myeloablative conditioning (MAC) or reduced-intensity conditioning (RIC). MAC and RIC were defined as proposed by Giralt et al.[23] and Bacigalupo et al,[24] respectively. Pre-engraftment immune reaction (PIR) was characterized by the presence of at least three of the following symptoms with no direct consequences of infection or adverse effects of medication six or more days before engraftment, as described previously: a high fever ($>38.5^{\circ}\text{C}$), skin eruptions, body weight gain greater than 5% of baseline, or peripheral edema.[25, 26] HHV-6 reactivation was defined as plasma HHV-6 DNA level $\geq 10^4$ copies/mL. CMV reactivation was defined as the detection of at least two CMV pp65 antigen-positive cells per 50,000 leucocytes.

Data analysis

Descriptive statistics were used to summarize the variables related to patient demographics and transplantation characteristics. The probability of the OS time was estimated according to the Kaplan–Meier method.[27] To evaluate the influences of confounding factors on OS,

133 log-rank tests were used for the univariate analysis.[28] The competing risk regression model
134 was used for the multivariate analysis of acute GVHD.[29] The cumulative incidences of
135 engraftment, relapse, and GVHD were calculated, and death without events were considered
136 a competing risk. Landmark analysis was performed to evaluate the effect of MPA level on
137 days 7 and 21, with the landmark day set to days 7 and 21. Mann–Whitney *U* tests were used
138 as non-parametric tests for two groups. Correlation of AUC and single point concentration
139 was assessed by Spearman’s correlation coefficient test.[30] The results are expressed as
140 hazard ratios (HRs) and their 95% confidence intervals (CIs). All tests were two sided, and
141 *P*-values <0.05 were considered to indicate statistical significance. All statistical analyses
142 were performed using Stata (version 13.0, Stata Corporation) and EZR (Saitama Medical
143 Center, Jichi Medical University), a graphical user interface for R (The R Foundation for
144 Statistical Computing, version 2.3.0).[31]

Results

Patient characteristics

The median observation time of the survivors was 1,352 (range: 103–2,487) days. The median age was 50 (range: 20–66) years. The percentages of patients diagnosed with acute myeloid leukemia, acute lymphoid leukemia, myelodysplastic syndrome, and malignant lymphoma were 43%, 13%, 15%, and 17%, respectively. The percentage of patients with CR was 52%. Fifty-two percent of patients received MAC and 93% received total body irradiation (TBI) (Table 1).

Outcomes of all patients

The cumulative incidence of neutrophil engraftment was 94%. A median of 25 (range: 11–43) days was required for neutrophil engraftment (Figure 1A). Engraftment failure was seen in 3 patients, and they had risks for engraftment failure; one patient had high tumor burden, another patient received weak conditioning (fludarabine 125 mg/m² plus melphalan 80 mg/m² without TBI) and the other patient transplanted the minimum count of CD34⁺ cells (0.18×10⁵/kg) in this cohort. The cumulative incidences of PIR, grade II–IV acute GVHD, grade III–IV acute GVHD, limited chronic GVHD, and extensive chronic GVHD were 30%, 63%, 11%, 15%, and 13%, respectively. The 3-year OS, cumulative incidence of relapse, and non-relapse mortality rates were 78% (95% CI: 62–87%), 20% (95% CI: 10–33%), and 11%

(95% CI: 4–22%), respectively. After excluding two patients with aplastic anemia, we divided patients with respect to disease status at SCT. The 3-year OS, cumulative incidence of relapse, and non-relapse mortality rates of patients in CR (n = 24) were 82%, 22%, and 13%, (95% CI: 59–93%, 8–41%, and 3–30%), respectively. The 3-year OS, cumulative incidence of relapse, and non-relapse mortality rates of patients in non-CR (n = 20) were 70% (95% CI: 45–85%), 20% (95% CI: 6–40%), and 10% (95% CI: 2–28%), respectively (Figure 1B, C, D). Next, we divided the patients with respect to the number of human leucocyte antigen (HLA) mismatches for graft-versus-host direction. The cumulative incidence of grade II–IV acute GVHD in patients who underwent CBT with more than two HLA antigen mismatches in the HLA-A, B, C, and DRB1 loci (n = 15) was 80% (95% CI: 46–94%), which was significantly higher than that of the other patients (48%, 95% CI: 30–65%, n = 31) ($p = 0.02$) (Figure 1E). The cumulative incidence of grade II–IV acute GVHD in patients who underwent CBT with more than two HLA allele mismatches in the HLA-A, B, C, and DRB1 loci (n = 24) was 75% (95% CI: 51–88%), which was significantly higher than that of the other patients (41%, 95% CI: 20–61%, n = 22) ($p = 0.02$) (Figure 1F).

Impact of MPA AUCs on outcomes

None of the cases discontinued MMF administration, except for one case of death before day 30. No patients experienced MMF-related severe toxicities such as diarrhea and myelosuppression. The AUC_{0-24} of MPA on day 7 (MPA7) was $49.4 \pm 24.7 \mu\text{g h/mL}$ (mean $\pm$

standard deviation). There was no association between MPA7 and the incidence of PIR, engraftment rate, engraftment speed or acute GVHD after day 7 (data not shown). In the landmark analysis on day 7, the rate of sepsis between day 7 and engraftment in patients with MPA7 >60 µg h/mL was 33% (95% CI: 9–60%), which was higher than that in the other patients (6%, 95% CI: 1–18%) ($p = 0.02$) (Figure 2A). The AUC₀₋₂₄ of MPA on day 21 (MPA21) was 47.7 ± 20.3 µg h/mL. After excluding three cases of graft failure, we divided patients with respect to MPA21. In the landmark analysis on day 21, the cumulative incidence of grade II–IV acute GVHD after day 21 in patients with MPA21 ≤ 30 µg h/mL was 80% (95% CI: 34–96%), which was higher than that in the other patients (50%, 95% CI: 31–67%) ($p = 0.04$) (Figure 2B). In the landmark analysis on day 21, the cumulative incidence of HHV-6 reactivation after day 21 in patients with MPA21 ≤ 48 µg h/mL (median) was 50% (95% CI: 28–68%), which was higher than that in the other patients (19%, 95% CI: 6–38%) ($p = 0.03$) (Figure 2C). Only two patients developed HHV-6 encephalitis, and there was no significant association between the development of HHV-6 encephalitis and the blood level of MPA (data not shown). In the analysis of OS, the cumulative incidence of relapse, chronic GVHD, and reactivation of cytomegalovirus, no significant association was found with the blood level of MPA (data not shown).

Next, we surveyed the causes of low MPA levels. The MPA21 of patients with diarrhea on day 21 was significantly lower than that of patients without diarrhea (35.8 vs. 54.5 µg h/mL,

$p = 0.006$) (Figure 3A). Patients with serum albumin levels <3 g/dL on day 21 tended to have low MPA21 (37.8 vs. 53.8 $\mu\text{g h/mL}$, $p = 0.10$) (Figure 3B).

We assessed the efficacy of the dose escalation of MMF. After 2014, the MPA7 of eight patients was <30 $\mu\text{g h/mL}$ and the MMF dose was escalated after day 7. In these patients, the MPA21 of six patients was higher than their MPA7; the MPA 21 of one patient who had moderate diarrhea was less than MPA7 and one patient died before day 21 (Figure 4A). The cumulative incidence of grade II–IV acute GVHD of these eight patients was 43%, which was lower than the incidence of all patients. The MPA7 of 13 patients was 30–60 $\mu\text{g h/mL}$ and the dose of MMF was not escalated after day 7. Among these patients, nine and four patients had MPA21 less and more than their MPA7s, respectively (Figure 4B). The cumulative incidence of grade II–IV acute GVHD of these 13 patients was 78%, which was higher than the incidence of all patients.

Correlation between AUC_{0-24} and single point concentration

To analyze whether we could reduce the number of blood collections, we investigated the correlation between AUC_{0-24} and single point concentration of MPA. We found that there was stronger correlation between AUC_{0-24} and C_2 (Correlation coefficient = 0.737, $p = 8.69 \times 10^{-17}$), compared with C_0 , C_1 and C_4 (Supplemental Figure 1A-D). However, there was no significant association between C_2 on day 7 and the incidence of sepsis before engraftment,

and between C₂ on day 21 and the incidence of grade II-IV acute GVHD or HHV-6 reactivation (data not shown).

Univariate and multivariate analyses for grade II–IV acute GVHD

After excluding the patients with graft failure and the patients who developed grade II-IV acute GVHD before day 21, we performed univariate and multivariate analyses for grade II–IV acute GVHD after day 21. In the univariate analysis, HLA allele mismatch, non-CR at SCT, and MPA level were associated with a high incidence of grade II–IV acute GVHD (Table 2). Multivariate analysis revealed that non-CR at SCT was associated with a high incidence of grade II–IV acute GVHD (Table 3).

Discussion

In the present study, GVHD prophylaxis with tacrolimus and MMF enabled a high incidence of neutrophil engraftment and prolonged OS compared to those reported in a nationwide survey in Japan.[4, 32] A previous report indicated that 3,000 mg/day of MMF was associated with delayed neutrophil engraftment in CBT,[33] but there was no evidence of an association between the dose of MMF and the rate of neutrophil engraftment. TBI is reportedly necessary for enhanced engraftment in CBT.[34] We administered MMF at 30 mg/kg/day, and 93% of patients received 3–12 Gy of TBI, which may have improved the incidence of neutrophil engraftment. Randomized studies are required to determine the optimal dose of MMF and the best conditioning regimen.

The cumulative incidence of grade II–IV acute GVHD in the present study was higher than that in previous reports.[4, 32] The main reason for this result might be the high percentage of patients who underwent CBT with more than two antigen or allele mismatches in the HLA-A, B, C, and DRB1 loci. In Japan, HLA mismatches are generally evaluated with HLA-A, B, and DRB1 antigens, and fewer than three mismatches were permitted;[35] our institute evaluated HLA mismatches in the same way. Our results suggest that HLA mismatch in CBT should be evaluated with the HLA-A, B, C, and DRB1 alleles to avoid acute GVHD.

In the present study, we observed that the blood levels of MPA fluctuated widely among patients after CBT. The kinetics of MMF are influenced by gastrointestinal damage caused by

conditioning, loss of appetite, decreased oral intake, and other drug interactions.[36] Change from oral to intravenous MMF may be one strategy to stabilize blood level;[37] however, intravenous MMF is not available in Japan.

The appropriate level of MPA in CBT is still not confirmed. We suggest that the AUC_{0-24} on day 7 should be $<60 \mu\text{g h/mL}$ to avoid sepsis before engraftment and that the AUC_{0-24} on day 21 should be $>30 \mu\text{g h/mL}$ to prevent acute GVHD. This range is similar to the adequate AUC_{0-24} used in kidney transplantation.[6, 7]

CBT is one risk factor of HHV-6 encephalitis,[38] and epidemiologic evidence indicates the association between HHV-6 reactivation and acute GVHD.[39] In the present study, the cumulative incidence of HHV-6 reactivation was higher in patients with AUC_{0-24} on day 21 of $\leq 48 \mu\text{g h/mL}$ (median) than in the other patients. Therefore, maintaining a high level of MPA and suppressing graft-versus-host reaction may prevent HHV-6 reactivation.

Based on these results, we hypothesize that MPA level should be changed before and after engraftment. In kidney transplantation, high MPA level is associated with myelosuppression and infection.[7] Similarly, MPA level in CBT should not be high to prevent bacterial infection before engraftment. After engraftment, alloreactive T cells become activated and excessive activation of alloreactive T cells leads to acute GVHD and delays immune reconstitution. Acute GVHD and delay of immune recovery cause HHV-6 reactivation.[39, 40] Moreover, treatment for GVHD also favors HHV-6 reactivation, as described with other

herpesviruses.[41] Therefore, MPA level should be high to suppress alloreactive T cells and prevent acute GVHD and HHV-6 reactivation. These are hypotheses and should be further explored.

After 2014, we escalated the dose of MMF of patients with low blood levels of MPA on day 7 and succeeded in reducing the incidence of grade II–IV acute GVHD in those patients. However, the levels on day 21 in some patients without dose escalation fell below the levels measured on day 7 and many of them developed grade II–IV acute GVHD. More frequent monitoring of MPA and dose adjustment of MMF is necessary, especially for patients with diarrhea or poor nutrition, whose blood levels of MPA tend to be low.

We focused on AUC and collected blood from patients 4 times per day. To proceed to multicenter study, we analyzed whether we could make the measurement simpler. As a result, there was no significant association between single point concentration and the incidence of acute GVHD or infection. Thus, AUC is a better marker to predict the risk of acute GVHD and infection compared with single point concentration, and 4 times blood collection per day is recommended.

Multivariate analysis revealed that non-CR at SCT was associated with the development of grade II–IV acute GVHD, while a low MPA AUC₀₋₂₄ on day 21 was not. In some patients with non-CR at SCT, the physicians decided to maintain low levels of MPA, expecting a graft-versus-tumor effect of cord blood. This was the limitation of this small-scale

retrospective study.

The present study also had additional limitations. We measured whole-blood levels of MPA. Levels of unbound form of MPA should be measured to evaluate the direct effect of MMF on lymphocytes.[42] Moreover, we did not survey the single-nucleotide polymorphism (SNP) for the genes coding inosine-5-monophosphate dehydrogenase (IMPDH). IMPDH is an enzyme inhibited by MPA; SNPs in those genes influence the activity of IMPDH and suppress lymphocyte proliferation.[43-45] In addition, our results should be interpreted with caution because of the small sample size and the heterogeneous patient and transplant backgrounds.

In conclusion, blood AUCs of MPA affected the incidence of grade II–IV acute GVHD, sepsis before engraftment, and HHV-6 reactivation in CBT. A multicenter prospective study is required to clarify the appropriate range of MPA AUCs.

Acknowledgment

The authors thank Emi Furusaka, Tomoko Okuda and Megumi Oka for their expert data management and the hematopoietic stem cell transplantation team members of Kyoto University Hospital for their dedicated care of the patients.

Conflicts of interest statement

The authors declare no conflicts of interest associated with this manuscript.

309 **Funding information**

310 There is no funding associated with this study.

311 **Data availability statement**

312 The data are not publicly available due to privacy or ethical restrictions.

313 **Author contributions**

314 HM, JK, YA and TK designed the research, organized the project, and drafted the
315 manuscript. HM and JK gathered the data and performed the statistical analysis. TS, MH, TY,
316 KY, KM, and ATK contributed to the data analysis and writing of the manuscript. All authors
317 approved the final version of the manuscript for submission for publication.

References

1. Storb R, Deeg HJ, Whitehead J, Appelbaum F, Beatty P, Bensinger W, Buckner CD, Clift R, Doney K, Farewell V. Methotrexate and cyclosporine compared with cyclosporine alone for prophylaxis of acute graft versus host disease after marrow transplantation for leukemia. *N Engl J Med* 1986; 314: 729-35.
2. Storb R, Deeg HJ, Pepe M, Appelbaum F, Anasetti C, Beatty P, Bensinger W, Berenson R, Buckner CD, Clift R. Methotrexate and cyclosporine versus cyclosporine alone for prophylaxis of graft-versus-host disease in patients given HLA-identical marrow grafts for leukemia: long-term follow-up of a controlled trial. *Blood* 1989; 73: 1729-34.
3. Uchida N, Wake A, Nakano N, Ishiwata K, Takagi S, Tsuji M, Yamamoto H, Kato D, Matsuno N, Masuoka K, Araoka H, Asano-Mori Y, Izutsu K, Makino S, Yoneyama A, Taniguchi S. Mycophenolate and tacrolimus for graft-versus-host disease prophylaxis for elderly after cord blood transplantation: a matched pair comparison with tacrolimus alone. *Transplantation* 2011; 92: 366-71.
4. Terakura S, Kuwatsuka Y, Yamasaki S, Wake A, Kanda J, Inamoto Y, Mizuta S, Yamaguchi T, Uchida N, Kouzai Y, Aotsuka N, Ogawa H, Kanamori H, Nishiwaki K, Miyakoshi S, Onizuka M, Amano I, Fukuda T, Ichinohe T, Atsuta Y, Murata M, Teshima T. GvHD prophylaxis after single-unit reduced intensity conditioning cord blood transplantation in adults with acute leukemia. *Bone Marrow Transplant* 2017; 52: 1261-67.
5. Giaccone L, McCune JS, Maris MB, Gooley TA, Sandmaier BM, Slattery JT, Cole S, Nash RA, Storb RF, Georges GE. Pharmacodynamics of mycophenolate mofetil after nonmyeloablative conditioning and unrelated donor hematopoietic cell transplantation. *Blood* 2005; 106: 4381-8.
6. Shaw LM, Figurski M, Milone MC, Trofe J, Bloom RD. Therapeutic drug monitoring of mycophenolic acid. *Clin J Am Soc Nephrol* 2007; 2: 1062-72.
7. Kuypers DR, Le Meur Y, Cantarovich M, Tredger MJ, Tett SE, Cattaneo D, Tönshoff B, Holt DW, Chapman J, Gelder T, MPA TSTCGoTo. Consensus report on therapeutic drug monitoring of mycophenolic acid in solid organ transplantation. *Clin J Am Soc Nephrol* 2010; 5: 341-58.
8. Haentzschel I, Freiberg-Richter J, Platzbecker U, Kiani A, Schetelig J, Illmer T, Ehninger G, Schleyer E, Bornhäuser M. Targeting mycophenolate mofetil for graft-versus-host disease prophylaxis after allogeneic blood stem cell transplantation. *Bone Marrow Transplant* 2008; 42: 113-20.
9. Wakahashi K, Yamamori M, Minagawa K, Ishii S, Nishikawa S, Shimoyama M, Kawano H, Kawano Y, Kawamori Y, Sada A, Matsui T, Katayama Y. Pharmacokinetics-based optimal dose prediction of donor source-dependent response to mycophenolate mofetil in unrelated hematopoietic cell transplantation. *Int J Hematol* 2011; 94: 193-202.
10. Jacobson P, Rogosheske J, Barker JN, Green K, Ng J, Weisdorf D, Tan Y, Long J, Remmel R, Sawchuk R, McGlave P. Relationship of mycophenolic acid exposure to clinical outcome after

354 hematopoietic cell transplantation. *Clin Pharmacol Ther* 2005; 78: 486-500.

355 11. McDermott CL, Sandmaier BM, Storer B, Li H, Mager DE, Boeckh MJ, Bemer MJ, Knutson J,
356 McCune JS. Nonrelapse mortality and mycophenolic acid exposure in nonmyeloablative hematopoietic
357 cell transplantation. *Biol Blood Marrow Transplant* 2013; 19: 1159-66.

358 12. Harnicar S, Ponce DM, Hilden P, Zheng J, Devlin SM, Lubin M, Pozotrigio M, Mathew S,
359 Adel N, Kernan NA, O'Reilly R, Prockop S, Scaradavou A, Hanash A, Jenq R, van den Brink M, Giralt S,
360 Perales MA, Young JW, Barker JN. Intensified Mycophenolate Mofetil Dosing and Higher
361 Mycophenolic Acid Trough Levels Reduce Severe Acute Graft-versus-Host Disease after Double-Unit
362 Cord Blood Transplantation. *Biol Blood Marrow Transplant* 2015; 21: 920-5.

363 13. Arai Y, Kondo T, Kitano T, Hishizawa M, Yamashita K, Kadowaki N, Yamamoto T, Yano I,
364 Matsubara K, Takaori-Kondo A. Monitoring mycophenolate mofetil is necessary for the effective
365 prophylaxis of acute GVHD after cord blood transplantation. *Bone Marrow Transplant* 2015; 50:
366 312-4.

367 14. Beal JL, Jones CE, Taylor PJ, Tett SE. Evaluation of an immunoassay (EMIT) for
368 mycophenolic acid in plasma from renal transplant recipients compared with a high-performance liquid
369 chromatography assay. *Ther Drug Monit* 1998; 20: 685-90.

370 15. Brunet M, Oppenheimer F, Martorell J, Vilardell J, Carreño MC, Carrillo M, Corbella J.
371 Mycophenolic acid monitoring: evaluation of the EMIT MPA immunoassay in kidney and lung
372 transplantation. *Transplant Proc* 1999; 31: 2275-6.

373 16. Parant F, Ranchin B, Gagnieu MC. The Roche Total Mycophenolic Acid® assay: An
374 application protocol for the ABX Pentra 400 analyzer and comparison with LC-MS in children with
375 idiopathic nephrotic syndrome. *Pract Lab Med* 2017; 7: 19-26.

376 17. van Hest RM, Doorduijn JK, de Winter BC, Cornelissen JJ, Vulto AG, Oellerich M,
377 Löwenberg B, Mathot RA, Armstrong VW, van Gelder T. Pharmacokinetics of mycophenolate mofetil
378 in hematopoietic stem cell transplant recipients. *Ther Drug Monit* 2007; 29: 353-60.

379 18. Jacobson P, El-Massah SF, Rogosheske J, Kerr A, Long-Boyle J, DeFor T, Jennissen C,
380 Brunstein C, Wagner J, Tomblyn M, Weisdorf D. Comparison of two mycophenolate mofetil dosing
381 regimens after hematopoietic cell transplantation. *Bone Marrow Transplant* 2009; 44: 113-20.

382 19. Li H, Mager DE, Sandmaier BM, Maloney DG, Bemer MJ, McCune JS. Population
383 pharmacokinetics and dose optimization of mycophenolic acid in HCT recipients receiving oral
384 mycophenolate mofetil. *J Clin Pharmacol* 2013; 53: 393-402.

385 20. Ogata M, Kikuchi H, Satou T, Kawano R, Ikewaki J, Kohno K, Kashima K, Ohtsuka E, Kadota
386 J. Human herpesvirus 6 DNA in plasma after allogeneic stem cell transplantation: incidence and
387 clinical significance. *J Infect Dis* 2006; 193: 68-79.

388 21. Cheson BD, Bennett JM, Kopecky KJ, Büchner T, Willman CL, Estey EH, Schiffer CA,
389 Doehner H, Tallman MS, Lister TA, Lo-Coco F, Willemze R, Biondi A, Hiddemann W, Larson RA,
390 Löwenberg B, Sanz MA, Head DR, Ohno R, Bloomfield CD, LoCocco F, International Working Group
391 for Diagnosis SoRC, Treatment Outcomes, and Reporting Standards for Therapeutic Trials in Acute

392 Myeloid Leukemia. Revised recommendations of the International Working Group for Diagnosis,
393 Standardization of Response Criteria, Treatment Outcomes, and Reporting Standards for Therapeutic
394 Trials in Acute Myeloid Leukemia. *J Clin Oncol* 2003; 21: 4642-9.

395 22. Cheson BD, Fisher RI, Barrington SF, Cavalli F, Schwartz LH, Zucca E, Lister TA, Alliance
396 AsLaLG, Group ECO, Consortium EMCL, Foundation IL, Research EOf, Group ToCDH-O, Ósea
397 GEdM, Group GH-GLS, Group GHsS, Group JLS, Association LS, Group NCT, Group NLS, Group
398 SO, Institute UKNCR. Recommendations for initial evaluation, staging, and response assessment of
399 Hodgkin and non-Hodgkin lymphoma: the Lugano classification. *J Clin Oncol* 2014; 32: 3059-68.

400 23. Giralt S, Ballen K, Rizzo D, Bacigalupo A, Horowitz M, Pasquini M, Sandmaier B.
401 Reduced-intensity conditioning regimen workshop: defining the dose spectrum. Report of a workshop
402 convened by the center for international blood and marrow transplant research. *Biol Blood Marrow*
403 *Transplant* 2009; 15: 367-9.

404 24. Bacigalupo A, Ballen K, Rizzo D, Giralt S, Lazarus H, Ho V, Apperley J, Slavin S, Pasquini M,
405 Sandmaier BM, Barrett J, Blaise D, Lowski R, Horowitz M. Defining the intensity of conditioning
406 regimens: working definitions. *Biol Blood Marrow Transplant* 2009; 15: 1628-33.

407 25. Miyakoshi S, Yuji K, Kami M, Kusumi E, Kishi Y, Kobayashi K, Murashige N, Hamaki T, Kim
408 SW, Ueyama J, Mori S, Morinaga S, Muto Y, Masuo S, Kanemaru M, Hayashi T, Takaue Y, Taniguchi
409 S. Successful engraftment after reduced-intensity umbilical cord blood transplantation for adult
410 patients with advanced hematological diseases. *Clin Cancer Res* 2004; 10: 3586-92.

411 26. Kishi Y, Kami M, Miyakoshi S, Kanda Y, Murashige N, Teshima T, Kusumi E, Hara S,
412 Matsumura T, Yuji K, Masuoka K, Wake A, Morinaga S, Kanemaru M, Hayashi T, Tanaka Y,
413 Taniguchi S, Consortium TSCT. Early immune reaction after reduced-intensity cord-blood
414 transplantation for adult patients. *Transplantation* 2005; 80: 34-40.

415 27. Kaplan EL, Meier P. Nonparametric Estimation from Incomplete Observations. *Journal of*
416 *the American Statistical Association* 1958; 53: 457-81.

417 28. Mantel N. Evaluation of survival data and two new rank order statistics arising in its
418 consideration. *Cancer Chemother Rep* 1966; 50: 163-70.

419 29. Fine JP, Gray RJ. A Proportional Hazards Model for the Subdistribution of a Competing Risk.
420 *Journal of the American Statistical Association* 1999; 94: 496-509.

421 30. Spearman C. The Proof and Measurement of Association between Two Things. *The*
422 *American Journal of Psychology* 1904; 15: 72-101.

423 31. Kanda Y. Investigation of the freely available easy-to-use software 'EZ' for medical statistics.
424 *Bone Marrow Transplant* 2013; 48: 452-8.

425 32. Terakura S, Wake A, Inamoto Y, Murata M, Sakai R, Yamaguchi T, Takahashi S, Uchida N,
426 Onishi Y, Ohashi K, Ozawa Y, Kanamori H, Yamaguchi H, Fukuda T, Ichinohe T, Takanashi M, Atsuta
427 Y, Teshima T. Exploratory research for optimal GvHD prophylaxis after single unit CBT in adults:
428 short-term methotrexate reduced the incidence of severe GvHD more than mycophenolate mofetil.
429 *Bone Marrow Transplant* 2017; 52: 423-30.

33. Okamura A, Shimoyama M, Ishii S, Wakahashi K, Asada N, Kawano H, Kawamori Y, Nishikawa S, Minagawa K, Katayama Y, Matsui T. Delayed neutrophil engraftment in cord blood transplantation with intensive administration of mycophenolate mofetil for GVHD prophylaxis. *Bone Marrow Transplant* 2011; 46: 148-9.
34. Nakasone H, Fuji S, Yakushijin K, Onizuka M, Shinohara A, Ohashi K, Miyamura K, Uchida N, Takanashi M, Ichinohe T, Atsuta Y, Fukuda T, Ogata M, Transplantation CWGoJSfHC. Impact of total body irradiation on successful neutrophil engraftment in unrelated bone marrow or cord blood transplantation. *Am J Hematol* 2017; 92: 171-78.
35. Kanda J, Atsuta Y, Wake A, Ichinohe T, Takanashi M, Morishima Y, Taniguchi S, Takahashi S, Ogawa H, Ohashi K, Ohno Y, Aotsuka N, Onishi Y, Kato K, Nagamura-Inoue T, Kanda Y, Transplantation HWGotJSfHC. Impact of the direction of HLA mismatch on transplantation outcomes in single unrelated cord blood transplantation. *Biol Blood Marrow Transplant* 2013; 19: 247-54.
36. Okamura A, Yamamori M, Shimoyama M, Kawano Y, Kawano H, Kawamori Y, Nishikawa S, Minagawa K, Yakushijin K, Katayama Y, Sakaeda T, Hirai M, Matsui T. Pharmacokinetics-based optimal dose-exploration of mycophenolate mofetil in allogeneic hematopoietic stem cell transplantation. *Int J Hematol* 2008; 88: 104-10.
37. Li H, Mager DE, Bemer MJ, Salinger DH, Vicini P, Sandmaier BM, Nash R, McCune JS. A limited sampling schedule to estimate mycophenolic Acid area under the concentration-time curve in hematopoietic cell transplantation recipients. *J Clin Pharmacol* 2012; 52: 1654-64.
38. Scheurer ME, Pritchett JC, Amirian ES, Zemke NR, Lusso P, Ljungman P. HHV-6 encephalitis in umbilical cord blood transplantation: a systematic review and meta-analysis. *Bone Marrow Transplant* 2013; 48: 574-80.
39. Pichereau C, Desseaux K, Janin A, Scieux C, Peffault de Latour R, Xhaard A, Robin M, Ribaud P, Agbalika F, Chevret S, Socié G. The complex relationship between human herpesvirus 6 and acute graft-versus-host disease. *Biol Blood Marrow Transplant* 2012; 18: 141-4.
40. Greco R, Crucitti L, Noviello M, Racca S, Mannina D, Forcina A, Lorentino F, Valtolina V, Rolla S, Dvir R, Morelli M, Giglio F, Barbanti MC, Lupo Stanghellini MT, Oltolini C, Vago L, Scarpellini P, Assanelli A, Carrabba MG, Marktel S, Bernardi M, Corti C, Clementi M, Peccatori J, Bonini C, Ciceri F. Human Herpesvirus 6 Infection Following Haploidentical Transplantation: Immune Recovery and Outcome. *Biol Blood Marrow Transplant* 2016; 22: 2250-55.
41. Ljungman P, Aschan J, Lewensohn-Fuchs I, Carlens S, Larsson K, Lönnqvist B, Mattsson J, Sparrelid E, Winiarski J, Ringdén O. Results of different strategies for reducing cytomegalovirus-associated mortality in allogeneic stem cell transplant recipients. *Transplantation* 1998; 66: 1330-4.
42. Frymoyer A, Verotta D, Jacobson P, Long-Boyle J. Population pharmacokinetics of unbound mycophenolic acid in adult allogeneic haematopoietic cell transplantation: effect of pharmacogenetic factors. *Br J Clin Pharmacol* 2013; 75: 463-75.
43. Wang J, Yang JW, Zeevi A, Webber SA, Girnita DM, Selby R, Fu J, Shah T, Pravica V,

468 Hutchinson IV, Burckart GJ. IMPDH1 gene polymorphisms and association with acute rejection in
469 renal transplant patients. Clin Pharmacol Ther 2008; 83: 711-7.
470 44. Sombogaard F, van Schaik RH, Mathot RA, Budde K, van der Werf M, Vulto AG, Weimar W,
471 Glander P, Essioux L, van Gelder T. Interpatient variability in IMPDH activity in MMF-treated renal
472 transplant patients is correlated with IMPDH type II 3757T > C polymorphism. Pharmacogenet
473 Genomics 2009; 19: 626-34.
474 45. McCune JS, Storer B, Thomas S, McKiernan J, Gupta R, Sandmaier BM. Inosine
475 Monophosphate Dehydrogenase Pharmacogenetics in Hematopoietic Cell Transplantation Patients.
476 Biol Blood Marrow Transplant 2018; 24: 1802-07.
477

Table 1. Patient characteristics

		Number (%) or Median (Range)
Age at SCT (years)		50 (20–66)
Sex	M/F	29 (63) / 17 (37)
Disease	AML/MDS	20 (43) / 17 (37)
	ALL	6 (13)
	NHL	8 (17)
	CML/MPN	1 (2) / 1 (2)
	ATL	1 (2)
	AA	2 (4)
PS at SCT	0/1	23 (50) / 14 (30)
	2–3	9 (20)
Serum Albumin at SCT (g/dL)		3.2 (1.9–4.2)
HLA Mismatch	0/1/2	1 (2) / 11 (24) / 10 (22)
(A, B, C, and DRB1 alleles)	3/4/5	11 (24) / 11 (24) / 2 (4)
Disease Status	CR/PR	24 (52) / 2 (4)
at SCT	Others	20 (43)
Days from Diagnosis	< 90	6 (13)
to SCT	90–180	14 (30)
	>180	26 (57)
Conditioning	MAC	24 (52)
	RIC	22 (48)
TBI	0	3 (7)
	3–4	21 (46)
	10–12	22 (48)
Total Nuclear Cell Count ($\times 10^7/\text{kg}$)		2.47 (1.94–4.27)
CD34 ⁺ Cell Count ($\times 10^5/\text{kg}$)		0.57 (0.18–1.91)

Abbreviations: SCT: stem cell transplantation, M: male, F: female, AML: acute myeloid leukemia, MDS: myelodysplastic syndrome, ALL: acute lymphoblastic leukemia, NHL:

484 non-Hodgkin lymphoma, CML: chronic myeloid leukemia, MPN: myeloproliferative
485 neoplasm, ATL: adult T cell leukemia/lymphoma, AA: aplastic anemia, PS: performance
486 status, HLA: human leucocyte antigen, CR: complete remission, PR: partial remission, MAC:
487 myeloablative conditioning, RIC: reduced-intensity conditioning, TBI: total body irradiation.

488

489 **Table 2. Univariate analysis for grade II–IV acute GVHD**

Variables	Number	HR	95% CI	<i>P</i>
Age at SCT (years)				
<50	20	1		
≥50	20	1.55	0.72–3.35	0.27
PS at SCT				
0–1	34	1		
2–4	6	0.56	0.16–1.90	0.35
HCT-CI at SCT				
0–1	31	1		
>2	9	0.6	0.22–1.63	0.31
Sex mismatch				
Match, M to F	34	1		
F to M	6	0.99	0.61–1.62	0.98
HLA allele mismatch				
0–2	20	1		
3–5	20	2.26	1.05–4.86	0.04
Disease status at SCT				
CR	22	1		
Non-CR	18	1.29	1.16–1.45	<0.001
Days from diagnosis to SCT				
0–180	16	1		
>180	24	1.01	0.58–1.75	0.98
Conditioning				
MAC	22	1		
RIC	18	0.92	0.41–2.07	0.85
TBI				
No	2	1		
Yes	38	2.41	0.31–18.55	0.40
AUC _{0–24} of MPA on day 21				
≤30 µg h/mL	10	1		
>30 µg h/mL	30	0.38	0.15–0.91	0.03
CD34 ⁺ cell count				
≤ 0.57×10 ⁵ /kg		1		
> 0.57×10 ⁵ /kg		1.27	0.62–2.60	0.51

490

491 Abbreviations: GVHD: graft-versus-host disease, HR: hazard ratio, CI: confidence interval,
492 SCT: stem cell transplantation, PS: performance status, HCT-CI: comorbidity index of
493 hematopoietic cell transplantation, M: male, F: female, HLA: human leucocyte antigen, CR:
494 complete remission, MAC: myeloablative conditioning, RIC: reduced-intensity conditioning,
495 TBI: total body irradiation, AUC₀₋₂₄: 24-hour area under the curve, MPA: mycophenolic acid.

Table 3. Multivariate analysis for grade II–IV acute GVHD

Variables	Number	HR	95% CI	<i>P</i>
HLA allele mismatch				
0–2	20	1		
3–5	20	2.14	0.96–4.79	0.06
Disease status at SCT				
CR	22	1		
Non-CR	18	1.26	1.07–1.48	0.006
AUC ₀₋₂₄ of MPA on day 21				
≤30 µg h/mL	10	1		
>30 µg h/mL	30	0.66	0.24–1.78	0.41

Abbreviations: GVHD: graft-versus-host disease, HR: hazard ratio, CI: confidence interval, HLA: human leucocyte antigen, SCT: stem cell transplantation, CR: complete remission, AUC₀₋₂₄: 24-hour area under the curve, MPA: mycophenolic acid.

Figure legends

Figure 1. Use of mycophenolate mofetil enabled high neutrophil engraftment and feasible overall survival rates.

(A) Cumulative incidence of engraftment. (B) Kaplan–Meier curves of overall survival of patients in CR and non-CR at transplantation. (C) Cumulative incidence of relapse and (D) non-relapse mortality of patients in CR and non-CR. (E) Cumulative incidence of grade II–IV graft-versus-host disease and the number of HLA antigen mismatches (A, B, C, and DRB1). (F) Cumulative incidence of grade II–IV graft-versus-host disease and the number of HLA allele mismatches (A, B, C and DRB1).

Abbreviations: CR: complete remission. HLA: human leukocyte antigen.

Figure 2. AUC₀₋₂₄ of MPA affected the incidence of sepsis, grade II–IV acute GVHD, and reactivation of human herpesvirus 6.

(A) AUC₀₋₂₄ on day 7 and cumulative incidence of sepsis before neutrophil engraftment. (B) AUC₀₋₂₄ on day 21 and cumulative incidence of grade II–IV graft-versus-host disease. (C) AUC₀₋₂₄ on day 21 and cumulative incidence of reactivation of human herpesvirus 6.

Abbreviations: AUC₀₋₂₄: 24-hour area under the curve, MPA: mycophenolic acid, GVHD: graft-versus-host disease.

Figure 3. Diarrhea and poor nutrition decreased AUC₀₋₂₄ of MPA on day 21.

(A) Diarrhea and box plot of AUC₀₋₂₄ on day 21. (B) Serum albumin levels and box plot of AUC₀₋₂₄ on day 21. Abbreviations: AUC₀₋₂₄: 24-hour area under the curve, MPA: mycophenolic acid.

Figure 4. Dose escalation for low AUC₀₋₂₄ on day 7 increased the AUC₀₋₂₄ on day 21.

(A) AUC₀₋₂₄ of MPA on days 7 and 21 among patients with MPA₇ (AUC₀₋₂₄ of MPA on day 7) <30 µg h/mL. The MMF dose of these patients was escalated after day 7. (B) AUC₀₋₂₄ on days 7 and 21 of patients with MPA₇ of 30–60 µg h/mL. The MMF dose of these patients was not escalated after day 7.

Abbreviations: AUC₀₋₂₄: 24-hour area under the curve, MPA: mycophenolic acid, MMF: mycophenolate mofetil.

Supplemental Figure 1. Correlation between AUC₀₋₂₄ and single point concentration.

Correlation between AUC₀₋₂₄ and (A) C₀, (B) C₁, (C) C₂ and (D) C₄.

Abbreviations: AUC₀₋₂₄: 24-hour area under the curve. *R*: correlation coefficient.