

精度保証付き数値計算による楕円型作用素の 逆作用素ノルム評価

渡部 善隆 (Yoshitaka Watanabe)*

九州大学 情報基盤研究開発センター
(Research Institute for Information Technology, Kyushu University)
独立行政法人科学技術振興機構, CREST
(CREST, Japan Science and Technology Agency)

Abstract

本稿では、2 階楕円型線形作用素に対する可逆性と逆作用素ノルムの上界値を数学的に厳密な意味で保証する数値計算法をいくつか紹介する。

1 Introduction

Let $\Omega \subset \mathbb{R}^d$ be a bounded polygonal or polyhedral domain ($d = 1, 2, 3$), and for some integer m , let $H^m(\Omega)$ denote the complex L^2 -Sobolev space of order m on Ω . We define the Hilbert space

$$H_0^1(\Omega) := \{u(x) \in H^1(\Omega) \mid u(x) = 0, x \in \partial\Omega\}$$

with the inner product $(\nabla u, \nabla v)_{L^2(\Omega)}$ and the norm $\|u\|_{H_0^1(\Omega)} := \|\nabla u\|_{L^2(\Omega)}$, where $(u, v)_{L^2(\Omega)}$ implies L^2 -inner product on Ω . Let

$$H(\Delta; L^2(\Omega)) := \{u(x) \in H_0^1(\Omega) \mid \Delta u \in L^2(\Omega)\}$$

be a Banach space with respect to the graph norm $\|u\|_{L^2(\Omega)} + \|\Delta u\|_{L^2(\Omega)}$. Since Ω is in a class of the bounded domain with a Lipschitz continuous boundary, the embedding $H(\Delta; L^2(\Omega)) \hookrightarrow H_0^1(\Omega)$ is compact by the Rellich compactness theorem.

Consider the linear elliptic operator

$$\mathcal{L}u := -\Delta u + b \cdot \nabla u + cu \tag{1}$$

for $b \in L^\infty(\Omega)^d$, $c \in L^\infty(\Omega)$ with norms

$$\|b\|_{L^\infty(\Omega)^d} = \operatorname{ess\,sup}_{x \in \Omega} \sqrt{|b_1(x)|^2 + \cdots + |b_d(x)|^2}, \quad \|c\|_{L^\infty(\Omega)} = \operatorname{ess\,sup}_{x \in \Omega} |c(x)|,$$

respectively.

The aim of this paper is to propose some procedures for verifying the invertibility of an $\mathcal{L}$ with a computable upper bound $M > 0$ satisfying

$$\|u\|_{H_0^1(\Omega)} \leq M \|\mathcal{L}u\|_{H^{-1}(\Omega)}, \quad \forall u \in H_0^1(\Omega) \tag{2}$$

or

$$\|u\|_{H_0^1(\Omega)} \leq M \|\mathcal{L}u\|_{L^2(\Omega)}, \quad \forall u \in H(\Delta; L^2(\Omega)) \tag{3}$$

or

$$\|\Delta u\|_{L^2(\Omega)} \leq M \|\mathcal{L}u\|_{L^2(\Omega)}, \quad \forall u \in H(\Delta; L^2(\Omega)). \tag{4}$$

*This is a joint work with Takehiko Kinoshita (Kyoto University) and Mitsuhiro T. Nakao (National Institute of Technology, Sasebo College)

For example, when one try to find $u \in H_0^1(\Omega)$ (weak sense) satisfying nonlinear problems

$$-\Delta u(x) = f(x, u, \nabla u), \quad x \in \Omega \quad (5)$$

with certain properties for f and apply infinite-dimensional verification approach for u , the norm estimations (2), (3), (4) are required [13, 16, 18, 19, 20]. We note that the upper bound M can also be applied to verified computations of eigenvalue exclosures in Hilbert spaces [25].

2 Approximation subspace and notations

Let S_h be a finite dimensional approximation subspace of $H_0^1(\Omega)$ dependent on the parameter $h > 0$. For example, S_h is taken to be a finite element subspace with mesh size h . Let $P_h : H_0^1(\Omega) \rightarrow S_h$ denote the H_0^1 -projection defined by

$$(\nabla(\phi - P_h\phi), \nabla v)_{L^2(\Omega)} = 0, \quad \forall v \in S_h, \quad (6)$$

and suppose that P_h has the following approximation properties.

$$\|v - P_h v\|_{H_0^1(\Omega)} \leq C(h) \|\Delta v\|_{L^2(\Omega)}, \quad \forall v \in H(\Delta; L^2(\Omega)), \quad (7)$$

$$\|v - P_h v\|_{L^2(\Omega)} \leq C(h) \|v - P_h v\|_{H_0^1(\Omega)}, \quad \forall v \in H_0^1(\Omega), \quad (8)$$

where $C(h) > 0$ is a positive constant which is *numerically* determined with the property that $C(h) \rightarrow 0$ as $h \rightarrow 0$. We emphasize that especially the estimate (7) is indispensable in our argument and the compactness of the embedding $H(\Delta; L^2(\Omega)) \hookrightarrow H^1(\Omega)$ is essential in getting the constant $C(h)$ with desired property. Usually the second estimation (8) for P_h is derived by using a technique so called Aubin-Nitsche's trick [1].

These assumptions (7) and (8) hold for many finite element subspaces of $H_0^1(\Omega)$ [1, 9, 10, 11, 12, 15] or function spaces of Fourier series with finite truncation [23]. For example it can be taken as $C(h) = h/\pi$ and $h/(2\pi)$ for bilinear and biquadratic element, respectively, for the rectangular mesh on the square domain [9], and $C(h) = 0.493h$ for the linear and uniform triangular mesh of the convex polygonal domain [3, 6]. Furthermore, a constructive a priori L^∞ error estimate for the projection P_h can also be obtained [7, 8]. In case of nonconvex polygonal domain, there are some useful techniques and consideration to obtain mathematically rigorous upper bounds for the constant $C(h)$ satisfying (7) with adequate order for such nonconvex domains [2, 5, 14, 26, 27, 28].

Define basis function of S_h by $\{\phi_i\}_{i=1}^N$ for $N := \dim S_h$ and $N \times N$ matrices G , D , L , and Hermitian matrix E by

$$[G]_{ij} = (\nabla \phi_j, \nabla \phi_i)_{L^2(\Omega)} + (b \cdot \nabla \phi_j + c \phi_j, \phi_i)_{L^2(\Omega)}, \quad (9)$$

$$[D]_{ij} = (\nabla \phi_j, \nabla \phi_i)_{L^2(\Omega)}, \quad (10)$$

$$[L]_{ij} = (\phi_j, \phi_i)_{L^2(\Omega)}, \quad (11)$$

$$[E]_{ij} = (b \cdot \nabla \phi_j + c \phi_j, b \cdot \nabla \phi_i + c \phi_i)_{L^2(\Omega)}, \quad (12)$$

respectively. Since D and L are positive definite, they can be decomposed as $D = D^{1/2} D^{H/2}$ and $L = L^{1/2} L^{H/2}$ where H indicates the conjugate transposition. Usually $D^{1/2}$ and $L^{1/2}$ are the lower triangular matrices. We assume that G has the inverse and let $C_p > 0$ denote the Poincaré or Rayleigh-Ritz constants which satisfies

$$\|u\|_{L^2(\Omega)} \leq C_p \|\nabla u\|_{L^2(\Omega)}, \quad u \in H_0^1(\Omega). \quad (13)$$

3 Estimation (2)

This section is devoted to an upper bound M satisfying

$$\|u\|_{H_0^1(\Omega)} \leq M \|\mathcal{L}u\|_{H^{-1}(\Omega)}, \quad \forall u \in H_0^1(\Omega)$$

with the invertibility of $\mathcal{L}$.

It is well-known that for each $\xi \in H^{-1}(\Omega)$ there exists a unique $\psi \in H_0^1(\Omega)$ satisfying

$$\begin{cases} -\Delta\psi &= \xi & \text{in } \Omega, \\ \psi &= 0 & \text{on } \partial\Omega. \end{cases}$$

By define this mapping $\xi \mapsto \psi$ by $(-\Delta)^{-1} : H^{-1}(\Omega) \rightarrow H_0^1(\Omega)$, a map $(-\Delta)^{-1}|_{L^2(\Omega)} : L^2(\Omega) \rightarrow H_0^1(\Omega)$ becomes compact because ψ belongs to $H(\Delta; L^2(\Omega))$ and the embedding $H(\Delta; L^2(\Omega)) \hookrightarrow H^1(\Omega)$ is compact. We define a linear compact operator $F : H_0^1(\Omega) \rightarrow H_0^1(\Omega)$ by

$$Fu := (-\Delta)^{-1}|_{L^2(\Omega)}(-b \cdot \nabla u - cu). \quad (14)$$

Then since the term $-b \cdot \nabla u - cu$ maps each bounded set of $H_0^1(\Omega)$ to a bounded set of $L^2(\Omega)$, the operator F becomes compact on $H_0^1(\Omega)$, and the following is true.

Lemma 1. [13, Theorem 2.3]

If $I - F$ on $H_0^1(\Omega)$ is invertible then so is $\mathcal{L}$, and $M > 0$ of (2) can be taken as satisfying

$$\|(I - F)^{-1}u\|_{H_0^1(\Omega)} \leq M\|u\|_{H_0^1(\Omega)}, \quad \forall u \in H_0^1(\Omega). \quad (15)$$

3.1 1st estimation of (2)

Our first result for (2) is as follows.

Theorem 1. [17, Theorem 1] For

$$C_1 := \|b\|_{L^\infty(\Omega)^d} + C_p\|c\|_{L^\infty(\Omega)}, \quad (16)$$

if $C_p C_1 < 1$ then $I - F$ is invertible and M of (2) can be taken as

$$M = \frac{1}{1 - C_p C_1}. \quad (17)$$

3.2 2nd estimation of (2)

We define

$$C_2 := \|b\|_{L^\infty(\Omega)^d} + C(h)\|c\|_{L^\infty(\Omega)}, \quad (18)$$

$$K := \begin{cases} C(h)(C_p\|\nabla \cdot b\|_{L^\infty(\Omega)} + C_1), & \text{if } b \in W^{1,\infty}(\Omega)^d, \\ C_p C_2, & \text{if } b \in L^\infty(\Omega)^d, \end{cases} \quad (19)$$

$$\rho := \|D^{T/2}G^{-1}D^{1/2}\|_2, \quad (20)$$

where $\|\cdot\|_2$ stands for matrix 2-norm. Note that ρ can be represented by

$$\rho^{-1} = \min\{|\lambda| \mid Gx = \lambda D x, \ 0 \neq x \in \mathbb{C}^n\},$$

and its verified upper bound can be computed [22]. The below is our second estimation of (2).

Theorem 2. [17, Theorem 2] If

$$\kappa := C(h)(\rho C_1 K + C_2) < 1 \quad (21)$$

then $I - F$ is invertible and $M > 0$ of (2) is obtained by

$$M = \frac{1}{1 - \kappa} \left\| \begin{bmatrix} \rho(1 - C_2 C(h)) & \rho K \\ \rho C_1 C(h) & 1 \end{bmatrix} \right\|_2.$$

3.3 3rd estimation of (2)

Defining

$$\begin{aligned}\tilde{K} &:= C(h) (\|b\|_{L^\infty(\Omega)^d} C_1 + \|c\|_{L^\infty(\Omega)}), \\ C_3 &:= C(h) \|b\|_{L^\infty(\Omega)^d},\end{aligned}$$

we have the following result.

Theorem 3. [17, Theorem 3] If $\tilde{\kappa} := \tilde{K}(\rho C_p K + C(h)) < 1$, $I - F$ is invertible and $M > 0$ of (2) is obtained by

$$M = \frac{1}{1 - \tilde{\kappa}} \left\| \begin{bmatrix} \rho(1 - \tilde{K}C(h) + KC_3) & \rho K(1 + C_3) \\ \rho \tilde{K}C_p + C_3 & 1 + C_3 \end{bmatrix} \right\|_2.$$

If $b \in W^{1,\infty}(\Omega)$, $K = O(C(h))$ and then $\tilde{\kappa} = O(C(h))^2$.

3.4 Numerical examples

3.4.1 One-dimensional operators

We use interval arithmetic toolbox INTLAB Version 7 [21] with MATLAB 8.0.0.783 (R2012b) on Intel Core i7 3.4GHz. Divide the interval $(0, 1)$ by equal partition size $h > 0$ and take S_h as the set of piecewise linear functions on each subinterval. We can take $C(h) = h/\pi$ and $C_p = 1/\pi$.

Table 1 and 2 show verification results. The bold letters indicate the smallest M in the theorems.

Table 1: Verification results for $b = \sin(\pi x)$, $c = 1$, $\rho = 1.0035$ ($1/h = 32$)

$1/h$	Theorem 1		Theorem 2		Theorem 3	
	$C_1 C_p$	M	κ	M	$\tilde{\kappa}$	M
4	0.4197	1.7231	0.1057	1.2507	0.0258	1.2186
8	0.4197	1.7231	0.0464	1.1106	0.0065	1.0976
16	0.4197	1.7231	0.0216	1.0521	0.0016	1.0461
32	0.4197	1.7231	0.0104	1.0258	0.0004	1.0229

Table 2: Verification results for $b = -\sin(\pi x)$, $c = -5$, $\rho = 2.0001$ ($1/h = 32$)

$1/h$	Theorem 1		Theorem 2		Theorem 3	
	$C_1 C_p$	M	κ	M	$\tilde{\kappa}$	M
4	0.8250	5.7116	0.2248	2.5155	0.1539	2.4918
8	0.8250	5.7116	0.0770	2.1125	0.0393	2.1122
16	0.8250	5.7116	0.0293	2.0280	0.0099	2.0285
32	0.8250	5.7116	0.0123	2.0082	0.0025	2.0084

3.4.2 Two-dimensional non-self-adjoint operators

Consider the case for

$$b = R \begin{bmatrix} -y + 1/2 \\ x - 1/2 \end{bmatrix}, \quad c \in \mathbb{C}, \quad \Omega = (0, 1) \times (0, 1) \quad (22)$$

We take linear and uniform triangular meshes on Ω with the element side length $h > 0$ for a given finite element mesh. We can take $C(h) = 0.493h$ and $C_p = 1/(\pi\sqrt{2})$. Table 3, 4, and 5 show verification results.

Table 3: Verification results for $R = 4$, $c = 0$, $\rho = 1.0001$ ($1/h = 10$)

$1/h$	Theorem 1		Theorem 2		Theorem 3	
	$C_1 C_p$	M	κ	M	$\tilde{\kappa}$	M
2	0.6367	2.7521	1.1835	—	0.7956	12.5322
5	0.6367	2.7521	0.3567	1.8230	0.1273	1.7994
10	0.6367	2.7521	0.1589	1.2914	0.0319	1.3180

Table 4: Verification results for $R = 6.75$, $c = -1 - 1.5i$, $\rho = 1.0487$ ($1/h = 10$)

$1/h$	Theorem 1		Theorem 2		Theorem 3	
	$C_1 C_p$	M	κ	M	$\tilde{\kappa}$	M
4	1.1658	—	1.0408	—	0.8928	23.7783
5	1.1658	—	0.7608	5.6411	0.5721	5.1856
10	1.1658	—	0.3081	1.7124	0.1433	1.8585

3.5 Report for estimation (2)

We consider three computer-assisted procedures to verify the invertibility of second order linear elliptic operators with a bound for the norm of its inverse. Although it has the limitation, the method of Theorem 1 does not need the computation of ρ (2-norm). The method based on Theorem 3 has the second order for $C(h)$ when $b \in W^{1,\infty}(\Omega)$ and some verification results show that it *could* be an alternative of Theorem 2, especially, some confirmation of the only invertibility for $\mathcal{L}$ are quite essential. We still conclude our second approach of Theorem 2 is robust and reliable than other two approaches.

4 Estimation (3)

Now we consider an upper bound M satisfying

$$\|u\|_{H_0^1(\Omega)} \leq M \|\mathcal{L}u\|_{L^2(\Omega)} \quad \forall u \in H(\Delta; L^2(\Omega)).$$

We have three approaches.

4.1 1st estimation of (3)

Our first result is a direct application of Theorem 2.

Theorem 4. [13, Theorem 2.3] If $\kappa = C(h)(\rho C_1 K + C_2) < 1$ then $\mathcal{L}$ is invertible and $M > 0$ of (3) is obtained by

$$M = \frac{C_p}{1 - \kappa} \left\| \begin{bmatrix} \rho(1 - C_2 C(h)) & \rho K \\ \rho C_1 C(h) & 1 \end{bmatrix} \right\|_2.$$

In Theorem 4, it is expected that $M \rightarrow C_p \max\{\rho, 1\}$.

4.2 2nd estimation of (3)

For

$$\hat{\rho} := \|D^{H/2} G^{-1} L^{1/2}\|_2, \quad (23)$$

we obtained the second estimation.

Table 5: Verification results for $R = 5$, $c = -15$, $\rho = 4.0804$ ($1/h = 20$)

$1/h$	Theorem 1		Theorem 2		Theorem 3	
	$C_1 C_p$	M	κ	M	$\tilde{\kappa}$	M
5	1.5558	—	1.9949	—	2.3104	—
10	1.5558	—	0.6596	11.0853	0.6723	13.9871
20	1.5558	—	0.2148	4.9111	0.1761	5.1964

Theorem 5. [24, Theorem 4.2] If

$$\hat{\kappa} := C(h)C_2(\hat{\rho}C_1 + 1) < 1 \quad (24)$$

then $\mathcal{L}$ is invertible and $M > 0$ of (3) is obtained by

$$M = \frac{\sqrt{\hat{\rho}^2 + C(h)^2(1 + \hat{\rho}C_1)^2}}{1 - \hat{\kappa}}.$$

In Theorem 5, it is expected that $M \rightarrow \hat{\rho}$.

4.3 3rd estimation of (3)

We also present the following estimate based on a fixed-point formulation.

Theorem 6. [4, Theorem 3] If $\kappa = C(h)(\rho C_1 K + C_2) < 1$ then $\mathcal{L}$ is invertible and $M > 0$ of (3) is obtained by

$$M = \frac{\sqrt{\rho^2(C_p + C(h)(K - C_p C_2))^2 + C(h)^2(1 + \rho C_p C_1)^2}}{1 - \kappa}.$$

In Theorem 6, it is expected that $M \rightarrow C_p \rho$.

Comparing three theorems for (3), Theorem 5 could converge to the exact operator norm for $\mathcal{L}^{-1}$. Because of it holds that $\hat{\rho} \leq C_p \rho$, when $\hat{\rho} \sim C_p \rho$, Theorem 6 would apply sufficient “good” M with low computational cost. From the *actual computational* point of view, since the criterion $\hat{\kappa} < 1$ is sometimes harder than $\kappa < 1$ for fixed h experimentally, Theorem 4 and 6 have a room to be effective.

4.4 Numerical examples

Our numerical environment and S_h for one- or two-dimensional operators are same as the previous section.

4.4.1 One-dimensional operators

Table 6, 7, 8, and 9 show verification results for some $b(x) = r \sin(\pi x)$ and $c \in \mathbb{R}$ in $\Omega = (0, 1)$.

4.4.2 Two-dimensional non-self-adjoint operators

Consider the case for (22). Table 10 and 11 show verification results.

4.4.3 Two-dimensional operators

We now report on a case for $b = 0$. Consider an operator: $\mathcal{L} = -\Delta - 1 - 2u_h + 3au_h^2$ which is the linearized the equation

$$\begin{cases} -\Delta u = 1 + u + u^2 - au^3 & \text{in } (0, 1) \times (0, 1), \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

Table 6: Verification results for $b = 2.5 \sin(\pi x)$, $c = -10$

$1/h$	ρ	$\hat{\rho}$	Theorem 4		Theorem 5		Theorem 6	
			κ	M	$\hat{\kappa}$	M	κ	M
10	12.6637	3.6970	0.6865	12.4285	1.9761	—	0.6865	12.2786
30	12.9669	3.8003	0.0956	4.4655	0.6249	10.1500	0.0956	4.4598
50	12.9916	3.8084	0.0409	4.2504	0.3696	6.0452	0.0409	4.2485
100	13.0020	3.8119	0.0142	4.1667	0.1827	4.6645	0.0142	4.1663
200	13.0047	3.8128	0.0056	4.1465	0.0908	4.1936	0.0056	4.1464
500	13.0054	3.8130	0.0019	4.1409	0.0362	3.9561	0.0019	4.1409
1000	13.0055	3.8131	0.0009	4.1401	0.0181	3.8832	0.0009	4.1401

Table 7: Verification results for $b = -20 \sin(\pi x)$, $c = -20$.

$1/h$	ρ	$\hat{\rho}$	Theorem 4		Theorem 5		Theorem 6	
			κ	M	$\hat{\kappa}$	M	κ	M
10	2.6420	0.3552	3.9293	—	6.8074	—	3.9293	—
30	2.5044	0.3542	0.5592	1.8684	2.2167	—	0.5592	1.5439
50	2.4950	0.3542	0.2518	1.0293	1.3246	—	0.2518	0.9502
100	2.4911	0.3542	0.0948	0.8417	0.6603	1.0469	0.0948	0.8249
200	2.4911	0.3542	0.0396	0.8040	0.3296	0.5289	0.0396	0.8002
500	2.4899	0.3542	0.0140	0.7943	0.1318	0.4080	0.0140	0.7938
1000	2.4899	0.3542	0.0067	0.7930	0.0659	0.3792	0.0067	0.7929

at two finite element approximate solutions u_h whose named “lower” and “upper.”

Table 12 and 13 show verification results.

4.5 Report for estimation (3)

The computer-assisted procedure (Theorem 6) is our latest approach to compute a verified bound of the norm for second order linear elliptic operators $\mathcal{L}$. The criterion for the invertibility of $\mathcal{L}$ is the same as Theorem 4, however, it has no limitation such that the lower bound of M is not less than 1. Although the proposed procedure would not converge to its exact operator norm, some verification examples show that it has a better bound than the approach in Theorem 5. We conclude that our proposed method should be a bridge the gap between the two previous approaches, and one may choice an appropriate procedure taking into consideration given problem or computational cost, and so on.

5 Estimation (4)

Finally we consider an upper bound M satisfying

$$\|\Delta u\|_{L^2(\Omega)} \leq M \|\mathcal{L}u\|_{L^2(\Omega)}, \quad \forall u \in H(\Delta; L^2(\Omega)).$$

We have two approaches.

5.1 1st estimation of (4)

Redefining $\rho_{10} := \|D^{H/2}G^{-1}L^{1/2}\|_2$ and defining $\rho_{00} := \|L^{H/2}G^{-1}L^{1/2}\|_2$, we have the first estimation.

Table 8: Verification results for $b = \sin(\pi x)$, $c = 100$.

$1/h$	ρ	$\hat{\rho}$	Theorem 4		Theorem 5		Theorem 6	
			κ	M	$\hat{\kappa}$	M	κ	M
10	0.9183	0.0500	1.1665	—	0.3516	0.1508	1.1665	—
30	0.9911	0.0499	0.1458	0.4977	0.0577	0.0608	0.1458	0.3920
50	0.9969	0.0499	0.0553	0.4060	0.0275	0.0542	0.0553	0.3426
100	0.9992	0.0499	0.0155	0.3568	0.0111	0.0512	0.0155	0.3242
200	0.9998	0.0499	0.0047	0.3365	0.0049	0.0504	0.0047	0.3198
500	1.0000	0.0499	0.0012	0.3254	0.0018	0.0501	0.0012	0.3186
1000	1.0000	0.0499	0.0005	0.3218	0.0009	0.0500	0.0005	0.3184

Table 9: Verification results for $b = \sin(\pi x)$, $c = -10$.

$1/h$	ρ	$\hat{\rho}$	Theorem 4		Theorem 5		Theorem 6	
			κ	M	$\hat{\kappa}$	M	κ	M
10	94.9621	29.6261	2.1281	—	5.2424	—	2.1281	—
30	231.4257	72.4346	0.5767	172.3900	3.5678	—	0.5767	172.4427
50	261.5470	81.8835	0.2366	108.4156	2.3262	—	0.2366	108.4277
100	276.7469	86.6517	0.0641	93.8348	1.1938	—	0.0641	93.8375
200	280.8268	87.9316	0.0171	90.7977	0.5964	217.8445	0.0171	90.7983
500	281.9909	88.2967	0.0032	89.9844	0.2373	115.7653	0.0032	89.9846
1000	282.1580	88.3491	0.0010	89.8696	0.1184	100.2071	0.0010	89.8697

Theorem 7. If $\kappa_7 := C(h)C_2(\rho_{10}C_1 + 1) < 1$ then $\mathcal{L}$ is invertible and $M > 0$ of (4) is obtained by

$$M = 1 + \|b\|_{L^\infty(\Omega)^d} A_1 + \|c\|_{L^\infty(\Omega)} A_0,$$

where

$$A_0 = \frac{\rho_{00} + C(h)^2(1 + \rho_{10}C_1)}{1 - \kappa_7}, \quad A_1 = \frac{\sqrt{\rho_{10}^2 + C(h)^2(1 + \rho_{10}C_1)^2}}{1 - \kappa_7}.$$

5.2 2nd estimation of (4)

Our second result is somewhat constructive than the previous approach.

Theorem 8. For

$$M_h := \sqrt{\|(G^{-1}L^{1/2})^H E(G^{-1}L^{1/2})\|_2}$$

if it holds that

$$\kappa_8 := C(h)C_2(1 + M_h) < 1$$

then $\mathcal{L}$ is invertible and a bound $M > 0$ of (4) can be taken as

$$M = \frac{1 + M_h}{1 - \kappa_8}.$$

Note that if E is positive definite, by using $E = E^{1/2}E^{H/2}$, it is true that

$$M_h = \|E^{H/2}G^{-1}L^{1/2}\|_2.$$

Table 10: Verification results for $R = 10$, $c = -10 - 5i$.

$1/h$	ρ	$\hat{\rho}$	Theorem 4		Theorem 5		Theorem 6	
			κ	M	$\hat{\kappa}$	M	κ	M
5	1.7039	0.3656	2.3287	—	3.6305	—	2.3287	—
10	1.7751	0.3946	0.7724	1.8734	1.7974	—	0.7724	1.6510
20	1.7941	0.4025	0.2814	0.5384	0.8798	3.4926	0.2814	0.5033
50	1.7995	0.4047	0.0869	0.4222	0.3456	0.6227	0.0869	0.4174
100	1.8001	0.4050	0.0392	0.4092	0.1716	0.4897	0.0392	0.4082
130	1.8004	0.4051	0.0294	0.4076	0.1318	0.4670	0.0294	0.4070

Table 11: Verification results for $R = 10$, $c = 15$.

$1/h$	ρ	$\hat{\rho}$	Theorem 4		Theorem 5		Theorem 6	
			κ	M	$\hat{\kappa}$	M	κ	M
5	0.9732	0.1270	1.8758	—	1.9610	—	1.8758	—
8	0.9903	0.1276	0.9032	3.3368	1.1493	—	0.9032	2.6387
10	0.9939	0.1277	0.6488	0.8671	0.8987	1.6951	0.6488	0.6589
20	0.9986	0.1279	0.2497	0.3543	0.4284	0.2453	0.2497	0.2760
50	0.9999	0.1279	0.0818	0.2632	0.1663	0.1559	0.0818	0.2316
100	1.0001	0.1279	0.0379	0.2426	0.0823	0.1400	0.0379	0.2267

5.3 Numerical examples

Consider the case for two-dimensional non-self-adjoint operators (22). Our numerical environment and S_h is same as the previous section. Table 14 and 15 show verification results.

5.4 Report for estimation (4)

We propose two computer-assisted procedures to compute a verified bound $M > 0$ satisfying (4). Some verification examples show that Theorem 8 has a better bound than the approach in Theorem 7. If we are indifferent to computational costs, instead of an estimation

$$\|b \cdot \nabla u_h + cu_h\|_{L^2(\Omega)} \leq M_h(C(h)C_2\|\Delta u\|_{L^2(\Omega)} + \|f\|_{L^2(\Omega)})$$

in the proof of the Theorem 8, it can be possible to use a bound such that

$$\|b \cdot \nabla u_h + cu_h + f\|_{L^2(\Omega)} \leq \hat{M}_h\|f\|_{L^2(\Omega)}$$

with numerically determined $\hat{M}_h > 0$ directly (more constructive).

Acknowledgments

This work was supported by a Grant-in-Aid from the Ministry of Education, Culture, Sports, Science and Technology of Japan (No. 24340018, 23740074, and 24540151).

References

- [1] P.G. Ciarlet, *The Finite Element Method for Elliptic Problems*, North-Holland, Amsterdam, 1978.
- [2] K. Hashimoto, K. Nagatou, and M.T. Nakao, A computational approach to constructive a priori error estimate for finite element approximations of bi-harmonic problems in nonconvex polygonal domains, *Information* **9** 573–580 (2006).

Table 12: Verification results for “lower” u_h at $a = 0.001$. $\hat{\rho}/(C_p \rho) \sim 0.9995$ ($1/h = 50$).

$1/h$	ρ	$\hat{\rho}$	Theorem 4		Theorem 5		Theorem 6	
			κ	M	$\hat{\kappa}$	M	κ	M
10	1.0586	0.2356	0.0030	0.2391	0.0030	0.2421	0.0030	0.2447
20	1.0599	0.2379	0.0008	0.2388	0.0008	0.2395	0.0008	0.2402
30	1.0601	0.2383	0.0004	0.2387	0.0004	0.2391	0.0004	0.2394
40	1.0602	0.2385	0.0002	0.2387	0.0002	0.2389	0.0002	0.2391
50	1.0603	0.2386	0.0002	0.2387	0.0002	0.2388	0.0002	0.2389

Table 13: Verification results for “upper” u_h at $a = 0.001$. $\hat{\rho}/(C_p \rho) \sim 0.6040$ ($1/h = 50$).

$1/h$	ρ	$\hat{\rho}$	Theorem 4		Theorem 5		Theorem 6	
			κ	M	$\hat{\kappa}$	M	κ	M
10	2.5948	0.3545	1.1823	—	0.7722	1.9668	1.1823	—
20	2.6622	0.3624	0.2856	0.9204	0.1861	0.4756	0.2856	0.8883
30	2.6758	0.3640	0.1262	0.7216	0.0822	0.4087	0.1262	0.7074
40	2.6807	0.3645	0.0709	0.6671	0.0461	0.3887	0.0709	0.6590
50	2.6830	0.3648	0.0453	0.6438	0.0295	0.3800	0.0453	0.6386

- [3] F. Kikuchi and X. Liu, Determination of the Babuska-Aziz constant for the linear triangular finite element, *Japan Journal of Industrial and Applied Mathematics* **23** 75–82 (2006).
- [4] T. Kinoshita, Y. Watanabe, and M.T. Nakao, Some remarks on the rigorous estimation of inverse linear elliptic operators, submitted.
- [5] K. Kobayashi, A constructive a priori error estimation for finite element discretizations in a non-convex domain using singular functions, *Japan Journal of Industrial and Applied Mathematics* **26** 493–516 (2009).
- [6] X. Liu and S. Oishi, Verified eigenvalue evaluation for the Laplacian over polygonal domains of arbitrary shape, *SIAM Journal on Numerical Analysis* **51** 1634–1654 (2013).
- [7] M.T. Nakao, Computable L^∞ error estimates in the finite element method with application to non-linear elliptic problems, in *Contributions in Numerical Mathematics*, pp. 309–319, ed. R. P. Agarwal, World Scientific, Singapore, 1993.
- [8] M.T. Nakao and N. Yamamoto, Numerical verification of solutions for nonlinear elliptic problems using L^∞ residual method, *Journal of Mathematical Analysis and Applications* **217** 246–262 (1998).
- [9] M.T. Nakao, N. Yamamoto, and S. Kimura, On best constant in the error bound for the H_0^1 -projection into piecewise polynomial spaces, *Journal of Approximation Theory* **93** 491–500 (1998).
- [10] M.T. Nakao and N. Yamamoto, A guaranteed bound of the optimal constant in the error estimates for linear triangular element, in *Topics in Numerical Analysis With Special Emphasis on Nonlinear Problems*, eds. G. Alfeld, and X. Chen, pp. 163–173, Computing Supplement, vol. 15, Springer, Wien, New York, 2001.
- [11] M.T. Nakao and N. Yamamoto, A guaranteed bound of the optimal constant in the error estimates for linear triangular element Part II: Details, in *Perspectives on Enclosure Methods*, eds. U. Kulisch, R. Lohner, and A. Facius, pp. 265–276, Springer, Wien, New York, 2001.
- [12] M.T. Nakao, Numerical verification methods for solutions of ordinary and partial differential equations, *Numerical Functional Analysis and Optimization* **22** 321–356 (2001).

Table 14: Verification results for $R = 20$, $c = 0$.

$1/h$	Theorem 7	Theorem 8	M_h	ρ_{10}	ρ_{00}	A_0	A_1
20	—	3.5386	0.5843	0.2238	0.0501	—	—
30	108.0393	2.5102	0.5854	0.2242	0.0503	1.6592	7.5689
40	12.9217	2.1922	0.5861	0.2243	0.0504	0.1867	0.8430
50	8.7114	2.0372	0.5865	0.2244	0.0504	0.1214	0.5453
100	5.4943	1.7847	0.5872	0.2244	0.0504	0.0712	0.3178
130	5.0987	1.7350	0.5873	0.2244	0.0504	0.0650	0.2899

Table 15: Verification results for $R = 10$, $c = -10 - 10i$.

$1/h$	Theorem 7	Theorem 8	M_h	ρ_{10}	ρ_{00}	A_0	A_1
20	16.3072	3.2671	1.0451	0.3172	0.0712	0.3314	1.5020
30	7.7991	2.7135	1.0469	0.3177	0.0714	0.1483	0.6651
40	6.3207	2.5057	1.0475	0.3179	0.0715	0.1164	0.5199
50	5.7107	2.3968	1.0478	0.3180	0.0715	0.1032	0.4600
100	4.8430	2.2077	1.0482	0.3181	0.0716	0.0843	0.3750
130	4.6888	2.1683	1.0483	0.3181	0.0716	0.0809	0.3599

- [13] M.T. Nakao, K. Hashimoto, and Y. Watanabe, A numerical method to verify the invertibility of linear elliptic operators with applications to nonlinear problems, *Computing* **75** 1–14 (2005).
- [14] M.T. Nakao, K. Hashimoto, and K. Kobayashi, Verified numerical computation of solutions for the stationary Navier-Stokes equation in nonconvex polygonal domains, *Hokkaido Mathematical Journal* **36** 777–799 (2007).
- [15] M.T. Nakao, K. Hashimoto, and K. Nagatou, A computational approach to constructive a priori and a posteriori error estimates for finite element approximations of bi-harmonic problems, in *Proceedings of the 4th JSIAM-SIMAI Seminar on Industrial and Applied Mathematics*, GAKUTO International Series, Mathematical Sciences and Applications, vol. 28, pp. 139–148, Gakkotosho, Tokyo, Japan, 2008.
- [16] M.T. Nakao and Y. Watanabe, Numerical verification methods for solutions of semilinear elliptic boundary value problems, *Nonlinear Theory and Its Applications, IEICE* **2** 2–31 (2011).
- [17] M.T. Nakao, Y. Watanabe, T. Kinoshita, T. Kimura, and N. Yamamoto, Some considerations of the invertibility verifications for linear elliptic operators, *Japan Journal of Industrial and Applied Mathematics*, to appear.
- [18] M. Plum, Explicit H_2 -estimates and pointwise bounds for solutions of second-order elliptic boundary value problems, *Journal of Mathematical Analysis and Applications* **165** 36–61 (1992).
- [19] M. Plum, Numerical existence proofs and explicit bounds for solutions of nonlinear elliptic boundary value problems, *Computing* **49** 25–44 (1992).
- [20] M. Plum, Existence and multiplicity proofs for semilinear elliptic boundary value problems by computer assistance, *Jahresbericht der Deutschen Mathematiker Vereinigung* **110** 19–54 (2008).
- [21] S.M. Rump, INTLAB – INTerval LABoratory, in *Developments in Reliable Computing*, ed. T. Csendes, pp. 77–104, Kluwer Academic Publishers, Dordrecht, 1999.
<http://www.ti3.tu-harburg.de/rump/>
- [22] S.M. Rump, Verified bounds for singular values, in particular for the spectral norm of a matrix and its inverse, *BIT Numerical Mathematics* **51** 367–384 (2011).

- [23] Y. Watanabe, N. Yamamoto, M.T. Nakao, and T. Nishida, A numerical verification of nontrivial solutions for the heat convection problem, *Journal of Mathematical Fluid Mechanics* **6** 1–20 (2004).
- [24] Y. Watanabe, T. Kinoshita, and M.T. Nakao, A posteriori estimates of inverse operators for boundary value problems in linear elliptic partial differential equations, *Mathematics of Computation* **82** 1543–1557 (2013).
- [25] Y. Watanabe, K. Nagatou, M. Plum, and M.T. Nakao, Verified computations of eigenvalue exclosures for eigenvalue problems in Hilbert spaces, *SIAM Journal on Numerical Analysis* **52** 975–992 (2014).
- [26] N. Yamamoto and M.T. Nakao, Numerical verifications of solutions for elliptic equations in non-convex polygonal domains, *Numerische Mathematik* **65** 503–521 (1993).
- [27] N. Yamamoto and K. Hayakawa, Error estimation with guaranteed accuracy of finite element method in nonconvex polygonal domains, *Journal of Computational and Applied Mathematics* **159** 173–183 (2003).
- [28] N. Yamamoto and K. Gemma, On error estimation of finite element approximations to the elliptic equations in nonconvex polygonal domains, *Journal of Computational and Applied Mathematics* **199** 286–296 (2007).