

Error Bounds of P-matrix Linear Complementarity Problems¹

Xiaojun Chen²

Shuhuang Xiang³

1 Introduction

The linear complementarity problem is to find a vector $x \in R^n$ such that

$$Mx + q \geq 0, \quad x \geq 0, \quad x^T(Mx + q) = 0,$$

or to show that no such vector exists, where $M \in R^{n \times n}$ and $q \in R^n$. We denote this problem by $LCP(M, q)$. A matrix M is called a P-matrix if

$$\max_{1 \leq i \leq n} x_i(Mx)_i > 0 \quad \text{for all } x \neq 0.$$

It is well-known that M is a P-matrix if and only if the $LCP(M, q)$ has a unique solution for any $q \in R^n$ [6]. Recall the following definitions for an $n \times n$ matrix.

M is called an M-matrix, if $M^{-1} \geq 0$ and $M_{ij} \leq 0$ ($i \neq j$) for $i, j = 1, 2, \dots, n$.

M is called an H-matrix, if its comparison matrix is an M-matrix.

It is known that an H-matrix with positive diagonals is a P-matrix. Moreover, if M is a P-matrix, then there is a neighborhood $\mathcal{M}$ of M , such that all matrices in $\mathcal{M}$ are P-matrices. Hence, we can define a solution function $x(A, b) : \mathcal{M} \times R^n \rightarrow R_+^n$, where $x(A, b)$ is the solution of $LCP(A, b)$ and $R_+^n = \{x \in R^n \mid x \geq 0\}$.

It is easy to verify that x^* solves the $LCP(M, q)$ if and only if x^* solves

$$r(x) := \min(x, Mx + q) = 0,$$

where the min operator denotes the componentwise minimum of two vectors. The function r is called the natural residual of the $LCP(M, q)$, and often used in error analysis. Error bounds for the $LCP(M, q)$ have been studied extensively, see [3, 6, 7, 11, 9, 12, 15].

¹This work is partly supported by a Grant-in-Aid from Japan Society for the Promotion of Science.

²Department of Mathematical System Science, Hirosaki University, Hirosaki 036-8561, Japan, chen@cc.hirosaki-u.ac.jp.

³Department of Applied Mathematics and Software, Central South University, Changsha, Hunan 410083, China, xiangsh@mail.csu.edu.cn

2 Global error bounds for P-matrix linear complementarity problems

For M being a P-matrix, Mathias and Pang [11] present the following error bound

$$\|x - x^*\|_\infty \leq \frac{1 + \|M\|_\infty}{c(M)} \|r(x)\|_\infty, \quad (2.1)$$

for any $x \in R^n$, where

$$c(M) = \min_{\|x\|_\infty=1} \left\{ \max_{1 \leq i \leq n} x_i (Mx)_i \right\}.$$

This error bound is well known and widely cited. However, the quantity $c(M)$ in (2.1) is not easy to find. For M being an H-matrix with positive diagonals, Mathias and Pang [11] gave a computable lower bound for $c(M)$,

$$c(M) \geq \frac{(\min_i b_i)(\min_i (\tilde{M}^{-1}b)_i)}{(\max_j (\tilde{M}^{-1}b)_j)^2} =: \tilde{c}(b), \quad (2.2)$$

for any vector $b > 0$, where $\tilde{M}$ is the comparison matrix of M , that is

$$\tilde{M}_{ii} = M_{ii} \quad \tilde{M}_{ij} = -|M_{ij}| \quad \text{for } i \neq j.$$

However, finding a large value of $\tilde{c}(b)$ is not easy. For some b , $\tilde{c}(b)$ can be very small, and thus the error coefficient

$$\mu(b) := \frac{1 + \|M\|_\infty}{\tilde{c}(b)} \quad (2.3)$$

can be very large.

Interval methods for validation of solution of the LCP(M, q) have been studied in [1, 14]. When a numerical validation condition for the existence of a solution holds, a numerical error bound is provided. However, the numerical validation condition is not ensured to be held at every point x .

In [4], we observed that for every $x, y \in R^n$,

$$\min(x_i, y_i) - \min(x_i^*, y_i^*) = (1 - d_i)(x_i - x_i^*) + d_i(y_i - y_i^*), \quad i \in N \quad (2.4)$$

where

$$d_i = \begin{cases} 0 & \text{if } y_i \geq x_i, y_i^* \geq x_i^* \\ 1 & \text{if } y_i \leq x_i, y_i^* \leq x_i^* \\ \frac{\min(x_i, y_i) - \min(x_i^*, y_i^*) + x_i^* - x_i}{y_i - y_i^* + x_i^* - x_i} & \text{otherwise.} \end{cases}$$

Moreover, we have $d_i \in [0, 1]$. Hence putting $y = Mx + q$ and $y^* = Mx^* + q$ in (2.4), we obtain

$$r(x) = (I - D + DM)(x - x^*), \quad (2.5)$$

where D is a diagonal matrix whose diagonal elements are $d = (d_1, d_2, \dots, d_n) \in [0, 1]^n$.

It is known that M is a P-matrix if and only if $I - D + DM$ is nonsingular for any diagonal matrix $D = \text{diag}(d)$ with $0 \leq d_i \leq 1$ [10]. This together with (2.5) yields upper and lower error bounds,

$$\frac{\|r(x)\|}{\max_{d \in [0, 1]^n} \|I - D + DM\|} \leq \|x - x^*\| \leq \max_{d \in [0, 1]^n} \|(I - D + DM)^{-1}\| \|r(x)\|. \quad (2.6)$$

Moreover, it is not difficult to verify that if M is a P-matrix and $D = \text{diag}(d)$ with $d \in [0, 1]^n$, we have

$$\max_{1 \leq i \leq n} x_i ((I - D + DM)x)_i > 0, \quad \text{for all } x \neq 0,$$

that is, $(I - D + DM)$ is a P-matrix. Therefore, computation of rigorous error bounds can be turned into $\|\cdot\|$ optimization problems over a P-matrix interval set, which is related to linear P-matrix interval systems.

The linear interval system has been studied intensively and some highly efficient numerical methods have been developed, see [13, 14] for references. In the rest part of this section, we give some simple upper bounds for

$$\max_{d \in [0, 1]^n} \|(I - D + DM)^{-1}\|.$$

Theorem 2.1 [4] *Suppose that M is an H-matrix with positive diagonals. Then we have*

$$\max_{d \in [0, 1]^n} \|(I - D + DM)^{-1}\| \leq \|\tilde{M}^{-1} \max(\Lambda, I)\|. \quad (2.7)$$

Remark 1. Since $\tilde{M}^{-1} \max(\Lambda, I) \geq 0$, we have

$$\|\tilde{M}^{-1} \max(\Lambda, I)\|_\infty = \|\tilde{M}^{-1} \max(\Lambda, I)e\|_\infty$$

and

$$\|\tilde{M}^{-1} \max(\Lambda, I)\|_1 = \|(e^T \tilde{M}^{-1} \max(\Lambda, I))^T\|_\infty.$$

The upper error bound in (2.7) with $\|\cdot\|_\infty$ or $\|\cdot\|_1$ can be computed by solving a linear system of equations $\min(\Lambda^{-1}, I)\tilde{M}x = e$ or $\tilde{M}^T \min(\Lambda^{-1}, I)x = e$.

Theorem 2.2 [4] Suppose that M is an M -matrix. Let $V = \{v \mid M^T v \leq e, v \geq 0\}$ and $f(v) = \max_{1 \leq i \leq n} (e + v - M^T v)_i$. Then we have

$$\max_{d \in [0,1]^n} \|(I - D + DM)^{-1}\|_1 = \max_{v \in V} f(v). \quad (2.8)$$

Theorem 2.3 [4] If M is a P -matrix, then for any $x \in R^n$, the following inequalities hold.

$$\begin{aligned} & \frac{1}{1 + \|M\|_\infty} \|r(x)\|_\infty \quad (\text{Mathias-Pang [11]}) \\ & \leq \frac{1}{\max(1, \|M\|_\infty)} \|r(x)\|_\infty \quad (\text{Cottle-Pang-Stone [6]}) \\ & = \frac{1}{\max_{d \in [0,1]^n} \|I - D + DM\|_\infty} \|r(x)\|_\infty \\ & \leq \|x - x^*\|_\infty \\ & \leq \max_{d \in [0,1]^n} \|(I - D + DM)^{-1}\|_\infty \|r(x)\|_\infty \\ & \leq \frac{\max(1, \|M\|_\infty)}{c(M)} \|r(x)\|_\infty \\ & = \frac{1 + \|M\|_\infty}{c(M)} \|r(x)\|_\infty - \frac{\min(1, \|M\|_\infty)}{c(M)} \|r(x)\|_\infty \\ & \leq \frac{1 + \|M\|_\infty}{c(M)} \|r(x)\|_\infty \quad (\text{Mathias-Pang [11]}). \end{aligned}$$

Theorem 2.4 [4] If M is an H -matrix with positive diagonals, then for any $x, b \in R^n$, $b > 0$, the following inequalities hold.

$$\begin{aligned} & \|x - x^*\|_\infty \\ & \leq \max_{d \in [0,1]^n} \|(I - D + DM)^{-1}\|_\infty \|r(x)\|_\infty \\ & \leq \|\tilde{M}^{-1} \max(\Lambda, I)\|_\infty \|r(x)\|_\infty \\ & \leq (\mu(b) - \|\tilde{M}^{-1} \min(\Lambda, I)\|_\infty) \|r(x)\|_\infty \\ & \leq \mu(b) \|r(x)\|_\infty \quad (\text{Mathias-Pang [11]}). \end{aligned}$$

In addition, if M is an M -matrix, then for any $x \in R^n$, the following inequalities hold.

$$\begin{aligned} & \|x - x^*\|_\infty \\ & \leq \|M^{-1} \max(\Lambda, I)\|_\infty \|r(x)\|_\infty \\ & \leq \left(\frac{1 + \|M\|_\infty}{c(M)} - \|M^{-1} \min(\Lambda, I)\|_\infty \right) \|r(x)\|_\infty \\ & \leq \frac{1 + \|M\|_\infty}{c(M)} \|r(x)\|_\infty \quad (\text{Mathias-Pang [11]}). \end{aligned}$$

Applying Theorem 2.1, we obtain the following relative error bounds

Corollary 2.1 [4] *Suppose M is an H-matrix with positive diagonals. For any $x \in R^n$, we have*

$$\frac{\|r(x)\|}{(1 + \|M\|)\|\tilde{M}^{-1} \max(\Lambda, I)\| \|(-q)_+\|} \leq \frac{\|x - x^*\|}{\|x^*\|} \leq \frac{\|M\| \|\tilde{M}^{-1} \max(\Lambda, I)\| \|r(x)\|}{\|(-q)_+\|}.$$

3 Perturbation bounds of P-matrix linear complementarity problems

In [6], Cottle, Pang and Stone introduced the following Lemma which has been widely applied in perturbation bounds based on the fundamental quantity associated with a P-matrix,

$$c(M) = \min_{\|x\|_\infty=1} \max_{1 \leq i \leq n} \{x_i(Mx)_i\}.$$

Lemma 3.1 [6] *Let $M \in R^{n \times n}$ be a P-matrix. The following statements hold:*

(i) *for any two vectors q and p in R^n ,*

$$\|x(M, q) - x(M, p)\|_\infty \leq \frac{1}{c(M)} \|q - p\|_\infty$$

(ii) *for each vector $q \in R^n$, there exist a neighborhood $\mathcal{U}$ of the pair (M, q) and a constant $c_0 > 0$ such that for any $(A, b), (B, p) \in \mathcal{U}$, A, B are P-matrices and*

$$\|x(A, b) - x(B, p)\|_\infty \leq c_0 (\|A - B\|_\infty + \|b - p\|_\infty).$$

Lemma 3.1 shows that when M is a P-matrix, for each q , $x(A, b)$ is a locally Lipschitzian function of (A, b) in a neighborhood of (M, q) , and $x(M, b)$ is a globally Lipschitzian function of b . This property plays a very important rule in the study of the LCP and mathematical programs with LCP constraints [8]. However, the constant $c(M)$ is difficult to compute, and c_0 is not specified. It is hard to use this lemma for verifying accuracy of a computed solution of the LCP when the data (M, q) contain errors.

For M being a P-matrix, we [5] introduce the following constant

$$\beta(M) = \max_{d \in [0, 1]^n} \|(I - D + DM)^{-1} D\|.$$

In the follows, we compare $\beta(M)$ with $c(M)^{-1}$ in $\|\cdot\|_\infty$ and give a simple version of $\beta(M)$ for M being an M-matrix, a symmetric positive definite matrix, and positive definite matrix.

Theorem 3.1 [5] *Let M be a P-matrix. Then*

$$\beta_{\infty}(M) := \max_{d \in [0,1]^n} \|(I - D + DM)^{-1}D\|_{\infty} \leq \frac{1}{c(M)}.$$

It is known that an H-matrix with positive diagonals is a P-matrix, and a positive definite matrix is a P-matrix [6]. Now, we consider the two subclasses of P-matrix.

Theorem 3.2 [5] *Let M be an H-matrix with positive diagonals. Then*

$$\beta(M) \leq \|\tilde{M}^{-1}\|,$$

where $\tilde{M}$ is the comparison matrix of M . In particular, if M is an M-matrix, then the equality holds.

Theorem 3.3 [5] *Let M be a symmetric positive definite matrix. Then*

$$\beta_2(M) := \max_{d \in [0,1]^n} \|(I - D + DM)^{-1}D\|_2 = \|M^{-1}\|_2.$$

In comparison to Lemma 3.1, the following theorem gives sharp perturbation error estimates for the P-matrix LCP

Theorem 3.4 [5] *Let $M \in R^{n \times n}$ be a P-matrix. Then the following statements hold:*

(i) *For any two vectors q and p in R^n ,*

$$\|x(M, q) - x(M, p)\| \leq \beta(M)\|q - p\|.$$

(ii) *Every matrix $A \in \mathcal{M} := \{A \mid \beta(M)\|M - A\| \leq \eta < 1\}$ is a P-matrix. Let*

$$\alpha(M) = \frac{1}{1 - \eta} \beta(M).$$

Then for any $A, B \in \mathcal{M}$ and $q, p \in R^n$

$$\|x(A, q) - x(B, p)\| \leq \alpha(M)^2 \|(-p)_+\| \|A - B\| + \alpha(M)\|q - p\|.$$

From Theorem 3.2 and Theorem 3.3, the Lipschitz constants $\beta(M)$ and $\alpha(M)$ can be estimated by matrix norms, if M is an H-matrix with positive diagonals or a symmetric positive definite matrix. In particular, we have the following two corollaries.

Corollary 3.1 [5] *Let $M \in R^{n \times n}$ be an H-matrix with positive diagonals. Then the following statements hold:*

(i) For any two vectors q and p in R^n ,

$$\|x(M, q) - x(M, p)\|_\infty \leq \|\tilde{M}^{-1}\|_\infty \|q - p\|_\infty$$

(ii) Every matrix $A \in \mathcal{M}_\infty := \{A \mid \|\tilde{M}^{-1}\|_\infty \|M - A\|_\infty \leq \eta < 1\}$ is an H -matrix with positive diagonals. Let

$$\alpha_\infty(M) = \frac{1}{1 - \eta} \|\tilde{M}^{-1}\|_\infty.$$

Then for any $A, B \in \mathcal{M}_\infty$ and $q, p \in R^n$

$$\|x(A, q) - x(B, p)\|_\infty \leq \alpha_\infty(M)^2 \|(-p)_+\|_\infty \|A - B\|_\infty + \alpha_\infty(M) \|q - p\|_\infty.$$

Corollary 3.2 [5] Let $M \in R^{n \times n}$ be a symmetric positive definite matrix. Then the following statements hold:

(i) For any two vectors q and p in R^n ,

$$\|x(M, q) - x(M, p)\|_2 \leq \|M^{-1}\|_2 \|q - p\|_2$$

(ii) Every matrix $A \in \mathcal{M}_2 := \{A \mid \|M^{-1}\|_2 \|M - A\|_2 \leq \eta < 1\}$ is a P -matrix. Let

$$\alpha_2(M) = \frac{1}{1 - \eta} \|M^{-1}\|_2.$$

Then for any $A, B \in \mathcal{M}_2$ and $q, p \in R^n$

$$\|x(A, q) - x(B, p)\|_2 \leq \alpha_2(M)^2 \|(-p)_+\|_2 \|A - B\|_2 + \alpha_2(M) \|q - p\|_2.$$

A matrix A is called positive definite if

$$x^T A x > 0, \quad 0 \neq x \in R^n.$$

Since $x^T A x = x^T \frac{A + A^T}{2} x$, A is positive definite if and only if $\frac{A + A^T}{2}$ is symmetric positive definite. Note that a positive definite matrix is not necessarily symmetric. Such asymmetric matrices frequently appear in the context of the LCP.

Combining the ideas of Mathias and Pang [11] and Corollary 3.2, we present perturbation bounds for the positive definite matrix LCP.

Theorem 3.5 [5] Let $M \in R^{n \times n}$ be a positive definite matrix. Then the following statements hold:

(i) For any two vectors q and p in R^n ,

$$\|x(M, q) - x(M, p)\|_2 \leq \left\| \left(\frac{M + M^T}{2} \right)^{-1} \right\|_2 \|q - p\|_2.$$

(ii) Every matrix $A \in \mathcal{M}_2 := \{A \mid \left\| \left(\frac{M + M^T}{2} \right)^{-1} \right\|_2 \|M - A\|_2 \leq \eta < 1\}$ is positive definite.

Let

$$\alpha_2(M) = \frac{1}{1 - \eta} \left\| \left(\frac{M + M^T}{2} \right)^{-1} \right\|_2.$$

Then for any $A, B \in \mathcal{M}_2$ and $q, p \in R^n$

$$\|x(A, q) - x(B, p)\|_2 \leq \alpha_2(M)^2 \|(-p)_+\|_2 \|A - B\|_2 + \alpha_2(M) \|q - p\|_2.$$

Example 3.1 Theorem 3.1 shows that for every P-matrix, $\beta_\infty(M) \leq c(M)^{-1}$. Now we show that $\beta_\infty(M)$ can be much smaller than $c(M)^{-1}$ in some case. Consider

$$M = \begin{pmatrix} 1 & -t \\ 0 & t \end{pmatrix}.$$

For $t \geq 1$, M is an M-matrix. By Theorem 3.2, $\beta_\infty(M) = \|M^{-1}\|_\infty = 2$. For $\bar{x} = (1, t^{-1})$, we have

$$c(M) \leq \max_{i \in N} \bar{x}_i (M\bar{x})_i = \frac{1}{t}.$$

Hence, $c(M)^{-1} \geq t \rightarrow \infty$, as $t \rightarrow \infty$.

Using the results in the last section, we derive relative perturbation bounds expressed in the term of $\beta(M)\|M\|$.

For the system of linear equations, A is nonsingular if and only if $Ax = b$ has a unique solution for any vector b . A system of linear equations is considered to be well-conditioned (ill-conditioned) if small changes in A or b result in small (large) changes in the solution x . The condition number of A is a measure of sensitivity of the solution of $Ax = b$ for A being a nonsingular matrix. For the linear complementarity problem, M is a P-matrix if and only if $\text{LCP}(M, q)$ has a unique solution for any vector q . A linear complementarity problem is considered to be well-conditioned (ill-conditioned) if small changes in M or q result in small (large) changes in the solution x . Based on the preceding analysis, we are able to give a perturbation theorem for the P-matrix LCP, and define a measure of sensitivity of the solution of $\text{LCP}(M, q)$ for M being a P-matrix.

Theorem 3.6 [5] *Suppose*

$$\begin{aligned} \min(x, Mx + q) &= 0 & M \in R^{n \times n}, \quad 0 \neq (-q)_+ \in R^n \\ \min(y, (M + \Delta M)y + q + \Delta q) &= 0 & \Delta M \in R^{n \times n}, \quad \Delta q \in R^n. \end{aligned}$$

with

$$\|\Delta M\| \leq \epsilon \|M\|, \quad \|\Delta q\| \leq \epsilon \max(\|(-q)_+\|, \|q\| - \|Mx + q\|).$$

If M is a P-matrix and $\epsilon\beta(M)\|M\| = \eta < 1$, then $M + \Delta M$ is a P-matrix and

$$\frac{\|y - x\|}{\|x\|} \leq \frac{2\epsilon}{1 - \eta} \beta(M) \|M\|.$$

Theorem 3.6 indicates that $\beta(M)\|M\|$ is a measure of sensitivity of the solution of the LCP(M, q) for M being a P-matrix. Application of Theorem 3.6 with Corollary 3.1, Corollary 3.2 and Theorem 3.5 gives $\beta(M)\|M\|$ in the term of condition number for the H-matrix LCP, symmetric positive definite LCP and positive definite LCP.

Corollary 3.3 [5] *Suppose*

$$\begin{aligned} \min(x, Mx + q) &= 0 & M \in R^{n \times n}, \quad 0 \neq (-q)_+ \in R^n \\ \min(y, (M + \Delta M)y + q + \Delta q) &= 0 & \Delta M \in R^{n \times n}, \quad \Delta q \in R^n. \end{aligned}$$

(i) *If M is an H-matrix with positive diagonals, $\epsilon\kappa_\infty(\tilde{M}) = \eta < 1$, and*

$$\|\Delta M\|_\infty \leq \epsilon \|\tilde{M}\|_\infty, \quad \|\Delta q\|_\infty \leq \epsilon \max(\|(-q)_+\|_\infty, \|q\|_\infty - \|Mx + q\|_\infty)$$

then $M + \Delta M$ is an H-matrix with positive diagonals and

$$\frac{\|y - x\|_\infty}{\|x\|_\infty} \leq \frac{2\epsilon}{1 - \eta} \kappa_\infty(\tilde{M}).$$

(ii) *If M is a symmetric positive definite matrix, $\epsilon\kappa_2(M) = \eta < 1$, and*

$$\|\Delta M\|_2 \leq \epsilon \|M\|_2, \quad \|\Delta q\|_2 \leq \epsilon \max(\|(-q)_+\|_2, \|q\|_2 - \|Mx + q\|_2),$$

then $M + \Delta M$ is a P-matrix and

$$\frac{\|y - x\|_2}{\|x\|_2} \leq \frac{2\epsilon}{1 - \eta} \kappa_2(M).$$

(iii) *If M is a positive definite matrix, $\epsilon\kappa_2(\frac{M+M^T}{2}) = \eta < 1$, and*

$$\|\Delta M\|_2 \leq \epsilon \left\| \frac{M + M^T}{2} \right\|_2, \quad \|\Delta q\|_2 \leq \epsilon \max(\|(-q)_+\|_2, \|q\|_2 - \|Mx + q\|_2) \frac{\|M + M^T\|_2}{2\|M\|_2},$$

then $M + \Delta M$ is a positive matrix, and

$$\frac{\|x - y\|_2}{\|x\|_2} \leq \frac{2\epsilon}{1 - \eta} \kappa_2\left(\frac{M + M^T}{2}\right).$$

Remark 3.1. If $Mx + q = 0$, then (i) of Corollary 3.3 for M being an M-matrix and (ii) of Corollary 3.3 reduce to the perturbation bounds for the system of linear equations.

For the H-matrix LCP, componentwise perturbation bounds based on the Skeel condition number $\|\tilde{M}^{-1}\|\tilde{M}\|_{\infty}$ can be represented as follows.

Theorem 3.7 [5] *Suppose*

$$\begin{aligned} \min(x, Mx + q) &= 0 & M \in R^{n \times n}, \quad 0 \neq (-q)_+ \in R^n \\ \min(y, (M + \Delta M)y + q + \Delta q) &= 0 & \Delta M \in R^{n \times n}, \quad \Delta q \in R^n. \end{aligned}$$

with

$$|\Delta M| \leq \epsilon |M|, \quad |\Delta q| \leq \epsilon \max((-q)_+, |q| - |Mx + q|). \quad (3.1)$$

If M is an H-matrix with positive diagonals and $\epsilon \kappa_{\infty}(\tilde{M}) = \eta < 1$, then $M + \Delta M$ is an H-matrix with positive diagonals and

$$\frac{\|y - x\|_{\infty}}{\|x\|_{\infty}} \leq \frac{2\epsilon}{1 - \eta} \|\tilde{M}^{-1}\|\tilde{M}\|_{\infty}. \quad (3.2)$$

References

- [1] G.E. Alefeld, X. Chen and F.A. Potra, Numerical validation of solutions of linear complementarity problems, *Numer. Math.*, 83 (1999) 1-23.
- [2] A. Berman and R.J. Plemmons, *Nonnegative Matrix in the Mathematical Sciences*, SIAM Publisher, Philadelphia, 1994.
- [3] B. Chen, Error bounds for R_0 -type and monotone nonlinear complementarity problems, *J. Optim. Theory Appl.*, 108 (2001) 297-316.
- [4] Xiaojun Chen and Shuhuang Xiang, Computation of error bounds for P-matrix linear complementarity problems, *Mathematical Programming, Series A* (to appear).
- [5] X. Chen and S. Xiang, Perturbation bounds of P-matrix linear complementarity problems, Technical Report, Department of Mathematical Science, Hirosaki University, January, 2005.
- [6] R.W.Cottle, J.-S.Pang and R.E.Stone, *The Linear Complementarity Problem*, Academic Press, Boston, MA, 1992.

- [7] M. C. Ferris and O. L. Mangasarian, Error bounds and strong upper semicontinuity for monotone affine variational inequalities, *Ann. Oper. Res.*, 47(1993) 293-305.
- [8] Z.Q. Luo, J.-S.Pang and D. Ralph, *Mathematical Programs with Equilibrium Constraints*, Cambridge University Press, Cambridge, 1996.
- [9] O.L. Mangasarian and J. Ren, New improved error bounds for the linear complementarity problem, *Math. Programming* 66(1994) 241-257.
- [10] S.A.Gabriel and J.J.Moré, Smoothing of mixed complementarity problems, *Complementarity and Variational Problems: State of the Art*, eds by M.C.Ferris and J.-S.Pang, (SIAM Publications, Philadelphia, PA, 1997), 105-116.
- [11] R. Mathias and J.-S. Pang, Error bounds for the linear complementarity problem with a P-matrix, *Linear Algebra Appl.*, 132(1990) 123-136.
- [12] J.-S.Pang, Error bounds in mathematical programming, *Math. Programming*, 79(1997) 299-332.
- [13] J. Rohn, Systems of linear interval equations, *Linear Algebra Appl.*, 126(1989) 39-78.
- [14] U. Schäfer, A linear complementarity problem with a P-matrix, *SIAM Review*, 46(2004) 189-201.
- [15] N. Xiu and J.Zhang, A characteristic quantity of P-matrices, *App. Math. Lett.*, 15(2002) 41-46.