

THE REAL REPRESENTATION ASSOCIATED WITH COPRIME NORMAL SUBGROUPS

Masaharu Morimoto

Graduate School of Natural Science and Technology, Okayama University

Dedicated to Professor K. Shimakawa on his 60th birthday

Abstract. Let G be a finite group. In this article, we introduce a mutually coprime family of normal subgroups of G and the real G -module associated with the family, and we report interesting results on the real G -module.

1. PRELIMINARY

Throughout this paper, G is a finite group. We mean by a *real G -module* a real G -representation space of finite dimension. Let $\mathcal{S}(G)$ denote the set of all subgroups of G .

In the study of smooth G -actions on disks and spheres, there are important families of normal subgroups of G : for examples, $\{G\}$, $\{G^{\{2\}}\}$, $\{G^{\text{nil}}\}$,

$$\mathcal{K}(G) = \{G^{\{p\}} \mid p \text{ is a prime}\}, \text{ and}$$

$$\mathcal{N}_p(G) = \{H \trianglelefteq G \mid |G/H| = 1 \text{ or } p\},$$

where $G^{\{p\}}$ is the smallest normal subgroup H such that G/H has order of p -power (possibly $|G/H| = 1$), and G^{nil} is the smallest normal subgroup N such that G/N is nilpotent.

Let $\mathcal{L}$ be a set of subgroups of G such that each minimal element of $\mathcal{L}$ is a normal subgroup of G . Let $\mathbb{R}[G]$ denote the regular representation of G and let $\mathbb{R}[G]^{\mathcal{L}}$ denote the smallest G -submodule of $\mathbb{R}[G]$ containing all $\mathbb{R}[G]^L$ with $L \in \mathcal{L}$. Let $\mathbb{R}[G]_{\mathcal{L}}$ be

2010 *Mathematics Subject Classification.* Primary 20C15; Secondary 57S17.

Key words and phrases. real representation, gap condition, real projective space.

Partially supported by a Grant-in-Aid for Scientific Research (KAKENHI) no. 22540085.

the orthogonal complement of $\mathbb{R}[G]^\mathcal{L}$ in $\mathbb{R}[G]$ with respect to some G -invariant inner product on $\mathbb{R}[G]$, i.e.

$$\mathbb{R}[G]_\mathcal{L} = \mathbb{R}[G] - \mathbb{R}[G]^\mathcal{L}.$$

In this paper we call $\mathbb{R}[G]_\mathcal{L}$ the *real G -module associated with $\mathcal{L}$* .

Definition 1.1. A nonempty family $\mathcal{K}$ of normal subgroups of G is called *mutually coprime* if either

- (1) $\mathcal{K} = \{G\}$, or
- (2) $G \notin \mathcal{K}$ and $|G/K|$'s are mutually prime integers, i.e.

$$(|G/K|, |G/K'|) = 1 \text{ for all } K, K' \in \mathcal{K} \text{ such that } K \neq K'.$$

If $\mathcal{K}$ is a mutually coprime family of normal subgroups of G , then the equality

$$(1.1) \quad \mathbb{R}[G]_\mathcal{K} = (\mathbb{R}[G] - \mathbb{R}) - \bigoplus_{K \in \mathcal{K}} (\mathbb{R}[G/K] - \mathbb{R})$$

holds, where $\mathbb{R}$ is the 1-dimensional trivial real G -module.

Definition 1.2. Let $\mathcal{L}$ be a set of subgroups of G . Then we define the *upper closure* $\overline{\mathcal{L}}$ of $\mathcal{L}$ by

$$(1.2) \quad \overline{\mathcal{L}} = \{H \in \mathcal{S}(G) \mid H \supset L \text{ for some } L \in \mathcal{L}\},$$

and the *exterior* $\underline{\mathcal{L}}$ of $\mathcal{L}$ by

$$(1.3) \quad \underline{\mathcal{L}} = \mathcal{S}(G) \setminus \overline{\mathcal{L}}.$$

With this notation, we have $\mathcal{L}(G) = \overline{\mathcal{K}(G)}$ and $\mathcal{M}(G) = \underline{\mathcal{K}(G)}$, cf. E. Laitinen-M. Morimoto [1].

Definition 1.3. Let V be a real G -module and $\mathcal{H}$ a family of subgroups of G . We say that V is $\mathcal{H}$ -*complete* if for each $H \in \mathcal{H}$, any irreducible real H -module is isomorphic to a submodule of $\text{res}_H^G V$.

The main results which will be reported in this article are Theorems 2.1, 2.2 and 3.2. The proofs will appear somewhere else.

2. COMPLETENESS AND GAP PROPERTY

Let $\mathcal{K}$ be a mutually coprime family of normal subgroups of G . We introduce two practically important properties of $\mathbb{R}[G]_{\mathcal{K}}$ as the theorems below.

Theorem 2.1. *Let G be a finite group and let $\mathcal{K}$ be a mutually coprime family of normal subgroups of G . Then for any $H \in \underline{\mathcal{K}}$, $\text{res}_H^G \mathbb{R}[G]_{\mathcal{K}}$ contains a real H -submodule isomorphic to $\mathbb{R}[H]$. Hence the real G -module $\mathbb{R}[G]_{\mathcal{K}}$ is $\underline{\mathcal{K}}$ -complete.*

Theorem 2.2. *Let G be a finite group and let $\mathcal{K}$ be a mutually coprime family of normal subgroups of G . Then the real G -module $\mathbb{R}[G]_{\mathcal{K}}$ possesses the following properties.*

- (1) $\mathbb{R}[G]_{\mathcal{K}}^H \neq 0$ if and only if $H \in \underline{\mathcal{K}}$.
- (2) Let p be a prime and $H < K \leq G$ with $|K : H| = p$. Then

$$\dim \mathbb{R}[G]_{\mathcal{K}}^H \geq p \dim \mathbb{R}[G]_{\mathcal{K}}^K$$

holds; the equality holds if and only if there exists $K_k \in \mathcal{K}$ such that $p \mid |G : K_k|$, $|KK_k : HK_k| = p$, and $HK_i = G$ for all $K_i \in \mathcal{K} \setminus \{K_k\}$.

- (3) Let $H < K \leq G$. Then

$$\dim \mathbb{R}[G]_{\mathcal{K}}^H \geq 2 \dim \mathbb{R}[G]_{\mathcal{K}}^K$$

holds; the equality holds if and only if

- (a) $H \in \overline{\mathcal{K}}$, or
- (b) $K \in \underline{\mathcal{K}}$, $|K : H| = 2$, there exists $K_k \in \mathcal{K}$ such that $2 \mid |G : K_k|$, $|KK_k : HK_k| = 2$ and $HK_i = G$ for all $K_i \in \mathcal{K} \setminus \{K_k\}$.

The next proposition has been used in the induction argument of the equivariant surgery theory, cf. [1, 4, 5].

Proposition 2.3. *Let G be an Oliver group, and let P, H_1, H_2 be subgroups of G such that $P \in \mathcal{P}(G)$, $P < H_1$, and $P < H_2$. If the equality*

$$(2.1) \quad 2 \dim \mathbb{R}[G]_{\mathcal{L}(G)}^{H_i} = \dim \mathbb{R}[G]_{\mathcal{L}(G)}^P$$

holds for each $i = 1$ and 2 , then the smallest subgroup K containing H_1 and H_2 belongs to $\mathcal{M}(G) = \mathcal{S}(G) \setminus \mathcal{L}(G)$.

3. CANONICAL LINE BUNDLE OF REAL PROJECTIVE SPACE

Let V be a real G -module (of finite dimension). The real projective space $P(V)$ is the space of all 1-dimensional real vector subspaces of V , and $P(V)$ has the canonically induced G -action. Let γ_M , where $M = P(V)$, denote the canonical line bundle of M .

Lemma 3.1. *Let V be a real G -module and $M = P(V)$. Then the following equalities hold as real G -vector bundles via canonical isomorphisms.*

- (1) $\text{Hom}(\gamma_M, \gamma_M) = \varepsilon_M(\mathbb{R})$.
- (2) $\text{Hom}(\gamma_M, \varepsilon_M(\mathbb{R})) = \gamma_M$.
- (3) $T(M) = \text{Hom}(\gamma_M, \gamma_M^\perp)$.
- (4) $T(M) \oplus \varepsilon_M(\mathbb{R}) = \text{Hom}(\gamma_M, \varepsilon_M(V))$.
- (5) $\text{Hom}(\gamma_M, \varepsilon_M(V)) = \gamma_M \otimes V$.

The equalities (1)–(4) above follow from the proof of [3, Lemma 4.4]. The equality (5) holds because

$$\text{Hom}(\gamma_M, \varepsilon_M(V)) = \text{Hom}(\gamma_M, \varepsilon_M(\mathbb{R})) \otimes_{\mathbb{R}} V = \gamma_M \otimes_{\mathbb{R}} V.$$

Theorem 3.2. *Let $\mathcal{K}$ be a mutually coprime family of normal subgroups of G and let V be a real G -module such that $V = V^{\mathcal{K}}$. Then for $K_i \in \mathcal{K}$,*

$$(1) \quad P(V)^{K_i} = \begin{cases} P(V^{K_i}) & \text{if } 2 \mid |G : K_i| \\ P(V^{K_i}) \amalg \coprod_{L \in \mathcal{A}_i} P(V_{G/L}^L) & \text{if } 2 \nmid |G : K_i| \end{cases}$$

and

$$(2) \quad \gamma_{P(V)}|_{P(V)^{K_i}} = \begin{cases} \gamma_{P(V^{K_i})} & \text{if } 2 \mid |G : K_i| \\ \gamma_{P(V^{K_i})} \amalg \coprod_{L \in \mathcal{A}_i} \gamma_{P(V_{G/L}^L)} & \text{if } 2 \nmid |G : K_i|, \end{cases}$$

where $\mathcal{A}_i$ is the set of all subgroups L such that $|G : L| = 2$ and $|K_i : K_i \cap L| = 2$.

In addition

$$(3) \quad \begin{aligned} & (\gamma_{P(V)} \otimes_{\mathbb{R}} V)^{K_i} \\ &= \begin{cases} \gamma_{P(V^{K_i})} \otimes_{\mathbb{R}} V^{K_i} & \text{if } 2 \mid |G : K_i| \\ \gamma_{P(V^{K_i})} \otimes_{\mathbb{R}} V^{K_i} \amalg \coprod_{L \in \mathcal{A}_i} \gamma_{P(V_{G/L}^L)} \otimes_{\mathbb{R}} V_{G/L}^L & \text{if } 2 \nmid |G : K_i| \end{cases} \\ &= T(P(V)^{K_i}) \oplus \varepsilon_{P(V)^{K_i}}(\mathbb{R}). \end{aligned}$$

REFERENCES

- [1] E. Laitinen and M. Morimoto, *Finite groups with smooth one fixed point actions on spheres*, Forum Math. **10** (1998), 479–520.
- [2] E. Laitinen, M. Morimoto and K. Pawałowski, *Deleting-inserting theorem for smooth actions of finite nonsolvable groups on spheres*, Comment. Math. Helv. **70** (1995), 10–38.
- [3] J. W. Milnor and J. D. Stasheff, *Characteristic Classes*, Ann. Math. Stud. 76, Princeton Univ. Press, Princeton, New Jersey, 1974.
- [4] M. Morimoto, *The Burnside ring revisited*, in Current Trends in Transformation Groups, A. Bak, M. Morimoto and F. Ushitaki (eds.), *K-Monographs in Math.* 7, Kluwer Academic Publ., Dordrecht-Boston, 2002, 129–145.
- [5] M. Morimoto, *Deleting and inserting fixed point manifolds under the weak gap condition*, to appear in Publ. RIMS **48** (2012) No. 3.
- [6] M. Morimoto and K. Pawałowski, *The equivariant bundle subtraction theorem and its applications*, Fund. Math. **161** (1999), 279–303.
- [7] M. Morimoto and K. Pawałowski, *Smooth actions of Oliver groups on spheres*, Topology **42** (2003), 395–421.
- [8] B. Oliver, *Fixed point sets and tangent bundles of actions on disks and Euclidean spaces*, Topology **35** (1996), 583–615.
- [9] R. Oliver, *Fixed point sets of groups on finite acyclic complexes*, Comment. Math. Helv. **50** (1975), 155–177.
- [10] R. Oliver and T. Petrie, *G-CW surgery and $K(\mathbb{Z}[G])$* , Math. Z. **179** (1982), 11–42.
- [11] T. Petrie and J. Randall, *Transformation Groups on Manifolds*, Marcel Dekker, Inc., New York and Basel, 1984.

Graduate School of Natural Science and Technology
Okayama University
Tsushima-aka 3-1-1
Okayama, 700-8530 Japan
E-mail address: morimoto@ems.okayama-u.ac.jp