

Title:

**The mutual relationship between the host immune system and radiotherapy:
Stimulating the action of immune cells by irradiation**

Authors: Tsubasa Watanabe^{1,2}, Genki Edward Sato³, Michio Yoshimura³, Minoru Suzuki¹, Takashi
Mizowaki³

Affiliations:

¹ Institute for Integrated Radiation and Nuclear Science, Kyoto University

² The Hakubi Project, Kyoto University

³ Department of Radiation Oncology and Image-Applied Therapy, Graduate School of Medicine,
Kyoto University

*Corresponding: Tsubasa Watanabe, Institute for Integrated Radiation and Nuclear Science, Kyoto
University. E-mail: watanabe.tsubasa.8x@kyoto-u.ac.jp; Phone: +81-72-451-2407; FAX: +81-72-
451-2627

Address: 2, Asashiro-Nishi, Kumatori-cho, Sennan-gun, Osaka 590-0494 JAPAN

Abstract:

The effects of irradiation on tumor tissue and the host immune system are interrelated. The antitumor effect of irradiation is attenuated in the immunocompromised hosts. In addition, radiation alone positively and negatively influences the host immune system. The positive effects of radiation are summarized by the ability to help induce and enhance tumor-antigen-specific immune responses. The cancer-immunity cycle is a multistep framework that illustrates how the tumor-antigen-specific immune responses are induced and how the induced antigen-specific immune cells exert their functions in tumor tissues. Irradiation affects each step of this cancer-immunity cycle, primarily in a positive manner. In contrast, radiation also has negative effects on the immune system. The first is that irradiation has the possibility to kill irradiated effector immune cells. The second is that irradiation upregulates immunosuppressive molecules in the tumor microenvironment, whereas the third is that irradiation to the tumor condenses immunosuppressor cells in the tumor microenvironment. When used in conjunction with radiotherapy, immune checkpoint inhibitors can further leverage the positive effects of radiation on the immune system and compensate for the negative effects of irradiation, which supports the rationale for the combination of radiotherapy and immune checkpoint inhibitors. In this review, we summarize the preclinical evidence for the reciprocal effects of radiation exposure and the immune system, and up-front topics of the combination therapy of immune checkpoint inhibitors and radiotherapy.

36

37 Key words: radiotherapy, immune cells, radiosensitivity, cancer-immunity cycle.

38

Introduction

Immune checkpoints are inhibitory pathways that are crucial for maintaining self-tolerance by regulating immune activation and by modulating the T-cell response to self-proteins [1]. In the tumor microenvironment, the immune checkpoint mechanisms driven by molecules, such as cytotoxic T lymphocyte antigen 4 (CTLA-4), programmed cell death protein 1 (PD-1), and its ligand, programmed death-ligand 1 (PD-L1), are activated to suppress the antitumor immune responses. Deactivating the checkpoint mechanisms with immune checkpoint inhibitors (ICIs), anti-CTLA4 antibody (aCTLA4), and anti-PD-1 antibody (aPD-1)/anti-PD-L1 antibody (aPD-L1), has significantly improved cancer patients' prognosis. The combination of ICIs with existing cancer treatment modalities is currently being evaluated to further improve overall survival [2].

Radiotherapy (RT) is one of the most promising cancer therapies combined with ICIs. Chemoradiotherapy followed by consolidation ICIs improves the prognosis of advanced-stage non-small cell lung cancer (NSCLC) patients without metastases [3–5]. According to an updated report, the 4-year overall survival rate for chemoradiotherapy + aPD-L1 for advanced-stage NSCLC is 49.6%, which represents a tremendous improvement compared with 36.3% for chemoradiotherapy alone [5]. Thus, this treatment protocol was established as a standard treatment of care for advanced-stage NSCLC patients. Radiation can have both positive and negative effects on host immunity. On the other

hand, ICIs can mitigate the negative impact of radiation on immunity. In addition, some immune activation mechanisms of radiation are different and non-redundant from those of ICIs, resulting in acting synergistically with ICIs.

In this review, we first provide an overview of the types of immune cells associated with tumor shrinkage following irradiation exposure and the radiosensitivity of immune cells to discuss the negative impact of radiation on host immunity. Next, we summarize the positive impact of irradiation on the immune responses focusing on multiple steps of inducing antigen-specific immune responses. In addition to the direct cell-kill of immune cells after irradiation, there are some other negative effects of radiation on the host immune responses. We discuss the potential other negative impacts of radiation on host immune responses and the rationale for combining RT and ICIs with consideration of both positive and negative impacts of radiation on immune cells. Finally, as up-front topics, we will discuss the effect of ICIs on the radiosensitivity of immune cells and the challenges of combining current ICIs with other immune-modulating agents as a new treatment strategy to discuss the potential role of RT in the era of cancer immunotherapy.

T-cell immune responses contribute to the antitumor effect of irradiation

In the past, the reason for tumor shrinkage after irradiation was considered to be mediated by only the

direct cell-killing effect of irradiation through deoxyribonucleic acid (DNA) damage and the subsequent DNA damage responses. However, it has become clear that the host immune responses also contribute to tumor shrinkage after irradiation [6]. Even when RT is given to the tumor derived from the same tumor cell line under the same irradiation conditions, RT is less effective when tumors are implanted in immunocompromised nude mice than when they are implanted in immunocompetent mice [6]. Nude mice used for *in vivo* experiments lack a thymus and have immature T-cell functions. This result suggests that T-cell immunity is involved in maintaining tumor volume reduction after irradiation.

Among T-cells, the cluster of differentiation eight positive (CD8⁺) T-cells have been shown to have a powerful influence on the antitumor effect after irradiation [6–9]. In an *in vivo* mouse tumor model, simultaneous elimination of CD8⁺T-cells with irradiation significantly attenuates the antitumor effect of RT, and the irradiated tumor rapidly regrows a few days later even after a curative dose of irradiation. CD4⁺T-cell depletion is less effective for modifying the antitumor effect of radiation, showing a weak or no significant difference compared with RT alone [7, 10]. On the other hand, the selective removal of the forkhead box P3 positive (Foxp3⁺)CD25⁺CD4⁺regulatory T-cells (Tregs), which act as CD8⁺T-cell suppressors, enhances the effect of irradiation [11–13]. Immune cells other than T-cells have also been shown to influence the antitumor effects of RT. The removal of dendritic cells (DCs) from mice

using genetic manipulation abrogated the effects of RT [7, 9], while the removal of myeloid-derived suppressor cells (MDSCs) enhances the effects of RT [14, 15]. Considering DCs are involved in the activation process of antigen-specific CD8+T-cells, and MDSCs suppress CD8+T-cell function [16], the effect of removing those immune cells on RT depends upon subsequent effects on CD8+T-cells mainly. Taken together, these data suggest that the immune responses mediated by CD8+T-cells contribute to the antitumor effect of RT in addition to the direct cell-killing effect of irradiation.

Activated tumor-infiltrating lymphocytes are radioresistant compared with naïve lymphocytes

The radiosensitivity of immune cells depends on whether they are in an activated or naïve state as well as the type of immune cell. Classically, lymphocytes are believed to be equally highly radiosensitive without considering their condition and situation [17]. Actually, irradiation of peripheral blood, bone marrow, and lymphoid tissues is immunosuppressive as a number of the immune cells in these tissues die, even at low doses of irradiation [18, 19]. However, phytohemagglutinin-treated activated lymphocytes have been reported to be radioresistant [20, 21]. Another activator, anti-CD3/CD28 antibody, also makes T-cells radioresistant by downregulating the expression of ataxia-telangiectasia mutated (ATM) kinase, a major regulator of the cellular response to DNA double-strand breaks, which results in decreased ATM phosphorylation following irradiation [22].

Among the T-cell subsets, memory T-cells survive eight times more than naïve T-cells after irradiation and exhibit resistance to irradiation-induced apoptosis [23]. In this study, after 18 h of whole-body irradiation (6 Gy), the number of naïve T-cells in the spleen decreased from 4×10^6 to 1×10^5 cells (1/40), whereas memory T-cells decreased from 1.5×10^6 to 3×10^5 (1/5) in the lymphocytic choriomeningitis virus (LCMV)-immune mouse model [23]. In addition to memory T-cells, Arina *et al.* demonstrated that tumor-infiltrating T-cells are radioresistant [24]. They examined the effects of a transforming growth factor beta (TGF- β) blockade on the mortality of the tumor-infiltrating T-cells following irradiation. TGF- β blockade transformed the radioresistant T-cells into a radiosensitive phenotype, indicating that this signal plays a role in the radioresistance of the tumor-infiltrating T-cells.

Even if T-cells survive radiation exposure, it is of no use if the irradiated T-cells lose their function as effector immune cells in the tumor microenvironment. There is one report that examined the motility and function of the tumor-infiltrating T-cells following irradiation [24]. The authors used longitudinal *in vivo* imaging and discovered that irradiated pre-existing intratumoral T-cells maintained, or even rather increased, their motility as well as IFN γ production [24]. These results were surprising, but irradiated intratumoral T-cells have the possibility to maintain their motility and cytotoxic function after irradiation [24].

Effects of irradiation on each step of the cancer-immunity cycle

“The cancer-immunity cycle” is a multistep framework used to describe how the tumor-antigen-specific immune cells are activated as well as recognize and kill tumor cells (Fig.1) [25]. In the first step (step 1), the cancer-immunity cycle is initiated by regulated cell death known as immunogenic cell death (ICD). ICD is a form of stress-driven cell death that elicits sufficiently immune responses through the extracellular release of tumor-associated antigens and damage-associated molecular patterns (DAMPs), including extracellularly secreted adenosine-5'-triphosphate (ATP) and high-mobility group box 1 (HMGB1) as well as surface-exposed calreticulin by translocation, from the dying tumor cells in the tumor microenvironment [26]. Radiation is a bona fide ICD inducer [27] as is a consequence of the production of reactive oxygen species and endoplasmic reticulum stress in the irradiated tumor (Fig.1) [28, 29]. The next step (step 2) is the tumor-antigen presentation by DCs. In addition to the release of DAMPs and tumor antigens, irradiation induces DC maturation, as measured by the increased expression of costimulatory molecules, CD80/CD86 (ligands to CD28 or CTLA4 on T-cells) on DCs following irradiation [7]. The release of DAMPs and tumor antigens, and the maturation of DCs cooperatively enable the DCs to present tumor antigens to responsive T-cells as the next step of the cancer-immunity cycle.

The third step of the cancer-immunity cycle is the T-cell priming and activation phase. After step 1 and step 2, DCs transport from the tumor tissue to draining lymph nodes and serve as antigen-presenting cells to effector cells in the lymph nodes, by cross-presenting captured antigens to effector cells and priming them [9, 30]. The efficient initiation of the priming step requires type I interferon (IFN) [31, 32], and starts from the activation of cyclic GMP-AMP synthase (cGAS) and the stimulator of IFN genes (STING) pathway in both tumor cells and DCs [32–34]. Micronuclei are extra-nuclear bodies that form following DNA damage whenever a chromosome or its fragment is not incorporated into the daughter nuclei during cell division. Micronuclei and chromatin bridges, which are bridges formed between the separating groups of anaphase chromosomes accompanied by acentric chromosome fragments, are vital cGAS activators [35, 36]. Radiation-induced DNA damage forms micronuclei and chromatin bridges in the irradiated tumor cells, and activates cGAS and subsequent STING pathway in a dose-dependent manner, which results in the production of type I IFN (Fig.1) [34, 37]. After priming, the activated CD8+T-cells divide and proliferate. Thus, the degree of priming is estimated *in vivo* by monitoring the division of the CD8+T cells in the tumor-draining lymph nodes. By measuring the enhanced division of CD8+T-cells after irradiation in the draining lymph nodes, irradiation of the tumor is demonstrated to induce robust priming compared with unirradiated tumors (Fig.1) [6].

Activated antigen-specific CD8⁺T-cells in the draining lymph nodes return to the tumor microenvironment and infiltrate the tumor tissue. This represents the next steps of the cancer-immunity cycle and is known as the T-cell trafficking (step 4) and infiltration (step 5) phase. Radiation-induced chemokines and integrin ligand receptors cooperatively enhance T-cell trafficking at the irradiated tumor site [38]. Chemokine receptors, C-X-C chemokine receptor 3 (CXCR3) and CXCR6, are receptors for C-X-C chemokine ligand 9 (CXCL9)/CXCL10 and CXCL16, respectively. CXCR3 and CXCR6 are expressed at low levels on naïve T cells, but are upregulated upon activation [39, 40]. Radiation induces CXCL9/CXCL10 and CXCL16 expression in the tumor microenvironment in an IFN γ -dependent manner, which helps activated T-cells to traffic to the irradiated tumor (step 4) [41, 42]. Regarding T-cell infiltration to the tumor (step 5), irradiation induces the expression of integrin ligand receptors, intercellular adhesion molecule-1 (ICAM-1), and vascular cell adhesion protein-1 (VCAM-1), on the endothelium in the tumor microenvironment in a radiation-dose-dependent manner from 2 Gy up to 20 Gy (Fig.1) [43, 44]. The induced integrin ligand receptors of the T-cells assist in migration to the irradiated tumor and prevent recirculation to draining lymph nodes, which results in the enhanced localization of antigen-specific CD8⁺T-cells and condensation of them in the tumor tissue [45, 46].

The final steps of the cancer-immunity cycle are tumor cell recognition (step 6) and killing (step 7) by

immune cells. The recognition of tumor cells by the activated antigen-specific CD8⁺T-cells requires major histocompatibility complex I (MHC-I) expression on the tumor cells. The CD8⁺T-cells respond better to tumor cells expressing abundant peptide-MHC-I complexes than those with low or no MHC-I expression [47, 48]. Importantly, radiation induces MHC-I expression on the tumor cell surface in a dose-dependent manner and enhances tumor immunogenicity (Fig.1) [49–51]. Moreover, radiation enhances not only MHC-I expression but also both the peptide repertoire in the tumor cells and the expression level of tumor-associated antigens [49, 52, 53]. These make irradiated tumor cells more likely to be recognized by immune cells, and help tumor-antigen-specific CD8⁺ T-cells to kill tumor cells (Fig.1) [52, 53].

Synergistic effects of combining ICIs and RT

Among the multistep of the cancer-immunity cycle, immune checkpoint molecules suppress especially step 3 (priming) and step 7 (killing of targets) [25]. T-cell mediated antigen recognition requires the interaction of T-cell receptor (TCR) with MHC molecules and costimulatory signals by CD28 on the T-cells through CD80/CD86 molecules expressed on antigen-presenting DCs (Fig. 1). CTLA4 is a CD28 homolog with the higher binding affinity of CD80/CD86 and competitively suppresses the binding of CD80/CD86 to CD28 [54, 55]. CTLA4 expression on both DCs and Tregs deprives the effector T-cells' chance to get a sufficient costimulatory signal via CD28, leading to the attenuation of

the antigen-specific activation processes (step 3). In contrast to CTLA4, mainly expressed on immune cells, PD-1 ligands, PD-L1 and PD-L2, are expressed on various normal tissues [56, 57], and these tissues protect themselves against the potentially autoreactive effector T-cells by suppressing their activities with PD-1 ligands [58]. By expressing the PD-L1, cancer cells escape the effector cell-killing step of the cancer-immunity cycle by inducing the exhaustion of effector immune cells and attenuating the effector functions [59]. ICIs, both aCTLA4 and aPD-1/aPD-L1, reverse the inhibitory responses on the cancer-immunity cycle and enhance tumor-antigen-specific immune responses. Irradiation activates multistep of the cancer-immunity cycle, while ICIs deactivate immunosuppressive responses in the cancer-immunity cycle. This is one of the rationales for the combination of RT and ICIs.

In addition, RT enhances the antitumor reactivity of T-cells in a non-redundant way of ICIs' mechanisms of action, which also strengthens the rationale for combining ICIs with RT. The effective antigen-specific T-cell responses are induced only if the intratumoral TCR repertoire is intrinsically tumor-reactive and appropriate T-cell clones are ready to respond to the tumor [60], which is consistent with the clinical finding that the increased diversity of TCR of tumor-infiltrating T-cells is predictive for the efficacy of ICIs in metastatic melanoma patients [61]. Victor *et al.* demonstrated that irradiation to the tumor diversifies the TCR repertoire of tumor-infiltrating lymphocytes compared with unirradiated tumors [62]. This RT-induced diversification of the TCR repertoire helps to broaden the

window of the T-cell response [63]. In summary, radiation increases the diversity of the T-cell clones, while ICIs, if combined with RT, expand the RT-induced variety of T-cell clones, working synergistically to enhance the more robust immune responses [62, 64].

In contrast to these positive effects of RT, there are some negative effects on the immune system. Combining RT with ICIs compensates for the negative impact of radiation on the antitumor immune responses. First, irradiation induces the expression of PD-L1 on tumor cells [65]. Sato *et al.* demonstrated that tumor cells upregulate PD-L1 in response to DNA double-strand break in an ATM/ataxia telangiectasia and Rad3-related protein (ATR)/checkpoint kinase 1 (Chk1) pathway-dependent manner [66]. Other studies indicate that increased tumor PD-L1 expression and subsequent T-cell exhaustion cause resistance to RT + aCTLA combination therapy using a genetic PD-L1-deficient tumor [62]. The combination of aPD-1/aPD-L1 with RT reverses the effect of irradiation-induced PD-L1. The second negative effect of radiation on the antitumor immune responses involves Tregs. Several reports have shown that Tregs are more radioresistant than other lymphocytes, resulting in their selection after irradiation [67, 68]. Irradiated Tregs in the tumor are able to exert suppressive effects equally and are also able to proliferate in the tumor tissue even after irradiation [68]. CTLA-4 is expressed by Tregs constitutively. aCTLA4 drives the loss of Treg stability in the tumor microenvironment [69]. Targeting Tregs with aCTLA4 or other drugs is a strategy to reverse the Treg-

related negative impact on the immune responses following irradiation [70–72].

Up-front topics and future perspectives of the combination therapy of ICIs and RT

The sequence of combination therapy of RT plus ICIs (e.g., RT 1st & ICIs later or ICIs 1st & RT later) was one of the controversial issues that need to be resolved to maximize the combination effect [73].

As for an up-front topic, a recent report provides clues to a solution to the optimal order of the combination in relation to the radiosensitivity of immune cells [13]. To determine whether aPD1 prior to RT affects radiation-induced DNA damage of T-cells, the amount of fluorescent-stained phosphorylation of histone 2A family member X (γ H2AX), a marker for DNA damage, of CD8+T-cells was evaluated by flow cytometry. T-cells treated with aPD1 showed a significantly higher level of γ H2AX median fluorescence intensity than those with isotype control. In addition, treatment with aPD1 before irradiation induced more apoptosis of intratumor T-cells than isotype control. These results suggest that RT first protocol may be better in combination with RT and aPD1. To elucidate the underlying mechanism, the authors used an unsupervised hierarchical clustering algorithm, and found that aPD-1 administration before irradiation expanded the intratumor CD8+T-cell population, which resulted in a more naïve/non-activated phenotype in the tumor tissue [13]. This is consistent with the previous reports demonstrating that the administration of aPD1 not only reinvigorates exhausted

immune cells, but also induces the proliferation of CD8+T-cells [64, 74]. Naïve/non-activating CD8+T-cells are more radiosensitive compared with the activating ones, as mentioned above. The administration of aPD1 prior to RT results in an increased percentage of radiosensitive, unactivated intratumor CD8+T-cell populations, and the following RT may eradicate the radiosensitive populations expanded in the tumor tissue by aPD1. The potential survival superiority of “RT 1st & ICIs later protocol” was demonstrated in a retrospective analysis of patients with resected melanoma brain metastases [75]. Although further translational researches and prospective clinical trials are still needed, these findings may imply that aPD1/aPD-L1 administration has the potential to be better used following irradiation.

The effect of the double combination of RT and ICIs is still insufficient in clinical situations, and triple combination therapy of RT, ICIs, and the addition of immunomodulatory and radiosensitizing drugs is a promising future strategy. One such candidate agent is an indoleamine 2, 3-dioxygenases (IDO) inhibitor, which compensates for the negative impact of RT on host immune function and improves the efficacy of the double combination. IDO is a tryptophan catabolic enzyme that catalyzes the essential amino acid tryptophan into its immunosuppressive metabolite, kynurenine. Irradiation induces not only PD-L1 on the tumor, but also IDO overexpression in the tumor cells through type I and II IFN signaling [76]. The depletion of tryptophan and the increase in kynurenine in the

microenvironment impair the proliferation of effector immune cells, induce their apoptosis, and stimulate the differentiation process into Tregs [77]. Additionally, IDO inhibitors improve the radiosensitivity of tumor cells, and synergistic effects are observed when used in combination with RT [78]. Free heme and heme-derived iron show pro-oxidant activity in the irradiated sites. IDO contains heme [79], and IDO inhibitors release IDO-bound heme, which increases the concentration of free heme in the microenvironment and contributes to increased radiosensitivity [80]. With the aforementioned mechanisms of action, we hypothesized that the combination of IDO inhibitors and RT as well as aPD1 would have a synergistic effect by mitigating the negative impact of radiation. We showed that the combination of RT + IDO inhibitor (1-methyl-D-tryptophan) + aPD1 enhances the effect of the double combination of RT and aPD1 in preclinical models [81]. The combination of RT and IDO inhibitors (+ chemotherapy) is being tested in the clinical studies based on the compatibility of each (NCT04049669, NCT02052648). The treatments in these clinical trials are not combined with ICIs, leaving room for further combination with them if the clinical trials are successful. Although the result of a randomized phase III study (ECHO-301) of the combination of ICIs and an IDO inhibitor (without RT) was negative [82], the triple treatment strategy with ICIs + IDO inhibitors + RT would still be promising considering that RT has immune-activating effects that do not overlap with immunotherapy and IDO inhibitors can mitigate the radioresistance of tumor cells.

Conclusion

We summarized the preclinical data related to the interaction of irradiation and the immune responses mainly from the viewpoint of the radiosensitivity of immune cells, the rationale for combining ICIs with RT considering the positive and negative impacts of radiation on the immune system, and future perspectives of the combination strategy of RT + ICIs. The results of preclinical studies may help us to effectively combine ICIs with RT to further improve treatment response in clinical practice.

Acknowledgments

This work was supported in part by JSPS KAKENHI (Grant numbers 19K17231, 21K15805, T. Watanabe).

Compliance with ethical standards

Conflict of interest

The authors declare no potential conflicts of interest.

Figure legend

Fig.1 The effects of radiation on multistep of cancer-immunity cycle

The cancer-immunity cycle is a multistep framework to induce antigen-specific host immune response,

starting from (1) initiation by immunogenic cell death of tumor cells phase to the subsequent phases: (2) cancer antigen presentation by DC, (3) T-cell priming and activation by antigen-presenting cells, (4) T-cell trafficking from draining lymph nodes, (5) infiltration of T-cells to the tumor tissue, (6) T-cell recognition and (7) killing of tumors. Radiotherapy (RT) enhances each step of the cancer-immunity cycle and antigen-specific immune responses.

References

1. Wei SC, Duffy CR, Allison JP (2018) Fundamental Mechanisms of Immune Checkpoint Blockade Therapy. *Cancer Discov* 8:1069–1086
2. Morad G, Helmink BA, Sharma P, et al (2021) Hallmarks of response, resistance, and toxicity to immune checkpoint blockade. *Cell* 184:5309–5337
3. Antonia SJ, Villegas A, Daniel D, et al (2017) Durvalumab after Chemoradiotherapy in Stage III Non-Small-Cell Lung Cancer. *N Engl J Med* 377:1919–1929
4. Antonia SJ, Villegas A, Daniel D, et al (2018) Overall Survival with Durvalumab after Chemoradiotherapy in Stage III NSCLC. *N Engl J Med* 379:2342–2350
5. Faivre-Finn C, Vicente D, Kurata T, et al (2021) Four-Year Survival With Durvalumab After Chemoradiotherapy in Stage III NSCLC-an Update From the PACIFIC Trial. *J Thorac Oncol* 16:860–867

- 327 6. Lee Y, Auh SL, Wang Y, et al (2009) Therapeutic effects of ablative radiation on local tumor
328 require CD8⁺ T cells: changing strategies for cancer treatment. *Blood* 114:589–595
- 329 7. Gupta A, Probst HC, Vuong V, et al (2012) Radiotherapy Promotes Tumor-Specific Effector CD8
330 + T Cells via Dendritic Cell Activation. *J Immunol* 189:558–566
- 331 8. Chen J, Cao Y, Markelc B, et al (2019) Type I IFN protects cancer cells from CD8⁺ T cell-
332 mediated cytotoxicity after radiation. *J Clin Invest* 129:4224–4238
- 333 9. Blair TC, Bambina S, Alice AF, et al (2020) Dendritic Cell Maturation Defines Immunological
334 Responsiveness of Tumors to Radiation Therapy. *J Immunol* 204:3416–3424
- 335 10. De La Maza L, Wu M, Wu L, et al (2017) In Situ Vaccination after Accelerated Hypofractionated
336 Radiation and Surgery in a Mesothelioma Mouse Model. *Clin Cancer Res* 23:5502–5513
- 337 11. Bos PD, Plitas G, Rudra D, et al (2013) Transient regulatory T cell ablation deters oncogene-
338 driven breast cancer and enhances radiotherapy. *J Exp Med* 210:2435–2466
- 339 12. Kohno M, Murakami J, Wu L, et al (2020) Foxp3⁺ Regulatory T Cell Depletion after Nonablative
340 Oligofractionated Irradiation Boosts the Abscopal Effects in Murine Malignant Mesothelioma. *J*
341 *Immunol* 205:2519–2531
- 342 13. Wei J, Montalvo-Ortiz W, Yu L, et al (2021) Sequence of α PD-1 relative to local tumor irradiation
343 determines the induction of abscopal antitumor immune responses. *Sci Immunol* 6:eabg0117
- 344 14. Liang H, Deng L, Hou Y, et al (2017) Host STING-dependent MDSC mobilization drives

extrinsic radiation resistance. *Nat Commun* 8:1736

15. Jones KI, Tiersma J, Yuzhalin AE, et al (2018) Radiation combined with macrophage depletion promotes adaptive immunity and potentiates checkpoint blockade. *EMBO Mol Med* 10:e9342

16. Baumann T, Dunkel A, Schmid C, et al (2020) Regulatory myeloid cells paralyze T cells through cell-cell transfer of the metabolite methylglyoxal. *Nat Immunol* 21:555–566

17. Bergonié J, Tribondeau L (2003) Interpretation of some results from radiotherapy and an attempt to determine a rational treatment technique. 1906. *Yale J Biol Med* 76:181–182

18. Dunn PL, North RJ (1991) Selective radiation resistance of immunologically induced T cells as the basis for irradiation-induced T-cell-mediated regression of immunogenic tumor. *J Leukoc Biol* 49:388–396

19. Alexandru M, Rodica A, Dragos-Eugen G, et al (2021) Assessing the Spleen as an Organ at Risk in Radiation Therapy and Its Relationship With Radiation-Induced Lymphopenia: A Retrospective Study and Literature Review. *Adv Radiat Oncol* 6:100761

20. Schrek R, Stefani S (1964) Radioresistance of Phytohemagglutinin-Treated Normal and Leukemic Lymphocytes I, 2. *Journal of the National Cancer Institute* 32:507–521

21. Carloni M, Meschini R, Ovidi L, Palitti F (2001) PHA-induced cell proliferation rescues human peripheral blood lymphocytes from X-ray-induced apoptosis. *Mutagenesis* 16:115–120

22. Heylmann D, Badura J, Becker H, et al (2018) Sensitivity of CD3/CD28-stimulated versus non-

stimulated lymphocytes to ionizing radiation and genotoxic anticancer drugs: key role of ATM in the differential radiation response. *Cell Death Dis* 9:1053

23. Grayson JM, Harrington LE, Lanier JG, et al (2002) Differential sensitivity of naive and memory CD8⁺ T cells to apoptosis in vivo. *J Immunol* 169:3760–3770

24. Arina A, Beckett M, Fernandez C, et al (2019) Tumor-reprogrammed resident T cells resist radiation to control tumors. *Nat Commun* 10:3959

25. Chen DS, Mellman I (2013) Oncology meets immunology: the cancer-immunity cycle. *Immunity* 39:1–10

26. Galluzzi L, Vitale I, Warren S, et al (2020) Consensus guidelines for the definition, detection and interpretation of immunogenic cell death. *J Immunother Cancer* 8:e000337

27. Galluzzi L, Buqué A, Kepp O, et al (2017) Immunogenic cell death in cancer and infectious disease. *Nat Rev Immunol* 17:97–111

28. Hernandez C, Huebener P, Schwabe RF (2016) Damage-associated molecular patterns in cancer: a double-edged sword. *Oncogene* 35:5931–5941

29. Golden EB, Frances D, Pellicciotta I, et al (2014) Radiation fosters dose-dependent and chemotherapy-induced immunogenic cell death. *Oncoimmunology* 3:e28518

30. Gardner A, Ruffell B (2016) Dendritic Cells and Cancer Immunity. *Trends Immunol* 37:855–865

31. Ablasser A, Schmid-Burgk JL, Hemmerling I, et al (2013) Cell intrinsic immunity spreads to

bystander cells via the intercellular transfer of cGAMP. *Nature* 503:530–534

32. Fuertes MB, Kacha AK, Kline J, et al (2011) Host type I IFN signals are required for antitumor CD8⁺ T cell responses through CD8^α⁺ dendritic cells. *J Exp Med* 208:2005–2016

33. Deng L, Liang H, Xu M, et al (2014) STING-Dependent Cytosolic DNA Sensing Promotes Radiation-Induced Type I Interferon-Dependent Antitumor Immunity in Immunogenic Tumors. *Immunity* 41:843–852

34. Vanpouille-Box C, Alard A, Aryankalayil MJ, et al (2017) DNA exonuclease Trex1 regulates radiotherapy-induced tumour immunogenicity. *Nat Commun* 8:15618

35. Mackenzie KJ, Carroll P, Martin C-A, et al (2017) cGAS surveillance of micronuclei links genome instability to innate immunity. *Nature* 548:461–465

36. Flynn PJ, Koch PD, Mitchison TJ (2021) Chromatin bridges, not micronuclei, activate cGAS after drug-induced mitotic errors in human cells. *Proc Natl Acad Sci U S A* 118:e2103585118

37. Schröder-Heurich B, Wieland B, Lavin MF, et al (2014) Protective role of RAD50 on chromatin bridges during abnormal cytokinesis. *FASEB J* 28:1331–1341

38. Zhao Y, Zhang T, Wang Y, et al (2021) ICAM-1 orchestrates the abscopal effect of tumor radiotherapy. *Proc Natl Acad Sci U S A* 118:e2010333118

39. Harlin H, Meng Y, Peterson AC, et al (2009) Chemokine expression in melanoma metastases associated with CD8⁺ T-cell recruitment. *Cancer Res* 69:3077–3085

- 399 40. Matsumura S, Wang B, Kawashima N, et al (2008) Radiation-induced CXCL16 release by breast
400 cancer cells attracts effector T cells. *J Immunol* 181:3099–3107
- 401 41. Lugade AA, Sorensen EW, Gerber SA, et al (2008) Radiation-induced IFN-gamma production
402 within the tumor microenvironment influences antitumor immunity. *J Immunol* 180:3132–3139
- 403 42. Matsumura S, Demaria S (2010) Up-regulation of the pro-inflammatory chemokine CXCL16 is
404 a common response of tumor cells to ionizing radiation. *Radiat Res* 173:418–425
- 405 43. Lugade AA, Moran JP, Gerber SA, et al (2005) Local Radiation Therapy of B16 Melanoma
406 Tumors Increases the Generation of Tumor Antigen-Specific Effector Cells That Traffic to the
407 Tumor. *J Immunol* 174:7516–7523
- 408 44. Rodriguez-Ruiz ME, Garasa S, Rodriguez I, et al (2017) Intercellular Adhesion Molecule-1 and
409 Vascular Cell Adhesion Molecule Are Induced by Ionizing Radiation on Lymphatic Endothelium.
410 *Int J Radiat Oncol Biol Phys* 97:389–400
- 411 45. Slaney CY, Kershaw MH, Darcy PK (2014) Trafficking of T cells into tumors. *Cancer Res*
412 74:7168–7174
- 413 46. Yanguas A, Garasa S, Teijeira Á, et al (2018) ICAM-1-LFA-1 Dependent CD8⁺ T-Lymphocyte
414 Aggregation in Tumor Tissue Prevents Recirculation to Draining Lymph Nodes. *Front Immunol*
415 9:2084
- 416 47. Riaz N, Havel JJ, Makarov V, et al (2017) Tumor and Microenvironment Evolution during

417 Immunotherapy with Nivolumab. *Cell* 171:934-949

418 48. Benci JL, Johnson LR, Choa R, et al (2019) Opposing Functions of Interferon Coordinate
419 Adaptive and Innate Immune Responses to Cancer Immune Checkpoint Blockade. *Cell* 178:933-
420 948

421 49. Reits EA, Hodge JW, Herberts CA, et al (2006) Radiation modulates the peptide repertoire,
422 enhances MHC class I expression, and induces successful antitumor immunotherapy. *J Exp Med*
423 203:1259–1271

424 50. Sharabi AB, Nirschl CJ, Kochel CM, et al (2015) Stereotactic Radiation Therapy Augments
425 Antigen-Specific PD-1-Mediated Antitumor Immune Responses via Cross-Presentation of
426 Tumor Antigen. *Cancer Immunol Res* 3:345–355

427 51. Zebertavage LK, Alice A, Crittenden MR, et al (2020) Transcriptional Upregulation of NLRC5
428 by Radiation Drives STING- and Interferon-Independent MHC-I Expression on Cancer Cells and
429 T Cell Cytotoxicity. *Sci Rep* 10:7376

430 52. Lhuillier C, Rudqvist N-P, Yamazaki T, et al (2021) Radiotherapy-exposed CD8⁺ and CD4⁺
431 neoantigens enhance tumor control. *J Clin Invest* 131:e138740

432 53. Lussier DM, Alspach E, Ward JP, et al (2021) Radiation-induced neoantigens broaden the
433 immunotherapeutic window of cancers with low mutational loads. *Proc Natl Acad Sci U S A*
434 118:e2102611118

- 435 54. van der Merwe PA, Bodian DL, Daenke S, et al (1997) CD80 (B7-1) binds both CD28 and CTLA-
436 4 with a low affinity and very fast kinetics. *J Exp Med* 185:393–403
- 437 55. Buchbinder EI, Desai A (2016) CTLA-4 and PD-1 Pathways: Similarities, Differences, and
438 Implications of Their Inhibition. *Am J Clin Oncol* 39:98–106
- 439 56. Freeman GJ, Long AJ, Iwai Y, et al (2000) Engagement of the PD-1 immunoinhibitory receptor
440 by a novel B7 family member leads to negative regulation of lymphocyte activation. *J Exp Med*
441 192:1027–1034
- 442 57. Latchman Y, Wood CR, Chernova T, et al (2001) PD-L2 is a second ligand for PD-1 and inhibits
443 T cell activation. *Nat Immunol* 2:261–268
- 444 58. Keir ME, Liang SC, Guleria I, et al (2006) Tissue expression of PD-L1 mediates peripheral T
445 cell tolerance. *J Exp Med* 203:883–895
- 446 59. Iwai Y, Ishida M, Tanaka Y, et al (2002) Involvement of PD-L1 on tumor cells in the escape from
447 host immune system and tumor immunotherapy by PD-L1 blockade. *Proc Natl Acad Sci U S A*
448 99:12293–12297
- 449 60. Scheper W, Kelderman S, Fanchi LF, et al (2019) Low and variable tumor reactivity of the
450 intratumoral TCR repertoire in human cancers. *Nat Med* 25:89–94
- 451 61. Valpione S, Mundra PA, Galvani E, et al (2021) The T cell receptor repertoire of tumor infiltrating
452 T cells is predictive and prognostic for cancer survival. *Nat Commun* 12:4098

- 453 62. Twyman-Saint Victor C, Rech AJ, Maity A, et al (2015) Radiation and dual checkpoint blockade
454 activate non-redundant immune mechanisms in cancer. *Nature* 520:373–377
- 455 63. Rudqvist N-P, Pilonis KA, Lhuillier C, et al (2018) Radiotherapy and CTLA-4 Blockade Shape
456 the TCR Repertoire of Tumor-Infiltrating T Cells. *Cancer Immunol Res* 6:139–150
- 457 64. Im SJ, Hashimoto M, Gerner MY, et al (2016) Defining CD8⁺ T cells that provide the
458 proliferative burst after PD-1 therapy. *Nature* 537:417–421
- 459 65. Sato H, Okonogi N, Nakano T (2020) Rationale of combination of anti-PD-1/PD-L1 antibody
460 therapy and radiotherapy for cancer treatment. *Int J Clin Oncol* 25:801–809
- 461 66. Sato H, Niimi A, Yasuhara T, et al (2017) DNA double-strand break repair pathway regulates PD-
462 L1 expression in cancer cells. *Nat Commun* 8:1751
- 463 67. Kachikwu EL, Iwamoto KS, Liao Y-P, et al (2011) Radiation Enhances Regulatory T Cell
464 Representation. *Int J Radiat Oncol Biol Phys* 81:1128–1135
- 465 68. Muroyama Y, Nirschl TR, Kochel CM, et al (2017) Stereotactic Radiotherapy Increases
466 Functionally Suppressive Regulatory T Cells in the Tumor Microenvironment. *Cancer Immunol*
467 *Res* 5:992–1004
- 468 69. Zappasodi R, Serganova I, Cohen JJ, et al (2021) CTLA-4 blockade drives loss of Treg stability
469 in glycolysis-low tumours. *Nature* 591:652–658
- 470 70. Demaria S, Kawashima N, Yang AM, et al (2005) Immune-mediated inhibition of metastases

after treatment with local radiation and CTLA-4 blockade in a mouse model of breast cancer.

Clin Cancer Res 11:728–734

71. Formenti SC, Rudqvist N-P, Golden E, et al (2018) Radiotherapy induces responses of lung cancer to CTLA-4 blockade. Nat Med 24:1845–1851

72. Sia J, Hagekyriakou J, Chindris I, et al (2021) Regulatory T Cells Shape the Differential Impact of Radiation Dose-Fractionation Schedules on Host Innate and Adaptive Antitumor Immune Defenses. Int J Radiat Oncol Biol Phys 111:502–514

73. Gunderson AJ, Young KH (2018) Exploring optimal sequencing of radiation and immunotherapy combinations. Adv Radiat Oncol 3:494–505

74. Siddiqui I, Schaeuble K, Chennupati V, et al (2019) Intratumoral Tcf1+PD-1+CD8+ T Cells with Stem-like Properties Promote Tumor Control in Response to Vaccination and Checkpoint Blockade Immunotherapy. Immunity 50:195–211

75. Pomeranz Krummel DA, Nasti TH, Izar B, et al (2020) Impact of Sequencing Radiation Therapy and Immune Checkpoint Inhibitors in the Treatment of Melanoma Brain Metastases. Int J Radiat Oncol Biol Phys 108:157–163

76. Munn DH, Mellor AL (2007) Indoleamine 2,3-dioxygenase and tumor-induced tolerance. J Clin Invest 117:1147–1154

77. Metz R, Rust S, Duhadaway JB, et al (2012) IDO inhibits a tryptophan sufficiency signal that

stimulates mTOR: A novel IDO effector pathway targeted by D-1-methyl-tryptophan.
 Oncoimmunology 1:1460–1468

78. Chen B, Alvarado DM, Iticovici M, et al (2020) Interferon-Induced IDO1 Mediates Radiation
 Resistance and Is a Therapeutic Target in Colorectal Cancer. *Cancer Immunol Res* 8:451–464

79. Thomas SR, Stocker R (1999) Redox reactions related to indoleamine 2,3-dioxygenase and
 tryptophan metabolism along the kynurenine pathway. *Redox Rep* 4:199–220

80. Ju J-M, Nam G, Lee Y-K, et al (2021) IDO1 scavenges reactive oxygen species in myeloid-
 derived suppressor cells to prevent graft-versus-host disease. *Proc Natl Acad Sci U S A*
 118:e2011170118

81. Watanabe T, Gaedicke S, Guffart E, et al (2020) Adding Indoximod to Hypofractionated
 Radiotherapy with Anti-PD-1 Checkpoint Blockade Enhances Early NK and CD8+ T-Cell–
 Dependent Tumor Activity Effects of Indoximod on Radiotherapy plus Anti-PD-1. *Clin Cancer*
Res 26:945–956

82. Van den Eynde BJ, van Baren N, Baurain J-F (2020) Is There a Clinical Future for IDO1
 Inhibitors After the Failure of Epacadostat in Melanoma? *Annu Rev Cancer Biol* 4:241–256

Tumor microenvironment

Radiotherapy (RT)

CD8+T-cell proliferation

[step 6, 7] RT enhances tumor immunogenicity and helps antigen-specific T-cells recognize and kill tumor cells.

RT-induced peptide- MHC I complex antigen-specific T-cells

[step 1] RT-induced immunogenic cell death

[step 4, 5] RT enhances T-cell trafficking & infiltration

Endothelium

ICAM-1
Integrin

[step 3] RT enhances priming of antigen-specific T cells

[step 2] RT-induced DC maturation and antigen release from tumors

cGAS-STING pathway

Tumor-antigen, DAMPs, IFNs

Irradiated site

Tumor-draining lymph nodes



Tumor cell



Dead tumor cell



Dendritic cell



CD8+T-cell



Immune suppressor cell



Activated cells