

実 2 次変換による Cuntz 環 $\mathcal{O}_2$ の表現
Representations of the Cuntz algebra $\mathcal{O}_2$
arising from real quadratic transformations

京大・数理解 川村 勝紀 (Katsunori Kawamura)
Research Institute for Mathematical Sciences,
Kyoto Univ.

Abstract

We construct representations of the Cuntz algebra $\mathcal{O}_2$ from real quadratic transformations of 1-dimensional dynamical systems. By intertwining relations among transformations, we derive relations among operators associated with representations of $\mathcal{O}_2$. From these relations, we show properties of these representations.

1 Introduction

In the theory of dynamical system, 1-dimensional dynamical systems of real quadratic functions on closed intervals are well known. In [?], such system is introduced as a typical example of chaotic dynamical system. The key point of this orbit analysis in [?] is some *intertwining relation* between a quadratic transformation and some piecewise linear transformation *but not conjugation between them*.

On the other hand, our interests are the construction of representations of Cuntz algebras and their analysis([?, ?, ?]). For these intentions, we construct a representation from a 1-dimensional dynamical system by a quadratic transformation as follows: Consider a real quadratic transformation Q on a closed interval $[-2, 2]$ by

$$Q : [-2, 2] \rightarrow [-2, 2]; \quad Q(x) \equiv x^2 - 2. \quad (1.1)$$

We define a representation $(L_2[-2, 2], \pi_q)$ of the Cuntz algebra $\mathcal{O}_2$ arising from Q by

$$(\pi_q(s_i)\phi)(x) = m_i(x)\phi(Q(x)) \quad (i = 1, 2, \phi \in L_2[-2, 2]) \quad (1.2)$$

where $m_i(x) = \chi_{D_i}(x)\sqrt{2|x|}$, χ_{D_i} is the characteristic function on D_i , $i = 1, 2$, $D_1 = [0, 2]$, $D_2 = [-2, 0]$ and s_1, s_2 are generators of $\mathcal{O}_2$.

Theorem 1.1 *Let $(L_2[-2, 2], \pi_q)$ be the representation of $\mathcal{O}_2$ defined in (??).*

(i) *Put a function*

$$\Omega(x) \equiv \frac{1}{\sqrt{\pi}} \frac{1}{(4 - x^2)^{1/4}} \quad (x \in [-2, 2]).$$

Then Ω is unique eigen vector in $L_2[-2, 2]$ of an operator $\pi_q(s_1) + \pi_q(s_2)$ up to scalar multiples and its eigen value is $\sqrt{2}$.

(ii) $(L_2[-2, 2], \pi_q)$ is unitarily equivalent to one of the GP representation of $\mathcal{O}_2$ in [?].

(iii) $(L_2[-2, 2], \pi_q)$ is irreducible.

We introduce several representations of Cuntz algebras in section ?? . In section ??, we show intertwining relations of transformations of dynamical systems and derive relations of operators from them. By using these relations, we show Theorem ??.

2 Representations of the Cuntz algebra

In this section we review some basic facts on the Cuntz algebras and their representations on measure spaces. For $N \geq 2$, a C^* -algebra with generators $s_1, \dots, s_N$ which satisfy the following relations([?])

$$s_i^* s_j = \delta_{ij} I \quad (i, j = 1, \dots, N), \quad \sum_{i=1}^N s_i s_i^* = I \quad (2.1)$$

is called the *Cuntz algebra* and is denoted by $\mathcal{O}_N$. $\mathcal{O}_N$ is non commutative, infinite dimensional, separable, simple and unique up to isomorphisms. Hence there is no finite dimensional representation of $\mathcal{O}_N$ except 0-representation.

2.1 GP representations of $\mathcal{O}_N$ with 1-cycle

We show only the 1-cycle case about GP representations of Cuntz algebras with cycle in [?]. The GP representation is used to characterize representation introduced in § ?? . In this paper, a representation always means a unital $*$ -representation on a complex Hilbert space.

Let $S(\mathbb{C}^N) \equiv \{z \in \mathbb{C}^N : \|z\| = 1\}$ be the complex sphere in a complex vector space $\mathbb{C}^N$.

Definition 2.1 $(\mathcal{H}, \pi, \Omega)$ is the GP(= generalized permutative) representation of $\mathcal{O}_N$ by $z = (z_1, \dots, z_N) \in S(\mathbb{C}^N)$ if $(\mathcal{H}, \pi)$ is a cyclic representation of $\mathcal{O}_N$ and Ω is the unit cyclic vector which satisfies the following equation:

$$\pi(z_1 s_1 + \dots + z_N s_N) \Omega = \Omega. \quad (2.2)$$

We denote $GP(z) \equiv (\mathcal{H}, \pi, \Omega)$.

For two representations $(\mathcal{H}_1, \pi_1)$ and $(\mathcal{H}_2, \pi_2)$ of $\mathcal{O}_N$, $(\mathcal{H}_1, \pi_1) \sim (\mathcal{H}_2, \pi_2)$ means the unitary equivalence between $(\mathcal{H}_1, \pi_1)$ and $(\mathcal{H}_2, \pi_2)$.

Theorem 2.2 (Properties of GP representation with 1-cycle)

(i) (Existence and Uniqueness) For any $z \in S(\mathbb{C}^N)$, $GP(z)$ exists uniquely up to unitary equivalence.

- (ii) (Irreducibility) For any $z \in S(\mathbf{C}^N)$, $GP(z)$ is irreducible.
- (iii) (Equivalence) For any $z, z' \in S(\mathbf{C}^N)$, $GP(z) \sim GP(z')$ if and only if $z = z'$.
- (iv) (Uniqueness of eigen vector) For $z \in S(\mathbf{C}^N)$, Ω in (??) is unique up to scalar multiples.

Proof. (i) Proposition 3.4 and Proposition 5.4 in [?]. (ii), (iii) Example 6.3 (ii) in [?]. (iv) Corollary 5.3 in [?]. ■

We show examples of GP representation. Let $l_2(\mathbf{N})$ be a Hilbert space with the canonical basis $\{e_n : n \in \mathbf{N}\}$ where $\mathbf{N} = \{1, 2, 3, \dots\}$. We make the following representation $(l_2(\mathbf{N}), \pi_S)$ of the Cuntz algebra $\mathcal{O}_N$ which is called *the standard representation of $\mathcal{O}_N$* :

$$\pi_S(s_i)e_n \equiv e_{N(n-1)+i} \quad (i = 1, \dots, N, n \in \mathbf{N}). \quad (2.3)$$

From this definition, we have $\pi_S(s_i)^*e_{N(n-1)+j} = \delta_{ij}e_n$ for $i, j = 1, \dots, N$ and $n \in \mathbf{N}$. Note that $(l_2(\mathbf{N}), \pi_S)$ is one of permutative representations of $\mathcal{O}_N$ by [?]. Since $\pi_S(s_1)e_1 = e_1$, $(l_2(\mathbf{N}), \pi_S, e_1)$ is $GP(z)$ for $z = (1, 0, \dots, 0) \in S(\mathbf{C}^N)$ and irreducible.

More about details for GP representations, see [?, ?, ?, ?].

2.2 Isometries arising from transformations on measure spaces

By simplicity and uniqueness of $\mathcal{O}_N$, it is sufficient to define operators $S_1, \dots, S_N$ on an infinite dimensional Hilbert space which satisfy (??) in order to construct a representation of $\mathcal{O}_N$.

We introduce an easy method to construct partial isometries from maps on a measure space.

Let (X, μ) be a measure space and $Y \subset X$ a measurable subset of X .

Definition 2.3 (i) $RN(Y, X)$ is the set of measurable maps on Y defined by

$$RN(Y, X) \equiv \{f : Y \rightarrow X \mid \text{there exists } \Phi_f \text{ and } \Phi_f > 0 \text{ a.e. } Y\}$$

where Φ_f is the Radon-Nikodým derivative of $\mu \circ f$ with respect to μ on Y .

- (ii) $RN_{loc}(X) \equiv \bigcup_{Y \subset X} RN(Y, X)$ where Y is taken from all measurable subsets of X .

For $f, g \in RN_{loc}(X)$, we denote the domain and the range of f by $D(f)$ and $R(f)$, respectively and $(f \circ g)(x) \equiv f(g(x))$ for $x \in D(g)$ when $D(f) \supset R(g)$.

Lemma 2.4 (i) If $f \in RN_{loc}(X)$, then $f^{-1} \in RN_{loc}(X)$.

- (ii) For $f, g \in RN_{loc}(X)$, $f \circ g \in RN_{loc}(X)$ when $D(f) \supset R(g)$.

- (iii) $\Phi_{f \circ g} = \Phi_g \cdot ((\Phi_f) \circ g)$.

Proof. These are easily checked by direct computation and property of Radon-Nikodým derivative. ■

Note that $RN_{loc}(X)$ is a groupoid by Lemma ?? (ii).

Definition 2.5 For $f \in RN_{loc}(X)$ define an operator $S(f) : L_2(X, \mu) \rightarrow L_2(X, \mu)$ by

$$(S(f)\phi)(x) \equiv \begin{cases} \{\Phi_f(f^{-1}(x))\}^{-1/2} \phi(f^{-1}(x)) & (x \in R(f)), \\ 0 & (\text{otherwise}) \end{cases} \quad (2.4)$$

for $\phi \in L_2(X, \mu)$ and $x \in X$.

We simply denote

$$S(f) = \mathcal{L}_f M_{\Phi_f^{-1/2}} \quad (2.5)$$

where M_g is the multiplication operator of $g \in L_\infty(X, \mu)$ defined by

$$(\mathcal{L}_f \phi)(x) \equiv \chi_{R(f)}(x) \phi(f^{-1}(x)) \quad (x \in X)$$

and χ_Y is the characteristic function on $Y \subset X$.

Lemma 2.6 (i) For $f \in RN_{loc}(X)$, $S(f)$ is a partial isometry on $L_2(X, \mu)$ with the initial projection $M_{\chi_{D(f)}}$ and the range projection $M_{\chi_{R(f)}}$.

(ii) For $f \in RN_{loc}(X)$, $S(f)^* = S(f^{-1})$.

(iii) $S(id_Y) = M_{\chi_Y}$.

(iv) For $f, g \in RN_{loc}(X)$, $\mathcal{L}_f \mathcal{L}_g = \mathcal{L}_{f \circ g}$ when $D(f) \supset R(g)$.

(v) For $f \in RN_{loc}(X)$ and $g \in L_\infty(X)$, $\mathcal{L}_f M_g = M_{\chi_{R(f)}} M_{g \circ f^{-1}} \mathcal{L}_f$.

Proof. (i), (iv) and (v) follow by simple computation. (ii) Since $R(f^{-1}) = D(f)$, $D(f^{-1}) = R(f)$ and the property of Radon-Nikodým derivative, it follows. (iii) Since $\Phi_{id_Y} = \chi_Y$, it follows. ■

Let $\text{PIso}(L_2(X, \mu))$ be the groupoid of partial isometries on $L_2(X, \mu)$ by usual product of operators.

Lemma 2.7 A map $S : RN_{loc}(X) \rightarrow \text{PIso}(L_2(X, \mu))$ is a groupoid homomorphism, that is,

$$S(f)S(g) = S(f \circ g) \quad (2.6)$$

when $f, g \in RN_{loc}(X)$ and $D(f) \supset R(g)$.

Proof. By Lemma ?? (iii) and Lemma ?? (iv),(v),

$$S(f)S(g) = \mathcal{L}_f M_{\Phi_f^{-1/2}} \mathcal{L}_g M_{\Phi_g^{-1/2}} = \mathcal{L}_{f \circ g} M_{((\Phi_f) \circ g)^{-1/2}} M_{\Phi_g^{-1/2}} = S(f \circ g).$$

Remark that $f \circ g$ in rhs of (??) is only the composition of two maps f and g but not special product of them. ■

2.3 Representations of Cuntz algebra on a measure space

The notion of branching function system was introduced in [?] in order to construct a representation of $\mathcal{O}_N$ from a family of maps. We introduce a branching function system to define the representation from a dynamical system in section ??.

Let $N \geq 2$ and (X, μ) a measure space.

Definition 2.8 $f = \{f_i\}_{i=1}^N$ is a branching function system over (X, μ) if $f_i \in RN_{loc}(X)$ and $D(f_i) = X$, put $R_i \equiv R(f_i)$, then $\mu(R_i \cap R_j) = 0$, $i \neq j$ and $\mu(X \setminus \bigcup_{i=1}^N R_i) = 0$.

Proposition 2.9 For a branching function system $f = \{f_i\}_{i=1}^N$ on (X, μ) ,

$$\pi_f(s_i) \equiv S(f_i) \quad (i = 1, \dots, N),$$

defines a representation $(L_2(X, \mu), \pi_f)$ of $\mathcal{O}_N$ where S is the map in Definition ??.

Proof. It is straightforward to show that $S(f_1), \dots, S(f_N)$ satisfy the relations (??) by Lemma ??, Lemma ?? and Definition ??. ■

We show several examples of branching function systems associated with 1-dimensional dynamical systems in [?].

Lemma 2.10 Let (Y, ν) be another measure space. Assume that $\varphi : X \rightarrow Y$ is a measurable bijection a.e. X and Y and its Radon-Nikodým derivative Φ_φ is positive a.e. X .

- (i) An operator $U_\varphi : L_2(X, \mu) \rightarrow L_2(Y, \nu)$ defined by $U_\varphi \equiv \mathcal{L}_\varphi M_{\Phi_\varphi^{1/2}}$ is a unitary.
- (ii) For $f \in RN_{loc}(X)$, $\varphi \circ f \circ \varphi^{-1} \in RN_{loc}(Y)$ and $U_\varphi S(f) U_\varphi^* = S(\varphi \circ f \circ \varphi^{-1})$.
- (iii) If $f = \{f_i\}_{i=1}^N$ is a branching function system over (X, μ) , then $\{\varphi \circ f_i \circ \varphi^{-1}\}_{i=1}^N$ is a branching function system over (Y, ν) , too.

Proof. These are easily proved by direct computation. ■

Proposition 2.11 Let $f = \{f_i\}_{i=1}^N$ and $g = \{g_i\}_{i=1}^N$ be branching function systems over measure spaces (X, μ) and (Y, ν) , respectively. Assume that there is a map $\varphi : X \rightarrow Y$ which satisfies the assumption in Lemma ?? and a map identity $g_i = \varphi \circ f_i \circ \varphi^{-1}$ holds for $i = 1, \dots, N$. Then $(L_2(X, \mu), \pi_f)$ and $(L_2(Y, \nu), \pi_g)$ are unitarily equivalent.

Proof. By Lemma ?? (ii), we can show $S(g_i) = U_\varphi S(f_i) U_\varphi^*$ for $i = 1, \dots, N$. These relations induce a unitary equivalence between π_f and π_g in Proposition ?? immediately. ■

3 Operator relations arising from intertwiner among transformations in dynamical systems

Under the preparation in section ??, we show the property of representation of $\mathcal{O}_2$ arising from a quadratic transformation. The first, we consider intertwining relations among transformations of dynamical systems. Next we derive relations of operator on a Hilbert space which are associated with representations of $\mathcal{O}_2$. The properties (??) of the map S plays important role in this procedure.

3.1 Intertwining relations of transformations

For study of dynamical system, the strongest equivalence relation between two systems is the conjugation([?]). For two dynamical systems (X, F) and (Y, G) are *conjugate* if there is a bijective map $\varphi : X \rightarrow Y$ such that $G = \varphi \circ F \circ \varphi^{-1}$ with some conditions (continuity, measurability, conservation of measure, differentiability and so on) of φ depending on the aim of study and class of dynamical systems. φ is called a *conjugacy*.

In spite that we do not know the existence of conjugation map between two dynamical systems, if we have an intertwiner $\varphi : X \rightarrow Y$ between F and G , that is, $G \circ \varphi = \varphi \circ F$, (where φ is not always injective), then it is useful to consider relations between (X, F) and (Y, G) in some situation. φ is called a *semiconjugacy*([?]).

In order to introduce intertwining relations of transformations associated with Q , we prepare two transformations except Q in (??) according to § 10.2 in [?]. Put two transformations

$$V, C : [-2, 2] \rightarrow [-2, 2]; \quad V(x) \equiv 2|x| - 2, \quad C(x) \equiv -2 \cos \frac{\pi}{2} x, \quad (3.1)$$

and $D_1 \equiv [0, 2]$, $D_2 \equiv [-2, 0]$. Then every restriction $V|_{D_i}, C|_{D_i}, Q|_{D_i} : D_i \rightarrow [-2, 2]$ is invertible for each $i = 1, 2$. Hence we can take inverse transformations $v_i, c_i, q_i : [-2, 2] \rightarrow D_i$ by

$$v_i \equiv (V|_{D_i})^{-1}, \quad c_i \equiv (C|_{D_i})^{-1}, \quad q_i \equiv (Q|_{D_i})^{-1} \quad (3.2)$$

for each $i = 1, 2$. Concretely we have the followings:

$$\begin{cases} v_1(x) = \frac{1}{2}x + 1, & \begin{cases} c_1(x) = \frac{2}{\pi} \arccos(-\frac{1}{2}x), \\ c_2(x) = -\frac{2}{\pi} \arccos(-\frac{1}{2}x), \end{cases} & \begin{cases} q_1(x) = \sqrt{x+2}, \\ q_2(x) = -\sqrt{x+2}. \end{cases} \end{cases}$$

Lemma 3.1 $\{v_1, v_2\}, \{c_1, c_2\}, \{q_1, q_2\}$ are branching function systems on $[-2, 2]$ for the case $N = 2$ with respect to Lebesgue measure in Definition ??.

Proof. Note

$$\Phi_{v_i}(x) = \frac{1}{2}, \quad \Phi_{c_i}(x) = \frac{2}{\pi} \frac{1}{\sqrt{4-x^2}}, \quad \Phi_{q_i}(x) = \frac{1}{2} \frac{1}{\sqrt{x+2}} \quad (3.3)$$

for $i = 1, 2$ and $x \in [-2, 2]$. Hence every Radon-Nikodým derivative of v_i, c_i, q_i is positive a.e. $[-2, 2]$. Other conditions in Definition ?? are easily checked. ■

Remark that the following intertwining relation holds on $[-2, 2]$:

$$C \circ V = Q \circ C. \quad (3.4)$$

(??) follows immediately from the double angle's formula and periodicity of cosine function. C is a semiconjugacy between V and Q but not a conjugacy because C is not injective on $[-2, 2]$. We derive several identities among v_i, c_i, q_i by dividing the domain of (??).


Let R_i be a subinterval of $[-2, 2]$ defined by


$$R_i \equiv [-2 + (i-1), -2 + i] \quad (i = 1, 2, 3, 4).$$

Then both sides of (??) are injective on R_i . V and C map R_i 's as follows:

$$V(R_i) = C(R_i) = D_1, \quad V(R_j) = C(R_j) = D_2 \quad (i = 1, 4, j = 2, 3)$$

where we note $D_1 = R_3 \cup R_4$, $D_2 = R_1 \cup R_2$.





Lemma 3.2 *The following relations hold for v_i, c_i, q_i in (??):*

$$v_2 \circ c_1 = c_2 \circ q_1, \quad v_1 \circ c_2 = c_1 \circ q_2, \quad v_2 \circ c_2 = c_2 \circ q_2, \quad v_1 \circ c_1 = c_1 \circ q_1. \quad (3.5)$$

Proof. By (??), we have

$$(C \circ V)|_{R_i} = (Q \circ C)|_{R_i} \quad (3.6)$$

for each $i = 1, 2, 3, 4$. When $i = 1$, $(C \circ V)|_{R_1} = (C \circ v_2^{-1})|_{R_1} = (C)|_{D_2} \circ (v_2^{-1})|_{R_1} = c_2^{-1} \circ (v_2^{-1})|_{R_1}$. In the same way, $(Q \circ C)|_{R_1} = q_2^{-1} \circ (c_2^{-1})|_{R_1}$. Therefore $c_2^{-1} \circ (v_2^{-1})|_{R_1} = q_2^{-1} \circ (c_2^{-1})|_{R_1}$. By taking inverse of both sides, we get the third identity in (??). In the same way, we get others by checking in cases $i = 2, 3, 4$ for (??). ■

By Lemma ??, we have the following relations:

$$v_1 = \begin{cases} c_1 \circ q_1 \circ c_1^{-1} & (\text{on } D_1), \\ c_1 \circ q_2 \circ c_2^{-1} & (\text{on } D_2), \end{cases} \quad v_2 = \begin{cases} c_2 \circ q_1 \circ c_1^{-1} & (\text{on } D_1), \\ c_2 \circ q_2 \circ c_2^{-1} & (\text{on } D_2). \end{cases} \quad (3.7)$$

3.2 Relations of operators arising from a semiconjugacy

By Lemma ?? and Proposition ??, we have three representations π_v, π_c, π_q of $\mathcal{O}_2$ on $L_2[-2, 2]$ associated with three branching function systems $v = \{v_1, v_2\}$, $c = \{c_1, c_2\}$, $q = \{q_1, q_2\}$. By definition of π_v, π_c, π_q , they are given as couples of operators $\{S(v_1), S(v_2)\}$, $\{S(c_1), S(c_2)\}$, $\{S(q_1), S(q_2)\}$ on $L_2[-2, 2]$, and they are concretely obtained as follows:

$$\begin{cases} (S(v_1)\phi)(x) = 2^{1/2}\chi_{[0,2]}(x)\phi(2x-2), \\ (S(v_2)\phi)(x) = 2^{1/2}\chi_{[-2,0]}(x)\phi(-2x-2), \end{cases}$$

$$\begin{cases} (S(c_1)\phi)(x) = \sqrt{\frac{\pi}{2}\sin\frac{\pi}{2}|x|}\chi_{[0,2]}(x)\phi(-2\cos(\pi/2)x), \\ (S(c_2)\phi)(x) = \sqrt{\frac{\pi}{2}\sin\frac{\pi}{2}|x|}\chi_{[-2,0]}(x)\phi(-2\cos(\pi/2)x), \end{cases} \quad (3.8)$$

$$\begin{cases} (S(q_1)\phi)(x) = 2^{1/2}|x|^{1/2}\chi_{[0,2]}(x)\phi(x^2-2), \\ (S(q_2)\phi)(x) = 2^{1/2}|x|^{1/2}\chi_{[-2,0]}(x)\phi(x^2-2) \end{cases}$$

for $\phi \in L_2[-2, 2]$ and $x \in [-2, 2]$ where we use simple notation like (??) instead of (??).

By Lemma ??, (??) and Lemma ?? (ii), we have the followings:

$$\begin{cases} S(v_1) = S(c_1)S(q_1)S(c_1)^* + S(c_1)S(q_2)S(c_2)^*, \\ S(v_2) = S(c_2)S(q_1)S(c_1)^* + S(c_2)S(q_2)S(c_2)^*. \end{cases} \quad (3.9)$$

Note that (??) is essentially derived from (??). Let

$$A_1 \equiv S(v_1) + S(v_2), \quad A_2 \equiv S(q_1) + S(q_2), \quad B \equiv S(c_1) + S(c_2)$$

and $\Omega_0 \equiv B^*1$ where 1 is the constant function on $[-2, 2]$ with value 1.

Lemma 3.3 (i) $A_1 \mathbf{1} = \sqrt{2} \mathbf{1}$. (ii) $A_2 \Omega_0 = \sqrt{2} \Omega_0$.

Proof. (i) By definition of $S(v_i)$, $S(v_i) \mathbf{1} = \sqrt{2} \chi_{D_i}$ for $i = 1, 2$. Hence the statement holds. (ii) Since $S(c_1)^* \mathbf{1} = S(c_2)^* \mathbf{1}$, $B^* \mathbf{1} = 2S(c_i)^* \mathbf{1}$ for $i = 1, 2$,

$$A_2 \Omega_0 = (S(q_1)B^* + S(q_2)B^*) \mathbf{1} = 2(S(q_1)S(c_1)^* + S(q_2)S(c_2)^*) \mathbf{1}.$$

Note $B^* A_1 = 2(S(q_1)S(c_1)^* + S(q_2)S(c_2)^*)$ by (??) with respect to $S(c_1), S(c_2)$. Hence $A_2 \Omega_0 = B^* A_1 \mathbf{1} = \sqrt{2} B^* \mathbf{1} = \sqrt{2} \Omega_0$. ■

Let Ω be the normalization of Ω_0 and denote $q_I = q_{i_1} \circ \dots \circ q_{i_k}$ for a multiindex $I = (i_1, \dots, i_k) \in \{1, 2\}^k$, $k \geq 1$.

Lemma 3.4 For $I \in \{1, 2\}^k$, $k \geq 1$, $(S(q_I)\Omega)(x) = \chi_{D_I}(x) 2^{k/2} \Omega(x)$ where $D_I \equiv q_I([-2, 2])$.

Proof. Since $q_i \circ c_i^{-1} = c_i^{-1} \circ v_i$, $S(q_i)\Omega = \chi_{D_i} \sqrt{2} \Omega$ for $i = 1, 2$. For any measurable subset $Y \subset [-2, 2]$, $S(q_i)M_{\chi_Y} = M_{\chi_{q_i(Y)}} S(q_i)$ for $i = 1, 2$ by Lemma ?? (v). Hence


$$S(q_I)\Omega = 2^{1/2} S(q_{I'}) M_{\chi_{D_{i_k}}} \Omega = (2^{1/2})^2 S(q_{I''}) M_{\chi_{D_{(i_{k-1}, i_k)}}} \Omega = \dots = 2^{k/2} M_{\chi_{D_I}} \Omega$$

when $I = (i_1, \dots, i_k)$ where $I' = (i_1, \dots, i_{k-1})$ and $I'' = (i_1, \dots, i_{k-2})$. ■

Since $S(c_i)^* \mathbf{1} = \Phi_{c_i}^{1/2}$, $i = 1, 2$, we have

$$\Omega(x) = \frac{1}{\sqrt{\pi}} \frac{1}{(4 - x^2)^{1/4}}. \quad (3.10)$$

Note that Ω is the following function:



$\Omega(x)$ is

positive on $-2 < x < 2$. Furthermore the followings hold:

$$\sum_{I \in \{1, 2\}^k} \chi_{D_I} \cdot \Omega = \Omega, \quad (\chi_{D_I} \cdot \Omega)(x) = \begin{cases} \Omega(x) & (x \in D_I), \\ 0 & (\text{otherwise}), \end{cases} \quad (I \in \{1, 2\}^k, k \geq 1).$$

Theorem 3.5 Let $(L_2[-2, 2], \pi_q)$ be the representation of $\mathcal{O}_2$ by $q = \{q_1, q_2\}$ in Proposition ?? and Ω in (??). Then $(L_2[-2, 2], \pi_q, \Omega)$ is $GP(z)$ for $z \equiv \frac{1}{\sqrt{2}}(1, 1)$ in Definition ??.

Proof. By definition, $\|\Omega\| = 1$. By Lemma ??,

$$\pi_q \left(\frac{1}{\sqrt{2}}(s_1 + s_2) \right) \Omega = \frac{1}{\sqrt{2}}(S(q_1) + S(q_2))\Omega = \Omega.$$

Hence $(L_2[-2, 2], \pi_q)$ satisfies (??) for $z = \frac{1}{\sqrt{2}}(1, 1)$. By Lemma ??, $K \equiv \{\chi_{D_I} \cdot \Omega : I \in \{1, 2\}^k, k \geq 1\} \subset \pi_q(\mathcal{O}_2)\Omega$. Since $\Omega(x) > 0$ for $-2 < x < 2$, $L_2[-2, 2] = \overline{\text{Lin}K} \subset \pi_q(\mathcal{O}_2)\Omega$. Therefore $(L_2[-2, 2], \pi_q)$ is a cyclic representation of $\mathcal{O}_2$ with a unit cyclic vector Ω . Hence $(L_2[-2, 2], \pi_q, \Omega)$ is $GP(z)$ by Definition ?? for $z = \frac{1}{\sqrt{2}}(1, 1)$. ■

Proof of Theorem ??. Note that the definition of π_q in (??) is same in (??). The uniqueness of eigen vector Ω of $\pi_q(s_1) + \pi_q(s_2)$ in Theorem ?? follows from Theorem ?? (iv). Hence (i) is proved. (ii) follows from Theorem ?? (iii) follows from Theorem ?? (ii). ■

We check the eigen equation (??) for Theorem ?? by direct computation according to definitions of operators in (??) and the eigen function in (??). Note

$$(\Phi_{c_i} \circ Q)(x) = \frac{1}{|x|} \Phi_{c_i}(x), \quad (\Phi_{q_i} \circ Q)(x) = \frac{1}{2|x|}, \quad (3.11)$$

$$\Omega(q_i^{-1}(x)) = \Omega(Q(x)) = \frac{1}{\sqrt{\pi}} \frac{1}{(4 - (x^2 - 2)^2)^{1/4}} = \frac{1}{\sqrt{|x|}} \Omega(x) \quad (3.12)$$

for $i = 1, 2$ and $x \in [-2, 2]$. From these relations,

$$(S(q_i)\Omega)(x) = \chi_{D_i}(x) \{\Phi_{q_i}(Q(x))\}^{-1/2} \Omega(Q(x)) = \chi_{D_i}(x) \sqrt{2} \Omega(x).$$

Hence $\frac{1}{\sqrt{2}}(S(q_1) + S(q_2))\Omega = \Omega$ and Ω is the eigen vector of $\frac{1}{\sqrt{2}}(S(q_1) + S(q_2))$ with eigen value 1 certainly.

4 Generalization

We generalize Theorem ?? slightly.

By Proposition ??, if we have a conjugacy between two branching function systems, then we have a unitary equivalence between two representations of $\mathcal{O}_N$. Therefore we construct a new branching function system from the system $([-2, 2], Q)$ in (??) by the following conjugacy $\varphi_{(a,b)}$: Let $a, b \in \mathbf{R}$, $a < b$ and $\varphi_{(a,b)} : [-2, 2] \rightarrow [a, b]$ defined by

$$\varphi_{(a,b)}(x) = \frac{b-a}{4}x + \frac{a+b}{2}, \quad \varphi_{(a,b)}^{-1}(x) = \frac{4}{b-a}x - 2\frac{a+b}{b-a}.$$

Note $\varphi_{(a,b)}$ satisfies the assumption in Lemma ?? . Put $Q^{(a,b)} \equiv \varphi_{(a,b)} \circ Q \circ \varphi_{(a,b)}^{-1}$, $q_{\pm}^{(a,b)} \equiv \left(Q^{(a,b)}|_{D_{\pm}^{(a,b)}} \right)^{-1}$ for $D_{-}^{(a,b)} \equiv [a, \frac{b+a}{2}]$, $D_{+}^{(a,b)} \equiv [\frac{b+a}{2}, b]$. Then

$$Q^{(a,b)}(x) = \frac{4}{b-a} \left(x - \frac{a+b}{2} \right)^2 + a, \quad q_{\pm}^{(a,b)}(x) = \frac{a+b \pm \sqrt{(b-a)(x-a)}}{2}.$$

The 1-dimensional dynamical system $([a, b], Q^{(a,b)})$ is conjugate with $([-2, 2], Q = Q^{(-2,2)})$ by construction. The representation $(L_2[a, b], \pi_{q^{(a,b)}})$ of $\mathcal{O}_2$ associated with the branching function system $q^{(a,b)} = \{q_{\pm}^{(a,b)}\}$ is given by

$$\pi_{q^{(a,b)}}(s_1) \equiv S(q_{+}^{(a,b)}), \quad \pi_{q^{(a,b)}}(s_2) \equiv S(q_{-}^{(a,b)}),$$

$$\left(S(q_{\pm}^{(a,b)}) \phi \right)(x) = 2 \sqrt{\frac{|2x-b-a|}{b-a}} \chi_{D_{\pm}}(x) \phi(Q^{(a,b)}(x)) \quad (4.1)$$

for $\phi \in L_2[a, b]$ and $x \in [a, b]$. Put $\Omega^{(a,b)} \equiv U_{\varphi_{(a,b)}} \Omega$ where $U_{\varphi_{(a,b)}}$ is in Lemma ?? (i). Then

$$\Omega^{(a,b)}(x) = \frac{1}{\sqrt{\pi}} \frac{1}{\{(3b-a-2x)(b-3a+2x)\}^{1/4}}.$$

For example, $Q^{(-1,1)}(x) = 2x^2 - 1$, $Q^{(-b,b)}(x) = \frac{4}{b}(x - \frac{b}{2})^2$, $Q^{(0,b)}(x) = \frac{2}{b}x^2 - b$ for $b > 0$.

By Theorem ?? and Proposition ??, we have the following result immediately.


Corollary 4.1 *The representation $(L_2[a, b], \pi_{q^{(a,b)}})$ of $\mathcal{O}_2$ is $GP(z)$ for $z = \frac{1}{\sqrt{2}}(1, 1)$. Specially, $(L_2[a, b], \pi_{q^{(a,b)}})$ is irreducible. $\Omega^{(a,b)}$ is unique eigen vector of $\frac{1}{\sqrt{2}}(\pi_{q^{(a,b)}}(s_1) + \pi_{q^{(a,b)}}(s_2))$ with eigen value 1.*

Remark that a given quadratic function does not always appears as $Q^{(a,b)}$. For example a transformation $P(x) = x^2$ on $[0, 1]$ is impossible to describe as $Q^{(a,b)}$.

Example 4.2 Denote $Q'(x) \equiv Q^{(0,1)}$, $q'_{\pm} \equiv q_{\pm}^{(0,1)}$, $V' \equiv \varphi_{(0,1)} \circ V \circ \varphi_{(0,1)}^{-1}$ and $v'_{\pm} \equiv \varphi_{(0,1)} \circ v_{\pm} \circ \varphi_{(0,1)}^{-1}$ where $v_{+} \equiv v_1$ and $v_{-} \equiv v_2$ in (??). Then

$$Q'(x) = (2x-1)^2, \quad q'_{\pm}(x) = \frac{1 \pm \sqrt{x}}{2}, \quad V'(x) = |2x-1|, \quad v'_{\pm}(x) = \frac{1 \pm x}{2}$$

for $x \in [0, 1]$.



For a branching function system $q' = \{q'_\pm\}$ on $[0, 1]$, we have a representation $\{S(q'_\pm)\}$ of $\mathcal{O}_2$ on $L_2[0, 1]$ as follows:

$$(S(q'_\pm)\phi)(x) = \chi_{D'_\pm}(x) \sqrt{2|2x-1|} \phi((2x-1)^2)$$

for $\phi \in L_2[0, 1]$ and $x \in [0, 1]$ where $D'_+ \equiv [\frac{1}{2}, 1]$ and $D'_- \equiv [0, \frac{1}{2}]$. The eigen vector is given by

$$\Omega'(x) \equiv \Omega^{(0,1)}(x) = \sqrt{\frac{2}{\pi}} \frac{1}{(x-x^2)^{1/4}}.$$

We can check directly that Ω' is the eigen vector of $\frac{1}{\sqrt{2}}(\pi_{q'}(s_1) + \pi_{q'}(s_2))$ with eigen vector 1.

5 Discussion

C*-algebras associated with dynamical systems are often studied([?]). However there are few studies about representations but not algebras themselves. It is not clear that relations between the properties of dynamical systems and those of representations of $\mathcal{O}_N$ by comparison with those between algebras themselves and dynamical systems. Our interest is the construction of representation of $\mathcal{O}_N$ and its characterization. Hence we list up forward problems in this point of view.

(1) In the theory of dynamical system, a transformation

$$Q_c(x) \equiv x^2 + c$$

is more general. We give a representation of $\mathcal{O}_2$ arising from a quadratic map Q_{-2} and characterize it. Can we construct similar representation from Q_c when $c \neq -2$? If it is possible, then how can we characterize it ?

For example, $f_1(x) = \frac{1}{2}x^2$ and $f_2(x) = \frac{1}{2}x^2 + \frac{1}{2}$ gives a branching function system $f = \{f_1, f_2\}$ on $[0, 1]$. However we have no idea to characterize the representation $(L_2[0, 1], \pi_f)$ of $\mathcal{O}_2$ yet.

Can we treat the dynamical system of polynomial (transformation) $P(x) = a_0 + a_1x + \dots + a_nx^n$?

(2) Equations (??) is derived from (??). That is, equations of transformations(or real functions) change operator relations by the map S . We are interest in other equation which is similar in this case.

(3) Equations (??) of six operators means a relation of three representations π_v, π_c, π_q of $\mathcal{O}_2$ on $L_2[-2, 2]$. This relation seems unique. How should they be studied ? We expect that similar relations will be derived from other semiconjugacies of dynamical systems.

(4) We can show easily that $(L_2[-2, 2], \pi_v, \frac{1}{2})$ in (??) is $GP(z)$ for $z = \frac{1}{\sqrt{2}}(1, 1)$ where $\frac{1}{2}$ is the constant function on $[-2, 2]$ with value $\frac{1}{2}$. But $(L_2[-2, 2], \pi_c)$ is not known yet. Is $(L_2[-2, 2], \pi_c)$ unitarily equivalent to $(L_2[-2, 2], \pi_v)$, too ?

References

- [1] O. Bratteli and P. E. T. Jørgensen, *Iterated Function Systems and Permutation Representations of the Cuntz Algebra*, Memoirs Amer. Math. Soc. **39** No. 663 (1999).

- [2] J. Cuntz, *Simple C^* -algebras generated by isometries*, Comm. Math. Phys. **57** (1977), 173-185.
- [3] Robert L. Devaney, *A first course in chaotic dynamical systems: Theory and experiment*, Addison-Wesley Publishing Company, Inc. (1992).
- [4] K.Kawamura, *Generalized permutative representation of Cuntz algebra. I —Generalization of cycle type—*, preprint, RIMS-1380 (2002).
- [5] K.Kawamura, *Generalized permutative representation of Cuntz algebra. II —Irreducible decomposition of periodic cycle—*, preprint, RIMS-1388 (2002).
- [6] K.Kawamura, *Generalized permutative representation of Cuntz algebra. III —Generalization of chain type—*, in preparation.
- [7] K.Kawamura, *Generalized permutative representation of Cuntz algebra. IV —Gauge transformation of representation—*, in preparation.
- [8] K.Kawamura, *Construction of representations of the Cuntz-Krieger algebra*, in preparation.
- [9] A.Connes, *Noncommutative Geometry*, Academic Press, Inc. (1994).