

Harmonic Univalent Functions for Which Analytic Part is Starlike

Yaşar POLATOĞLU

Abstract

In a simply connected domain $\mathcal{U} \subset \mathbb{C}$ a complex-valued harmonic function f has the representation $f = h + \bar{g}$, where h and g are analytic function in $\mathcal{U}$, and are called analytic part and co-analytic part of f , respectively.

Let $h(z) = a_0 + a_1z + a_2z^2 + \cdots$ and $g(z) = b_0 + b_1z + b_2z^2 + \cdots$ be analytic functions in the open unit disc $\mathbb{D} = \{z \in \mathbb{C} | |z| < 1\}$, if $(|h'(z)|^2 - |g'(z)|^2) > 0$ or $(|h'(z)|^2 - |g'(z)|^2) < 0$ then f is called a sense-preserving harmonic function in $\mathbb{D}$. The class of all sense-preserving harmonic functions in $\mathbb{D}$ with $a_0 = b_0 = 0$, and $a_1 = 1$ will be denote by $\mathcal{S}_H$. Thus $\mathcal{S}_H$ contains the standard class of analytic univalent functions $\mathcal{S}$.

The aim of this paper is to investigate the subclass of $\mathcal{S}_H$. By choosing only these functions whose analytic parts are starlike functions for such mappings we will find distortion theorems.

1 Introduction

Let $h(z) = z + a_2z^2 + \cdots$ be an analytic function in the open unit disc $\mathbb{D}$, if $h(z)$ satisfies the condition

$$\operatorname{Re} \left(z \frac{h'(z)}{h(z)} \right) > 0, \quad (1.1)$$

then $h(z)$ is called starlike function in $\mathbb{D}$, and the class of starlike functions in $\mathbb{D}$ is denoted by $\mathcal{S}^*$ [3].

2000 *Mathematics Subject Classification*: Primary 30C45.

Key words and phrases: harmonic univalent functions, starlike functions, sense-preserving.

Next Ω be the family of functions $\phi(z)$ which are regular and satisfying the conditions $\phi(0) = 0$, $|\phi(z)| < 1$ for every $z \in \mathbb{D}$, and let $\Omega(a)$ be the class of functions $w(z)$ which are analytic in $\mathbb{D}$ and satisfying the conditions $w(0) = a$, $|w(z)| < 1$ for all $z \in \mathbb{D}$. We note that $\Omega_{\mathcal{H}}(a)$ be the union of all classes $\Omega(a)$ whereas a ranges over $[0, 1)$ [4].

Moreover, a function f is said to be a complex-valued harmonic function in $\mathbb{D}$ if both $\operatorname{Re} f$ and $\operatorname{Im} f$ are real harmonic in $\mathbb{D}$. Every such f can be uniquely represented by

$$f = h + \bar{g}, \quad (1.2)$$

where h and g are analytic in $\mathbb{D}$ with $g(0) = 0$. A complex valued harmonic function f not identically constant, satisfying (1.2) is said to be sense-preserving in $\mathbb{D}$ if and only if satisfying the equation

$$g'(z) = w(z)h'(z), \quad (1.3)$$

where $w(z)$ is analytic in $\mathbb{D}$ with $|w(z)| < 1$ for every $z \in \mathbb{D}$ [2], and the function $w(z)$ is called the second dilatation of f . It is closely related to the Jacobian of f is defined by

$$J_f(z) = |h'(z)|^2 - |g'(z)|^2. \quad (1.4)$$

Finally, let $h(z) = \sum_{n=0}^{\infty} a_n z^n$ and $g(z) = \sum_{n=2}^{\infty} b_n z^n$ be analytic functions in $\mathbb{D}$. Choose $g(0) = 0$, i.e., $b_0 = 0$, so the representation (1.2) is unique in $\mathbb{D}$ and is called the canonical representation of f . For univalent and sense-preserving harmonic functions f in $\mathbb{D}$, it is convenient to make further normalization (with no loss generality) $h(0) = 0$, i.e., $a_0 = 0$ and $h'(0) = 1$, i.e., $a_1 = 1$. The family of all such functions f is denoted by $\mathcal{S}_{\mathcal{H}}$ [1], [2], [5].

In this paper we will study on the subclass of $\mathcal{S}_{\mathcal{H}}$ consisting of all univalent harmonic functions for which analytic part is starlike, and this class will be denoted by $\mathcal{S}_{\mathcal{H}}^*$.

2 Main Results

Lemma 2.1. *Let $w(z)$ be element of $\Omega(a)$, then*

$$\frac{|a - r|}{1 - ar} \leq |w(z)| \leq \frac{a + r}{1 + ar} \quad (2.1)$$

for all $z \in \mathbb{D}$, and $0 \leq a < 1$.

Proof. Let $w(z) \in \Omega(a)$, then $w(z)$ is analytic in $\mathbb{D}$ and satisfies the condition $w(0) = a$. Now we consider the function

$$\phi(z) = \frac{w(z) - w(0)}{1 - \overline{w(0)}w(z)}, \quad z \in \mathbb{D}. \quad (2.2)$$

Therefore $\phi(z)$ satisfies the conditions of Schwarz lemma. Using the estimate the Schwarz lemma $|\phi(z)| \leq |z| = r$, which gives

$$|\phi(z)| = \left| \frac{w(z) - w(0)}{1 - \overline{w(0)}w(z)} \right| \leq r. \quad (2.3)$$

The inequality (2.3) can be written in the following form

$$\left| \frac{w(z) - a}{1 - \overline{a}w(z)} \right| \leq r \Leftrightarrow |w(z) - a| \leq r|aw(z) - 1|. \quad (2.4)$$

The inequality (2.4) is equivalent

$$\left| w(z) - \frac{a(1 - r^2)}{1 - a^2 r^2} \right| \leq \frac{r(1 - a^2)}{1 - a^2 r^2}. \quad (2.5)$$

The equality holds in the inequality (2.5) only for the function

$$w(z) = e^{i\beta} \frac{e^{i\theta} z + a}{1 + \overline{a}e^{i\theta} z}, \quad z \in \mathbb{D}.$$

From the inequality (2.5) we have

$$\begin{aligned} |w(z)| &\geq \left| \frac{a(1 - r^2)}{1 - a^2 r^2} - \frac{r(1 - a^2)}{1 - a^2 r^2} \right| = \frac{|a - r|}{1 - ar}, \\ |w(z)| &\leq \frac{a(1 - r^2)}{1 - a^2 r^2} + \frac{r(1 - a^2)}{1 - a^2 r^2} = \frac{a + r}{1 - ar}. \end{aligned} \quad (2.6)$$

□

Corollary 2.2. Let $f \in \mathcal{S}_{\mathcal{H}}^*$, then

$$\frac{(1 - r)|a - r|}{(1 + r)^3(1 - ar)} \leq |g'(z)| \leq \frac{(1 + r)(a + r)}{(1 - r)^3(1 + ar)}. \quad (2.7)$$

Proof. Let $f \in \mathcal{S}_{\mathcal{H}}^*$, then

$$g'(z) = w(z)h'(z) \Leftrightarrow \frac{g'(z)}{h'(z)} = w(z). \quad (2.8)$$

Using lemma 2.1 we obtain

$$|h'(z)| \frac{|a-r|}{(1-ar)} \leq |g'(z)| \leq |h'(z)| \frac{a+r}{(1+ar)}. \quad (2.9)$$

On the other hand, since $h(z)$ is starlike, then $h(z)$ satisfies the following inequality

$$\frac{1-r}{(1+r)^3} \leq |h'(z)| \leq \frac{1+r}{(1-r)^3}. \quad (2.10)$$

Considering the inequalities (2.9) and (2.10) together we obtain (2.7). $\square$

Theorem 2.3. Let $f \in \mathcal{S}_{\mathcal{H}}^*$, then

$$|g(z)| \leq \log \left(\frac{1+ar}{1-r} \right)^{\left(\frac{a-1}{a+1}\right)^2} + \frac{a-3}{a+1} \cdot \frac{r}{1-r} + \frac{2r^2}{(1-r)^2}. \quad (2.11)$$

Proof. Applying the estimate (2.7) we have

$$|g(z)| = \left| \int_C g'(\zeta) d\zeta \right| \leq \int_C |g'(\zeta)| |d\zeta| \leq \int_0^r \frac{(1+\rho)(a+\rho)}{(1-\rho)^3(1+a\rho)} d\rho.$$

Integrating, we obtain the estimate (2.11), where $C = [0, z]$ is Jordan arc. $\square$

Theorem 2.4. Let f be element of $\mathcal{S}_{\mathcal{H}}^*$, then

$$\begin{aligned} & \log \left(\frac{1+r}{1+ar} \right)^{\left(\frac{1+a}{1-a}\right)^2} - \frac{2r[(1+a)r+2a]}{(1-a)(1+r)^2} \leq |f(z)| \\ & \leq \frac{r(1+2r)}{(1-r)^2} + \frac{a-3}{a+1} \cdot \frac{r}{1-r} + \log \left(\frac{1+ar}{1-r} \right)^{\left(\frac{a-1}{a+1}\right)^2}. \end{aligned}$$

Proof. Let $z \in \mathbb{D}$. We denote $|z| = r$ and $M(r) = \inf\{|f(z)| : |z| = r\}$. Then $|f(z)| \geq M(r)$ and $\{w : |w| \leq M(r)\} \subset f(\{\zeta : |\zeta| < r\}) \subset f(\mathbb{D})$, hence there exist z_r satisfying $|z_r| = r$ such that $M(r) = |f(z_r)|$ and $\gamma(t) = tf(z_r)$,

$t \in [0, 1]$. Therefore $f^{-1}(\gamma(t)) = \Gamma(t)$, $t \in [0, 1]$ is well defined Jordan arc, and

$$\begin{aligned} |f(z)| &\geq \int_{f^{-1}(\gamma(t))} (|h'(\zeta)| - |g'(z)|) |d\zeta| = \int_{f^{-1}(\gamma(t))} (|h'(\zeta)| - |w(\zeta)||h'(z)|) |d\zeta| \\ &= \int_{f^{-1}(\gamma(t))} |h'(\zeta)|(1 - |w'(z)|) |d\zeta| \geq \int_0^r \left(\frac{1-\rho}{(1+\rho)^3} \right) \left(1 - \frac{a+\rho}{1+a\rho} \right) d\zeta \Rightarrow \end{aligned}$$

$$|f(\zeta)| \geq \log \left(\frac{1+r}{1+ar} \right)^{\left(\frac{1+a}{1-a} \right)^2} - \frac{2r[(1+a)r + 2a]}{(1-a)(1+r)^2}.$$

On the other hand, we have

$$|f(z)| = |h(z) + \overline{g(z)}| \leq |h(z)| + |g(z)|,$$

then using the following inequality and theorem 2.3, we have

$$|f(z)| \leq \frac{r(1+2r)}{(1-r)^2} + \frac{a-3}{a+1} \cdot \frac{r}{1-r} + \log \left(\frac{1+ar}{1-r} \right)^{\left(\frac{a-1}{a+1} \right)^2}.$$

□

Theorem 2.5. Let $f \in \mathcal{S}_{\mathcal{H}}^*$, then

$$\frac{(1-a^2)(1-r)}{(1+ar)^2(1+r)^5} \leq J_f(z) \leq \frac{(1-ar)^2 - |a-r|^2}{(1-ar)^2(1-r)^6}.$$

Proof. Using lemma 2.1 and the following relations,

$$J_f(z) = |h'(z)|^2 - |g'(z)|^2,$$

$$g'(z) = w(z)h'(z),$$

and after the simple calculations we get

$$\frac{(1-a^2)(1-r)}{(1+ar)^2(1+r)^5} \leq |h'(z)|^2(1 - |w(z)|^2) \leq \frac{(1-ar)^2 - |a-r|^2}{(1-ar)^2(1-r)^6}.$$

□

References

- [1] J. Clunie, T. Sheil-Small, Harmonic univalent functions, *Ann. Acad. Sci. Fenn. Ser. A I Math.*, **9** (1984), 3-25.
- [2] P. Duren, *Harmonic Mappings in the Plane*, Cambridge University Press, UK, 2004.
- [3] A.W. Goodman, *Univalent Functions*, Vol I. and II, Mariner Publ. Comp., Tampa, Florida, 1984.
- [4] G.M. Golusin, Geometrische Funktionentheorie. (German) Hochschulbücher für Mathematik, Bd. 31. VEB Deutscher Verlag der Wissenschaften, Berlin, 1957.
- [5] H. Lewy, On the non-vanishing of the Jacobian in certain one-to-one mappings, *Bull. Amer. Math. Soc.*, **42** (1936), 689-692.

YAŞAR POLATOĞLU

Department of Mathematics and Computer Science,
İstanbul Kültür University, 34156 İstanbul, Turkey
e-mail: y.polatoglu@iku.edu.tr