

1

2

31Function of meiobenthos and microorganisms in cellulose breakdown in sediments

4

5

62of wetlands with different origins in Hokkaido

7

8

93

10

11

124

13

14

155Kyohei Yamada • Haruhiko Toyohara

16

17

186Division of Applied Biosciences, Graduate School of Agriculture, Kyoto University,

19

20

21

227Kyoto 606-8502, Japan

23

24

258

26

27

289Corresponding author

29

30

3110Haruhiko Toyohara

32

33

34

3511Tel/Fax: 81-075-753-6446

36

37

3812Email: toyohara@kais.kyoto-u.ac.jp

39

40

4113

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

1
2
3 14 Abstract
4
5

6 15 To validate the mechanism of cellulose breakdown in cold climate wetlands, we
7
8
9
10 16 investigated cellulase activity in sediments collected from 17 wetland sites in Hokkaido,
11
12
13 17 the northern area of Japan. We evaluated cellulase activity by quantitative analysis of
14
15
16 18 glucose released from carboxymethyl cellulose and found that sediments from peat fens
17
18
19 19 demonstrated high activity, followed by sediments from lagoons and estuaries.
20
21
22 20 Sediments from peat fens also contained greater amounts of organic matter, followed by
23
24
25 21 lagoons and estuaries, thereby suggesting a strong positive correlation between organic
26
27
28 22 matter content and cellulase activity. Evaluation of cellulase activity by qualitative
29
30
31 23 cellulose zymographic analysis showed that various cellulases with different molecular
32
33
34 24 sizes were implicated in cellulose breakdown in wetlands. Among them, cellulose
35
36
37
38 25 breakdown in Meguma Pond (peat fen), Notsuke Gulf (peat fen) and Lake Utonai
39
40
41 26 (lagoon) was potentially due to microorganism cellulase, while that in Lake Chobushi
42
43
44 27 (lagoon) was ascribed to meiobenthos (Oligochaeta species) cellulase. The findings
45
46
47 28 presented herein suggest that the origin and activity level of cellulase varied, depending
48
49
50 29 on the types of cold climate wetlands.
51
52
53

54 30 **Keywords:** Cellulase • Cellulose • Cold district • Microorganism • Hokkaido •
55
56

57 31 Meiobenthos • Sediment • Wetland
58
59
60
61 32

Introduction

Wetlands play ecologically important roles as breeding grounds and stopping points for migratory birds, as well as habitats for aquatic invertebrates, because of the richness of nutrients derived from rivers, lakes, and seas [1]. Cellulose, a component of plant cell walls, is a major organic material in the sediment of wetlands. Cellulose is a high-molecular-weight polysaccharide comprised of β -1,4-linked glucose residues and biochemically stable compared to starch, in which glucose residues are bound by α -1,4 linkages and α -1,6 linkages [2,3]. Cellulase, which is a general term for enzymes that belong to the glycoside hydrolase family and catalyzes the hydrolysis of the β -1,4-glycoside linkages of cellulose chains, includes endo- β -1,4-glucanase (EC 3.2.1.4) and cellobiohydrolase (EC 3.2.1.91). Endo- β -1,4-glucanase and cellobiohydrolase degrade cellulose to cellulodextrin or cellobiose, and another enzyme β -glucosidase (EC 3.2.1.21) further degrades them into glucose [4]. Cellulases from bacteria [5], filamentous fungi [6], basidiomycetes [7], myxomycetes [8], and protozoa [9] have been extensively studied. Occurrence of cellulase of which genes are encoded on chromosomes of their own have been reported from termite [10] and nematoda [11, 12]. Occurrence of these endogenous cellulases has also been reported in aquatic

1
2
3 51 animals, such as blue mussels, abalones, sea urchins [13, 14, 15], and brackish clam
4
5
6 52 [16].
7
8

9
10 53 Cellulase and β -acetylglucosaminidase activities in sediments collected from
11
12 54 various wetlands in Japan were measured as part of the research conducted for The
13
14
15
16 55 International Collaborative Research on the Management of Wetland Ecosystem of the
17
18
19 56 National Institute for Environmental Studies between 1998 and 2002 [17]. In this report,
20
21
22 57 high cellulase activities were detected in the sediments from Lake Furen and Biwase
23
24
25
26 58 River, located in the east area of Hokkaido Prefecture of Japan, and the activities were
27
28
29 59 assumed to be derived from microorganisms. Recently, it was shown that the cellulase
30
31
32 60 activities in these northern areas of Japan can be ascribed to meiobenthos, but not to
33
34
35 61 microorganisms, and suggested that meiobenthos play an important role in the
36
37
38 62 breakdown of cellulose, especially in cold climates [18]. Meiobenthos are defined as
39
40
41 63 animal that pass through a 1-mm mesh filter and are known to be composed of a variety
42
43
44 64 of fauna corresponding to 22 phyla [19].
45
46

47 65 There are many untouched wetlands in Hokkaido, which has the greatest
48
49
50
51 66 number of wetlands on the registry of the 500 most important wetlands in Japan
52
53
54 67 maintained by the Ministry of Environment [20] and Ramsar Convention [21]. Wetlands
55
56
57 68 are classified as lakes, rivers, or estuaries. Hokkaido has many lakes, most of which are
58
59
60
61
62
63
64
65

1
2
3 69 classified as lagoons that were formed when a part of the sea was enclosed by land.
4
5
6 70 Many lagoons are located in Hokkaido (e.g., Lake Saroma and Lake Furen).
7
8
9 71 Land-derived organic matter accumulates more easily in lagoons than in estuaries,
10
11
12 72 because lagoons have only a narrow mouth open to the sea [22]. Many peat fens are
13
14
15 73 localized in the eastern and northern parts of Hokkaido, because cellulose breakdown by
16
17
18 74 microorganisms is suppressed at low level due to low temperature throughout a year.
19
20
21 75 For example, annual mean temperatures around Meguma Pond and Notsuke Gulf in
22
23
24 76 2010 were 6.7°C and 6.3°C, respectively (Japan Meteorological Agency Web:
25
26
27
28 77 <http://www.jma.go.jp/> “Accessed 19 August 2011”). Because enough amount of
29
30
31 78 cellulose derived from undecayed plants in peat fens could be available, it is assumed
32
33
34 79 that various cellulose consumers inhabit there [23]. Although various types of wetlands
35
36
37 80 located in Hokkaido are presumed to be inhabited by diverse cellulose consumers such
38
39
40 81 as microorganisms and meiobenthos, it remains unknown what types of organisms are
41
42
43 82 mainly involved in cellulose breakdown in these wetlands.
44
45
46

47
48 83 In the present study, in order to evaluate cellulose breakdown in cold climate
49
50
51 84 wetlands, we compared the degree of cellulose breakdown among the different types of
52
53
54 85 wetlands in Hokkaido and tried to identify major cellulose consumers in these wetlands.
55
56

57 86
58
59
60
61
62
63
64
65

Materials and methods

Materials

Figure 1 shows the sampling sites and their latitude and longitude measured by

Fig. 1

a handy GPS (eTrex Vista HCx; Garmin, Olathe, KS, USA). Sampling was performed

from early to mid-August 2010 and from mid-September to early October 2010. We

collected sediments from 11 lagoons (Koetoi Onuma Pond, Lake Kuccharo, Lake

Saroma, Lake Noto, Lake Abashiri, Lake Furen, Mochirippu Pond, Lake Akkeshi,

Pashikuru Pond, Lake Chobushi, and Lake Utonai), 2 peat fens (Notsuke Gulf and

Meguma Pond), and 4 estuaries (Teshio River, Ishikari River, Mukawa River, and Saru

River). Sediments from Lake Saroma, Lake Noto, Lake Abashiri, Lake Akkeshi, Lake

Furen, Notsuke Gulf, Mochirippu Pond, Pashikuru Pond, and Lake Chobushi were

collected on August 9–12, 2010, and those from the other sites were collected from

September 29 to October 2, 2010. We collected approximately 1 kg of sediments from a

depth of 5 cm of each collecting site. We selected one collecting site apparently without

plants for each wetland and transported these samples at 4°C back to the laboratory at

Kyoto University. Sediment samples were stored at 4°C until analyses. Salt

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

105 concentration of environmental water from each sampling site was measured by a
106 salinometer (IS/Mill-E; AS ONE corporation, Osaka, Japan). Table 1 and Table 2 show
107 salinity and composition of grain sizes of each wetland, respectively. Unless otherwise
108 specified, special grades of reagents were commercially obtained from nacalai tesque
109 (Kyoto, Japan).
110
111 Measurement of sediment cellulase activity by quantitative analysis
112
113 Cellulase activity of sediments was measured within 2 weeks of collection,
114 according to the method of Hayano et al. [24], by using tetrazolium as a coloring agent
115 [25]. Five grams (wet weight) of sediment, passed through a 2 mm-filter, was collected
116 in a 50 ml-conical tube and added to 0.5 ml toluene for sterilization. Next, 10 ml of 0.2
117 M acetate buffer (pH 5.9) and 10 ml of 1% sodium carboxymethyl cellulose (CMC;
118 Sigma, St Louis, MO, US) were added and incubated in a water bath at 30°C for 24 h
119 with shaking. The same reaction mixture containing water instead of CMC was used as
120 a control. After incubation, tubes were centrifuged at $8,000 \times g$ for 5 min, and 100 μ l of
121 supernatant was added to a 1.5-ml tube. One milliliter of blue tetrazolium was added to
122 the tube and heated at 100°C for 4 min in a block incubator (Block Incubator BI-525;

Table 1
Table 2

ASTEC, Fukuoka, Japan), and the absorbance at 660 nm was measured by a spectrophotometer (UV mini 1240; Shimadzu Corporation, Kyoto, Japan) after cooling. The value of the absorbance was converted to glucose concentration by using a standard curve of glucose (0–180 µg/ml) created at the same time. The pellet obtained by centrifugation was dried in a dryer (PS-420; ADVANTEC, Tokyo, Japan) at 60°C overnight to determine the dry weight. Cellulase activity was represented as the amount of glucose released from CMC per 1 g sediment (dry weight) per 1 h.

Isolation of meiobenthos

Meiobenthos were isolated alive from sediments within 1 week of collection. Meiobenthos were recovered in the fraction that included materials small enough to pass through a 1-mm mesh filter but too large to pass through a 63-µm mesh filter. Each meiobenthos was isolated under observation with a microscope (S2X12; Olympus, Tokyo, Japan). Classification of meiobenthos was performed at the level of Class according to Robert et al. [19] except for nematoda due to the difficulty in classification of this species. Classification of arthropods was performed according to Joei et al. [26]. We used single body of meiobenthos for qualitative cellulase assay and two bodies for

quantitative assay.

Cellulase activity of oligochaeta from Notsuke Gulf was measured quantitatively according to the modified method of Niiyama and Toyohara [27]. Briefly, two bodies of living oligochaeta were homogenized with cold 110 μ l phosphate-buffered saline (PBS, containing 140 mM NaCl, 2.7 mM KCl, 8 mM Na_2HPO_4 , and 1.5 mM KH_2PO_4 , pH 7.4). Then, 3 μ l of meiobenthos extract, 3 μ l of 1 M sodium acetate buffer (pH 5.9), and 24 μ l of 1% CMC solution were mixed. Reactions were carried out at 30°C and 4°C for 1, 3, 7, 12, and 24 h with shaking. After incubation, the mixtures were heated at 100°C for 3 min in the block incubator described above to terminate the enzyme reaction. The amount of reducing sugar produced was measured by the tetrazolium blue method [25]. The absorbance at 660 nm was measured with a UV-mini 1240 spectrophotometer.

Preparation and culture of cellulose breakdown microorganisms

Sediment was spread on an agar plate (1.5% agar containing 0.5% CMC, 0.15% $\text{Ca}(\text{NO}_3)_2$, 0.05% MgSO_4 , 0.05% K_2HPO_4) and cultured at 25°C for 1 week. Autoclaved 0.1% soft agar was then added to the cultured plate, and the surface of the

plate containing microorganisms was scraped with a bacteria spreader. Soft agar
 containing cultured microorganisms was added to a liquid culture medium (0.5% CMC,
 0.15% $\text{Ca}(\text{NO}_3)_2$, 0.05% MgSO_4 , and 0.05% K_2HPO_4) and cultured at 25°C for 1 week.
 Culture medium was then filtered through paper filter (No. 1; Toyo Roshi Kaisha,
 Tokyo, Japan), and the filtrate was used for SDS-PAGE zymographic analysis.
 Measurement of cellulase activity by qualitative analysis with sodium dodecyl sulfate
 polyacrylamide gel electrophoresis (SDS-PAGE) zymography
 An aliquot of sediment and a 1/5 volume of 6 × SDS sample buffer (containing
 0.6 M Tris-HCl (pH 6.8), 60% glycerol, 6% SDS, and 0.06% bromophenol blue) were
 mixed with a homogenizer (HandySonic UR-20P; TOMY SEIKO, Tokyo, Japan),
 incubated on ice for 2 h, and centrifuged at 8,000 × *g* for 5 min. The supernatant was
 used for SDS-PAGE zymographic analysis.
 Meiobenthos were picked up from the sediments one by one using a pair of
 tweezers under a binocular microscope (S2X12; Olympus, Tokyo, Japan), and each was
 then homogenized alive with cold 20 µl PBS to prepare a meiobenthos extract for
 SDS-PAGE zymographic analysis. Approximate lengths of each meiobenthos are as

follows. A nematoda obtained from Meguma Pond is 2-3 mm long and that from Lake Utonai is 4 mm long. An oligochaeta species from Meguma Pond, Lake Notoro and Lake Utonai is 1-2 mm long, 4 mm long, and 8 mm long, respectively. A polychaeta species from Lake Utonai is 1-2 mm long. Maxillopoda species from Meguma Pond is 1 mm long.

Cellulase zymographic analysis was performed using 7.5% SDS-PAGE gels containing 0.1% CMC. After electrophoresis, the gels were soaked in 10 mM acetate buffer (pH 5.5) containing 0.1% TritonX-100 for 30 min to remove SDS from the gels. The gels were transferred to 10 mM acetate buffer (pH 5.5), incubated at 37°C or 4°C overnight, and then stained with 0.1% Congo Red. In case of sediment of Notsuke Gulf, the gel was incubated for 4 days because of low activity. The gels were destained using 1 M NaCl. The active bands were detected as nonstained bands.

Measurement of organic component ratio

Dried sediment obtained as mentioned above was heated in a mantle heater (KCA-10A; Koito, Tokyo, Japan) at 600°C for 3 h [28]. Organic component ratio (%)

was calculated according to the formula below.

$$\text{Organic component ratio (\%)} = [(dry\ weight - ignition\ weight)/(dry\ weight)] \times 100$$

Results

Comparison of cellulase activity level by quantitative cellulase analysis

Among 17 wetland sites in Hokkaido, Meguma Pond showed the highest cellulase activity (peat fen, 737.88 nmol/gh, Table 1), followed by Notsuke Gulf (peat fen, 92.39 nmol/gh), Lake Utonai (fresh water lagoon, 44.45 nmol/gh), Lake Saroma (lagoon, 28.48 nmol/gh), Lake Akkeshi (lagoon, 21.42 nmol/gh), and Lake Notoro (lagoon, 13.86 nmol/gh), as summarized in Table 1. Sediments from the estuaries of the Teshio River, Ishikari River, Mukawa River, and Saru River showed little or no cellulase activity.

Qualitative analysis of cellulases by SDS-PAGE zymography

Among 17 wetlands in Hokkaido, active cellulase bands were detected in all

Fig. 2

213 samples by SDS-PAGE zymographic analysis, except for sediments from Pashikuru
 214 Pond, Mukawa River, Saru River, and Lake Abashiri (data not shown). For meiobenthos,
 215 active cellulase bands were detected in the Oligochaeta species in Meguma Pond (Fig.
 216 2), Notsuke Gulf (Fig. 2), Lake Notoro and Lake Abashiri (data not shown), Lake
 217 Chobushi (Fig. 2), Lake Utonai (Fig. 2), Ishikari River, and Koetoi Onuma Pond (data
 218 not shown); Malacostraca species in Lake Kuccharo (data not shown); Nematoda
 219 species in Lake Saroma (data not shown); Foraminifera species in Lake Akkeshi (data
 220 not shown); and Polychaeta species in Teshio River (data not shown).

221 As shown in Fig. 2a, sediment from Meguma Pond demonstrated activity as a
 222 broad smear above 38 kDa. For meiobenthos, Oligochaeta species showed an active
 223 band at 48 kDa, but Nematoda species and Maxillopoda species showed no activity.
 224 However, culture medium of microorganisms showed an active band of high molecular
 225 weight (above 199 kDa).

226 Figure 2b shows the cellulase activity from the Notsuke Gulf sample. Sediment
 227 exhibited intensive active bands at 33 and 87 kDa and faint active bands at 49, 146, 172
 228 and 244 kDa, while Oligochaeta species showed at 26, 29 and 30 kDa. On the other
 229 hand, culture medium of microorganisms showed active bands at 108, 146, 172 and 244
 230 kDa.

Figure 2c shows the cellulase activity from the Lake Notoro sample. Sediment showed weak active bands at 24, 30, and 58 kDa. Oligochaeta species showed a strong active band at 28 kDa and a weak active band at 29 kDa. Ostracoda species demonstrated a weak active band at 27 kDa, while the culture medium of microorganisms showed active bands at 49, 108, and 230 kDa.

Figure 2d shows results from the Lake Chobushi sample. Sediment showed active bands at 33, 59, and 62 kDa, while Oligochaeta species showed active bands at 30, 33, 36, 38, 43, 59, and 62 kDa. Although smear active bands were detected by 24 h-incubation because of the intensive cellulase activity of Oligochaeta species, sharp bands could be detected by 10 h-incubation.

Figure 2e shows the results from the Lake Utonai sample. Sediment showed active bands at 46, 65, and 105 kDa. Nematoda species showed no activity, while Oligochaeta species showed an active band at 68 kDa.

Demonstration of cellulase activity of meiobenthos at low temperature

Fig. 3

As shown in Fig.3, Oligochaeta species demonstrated the substantial cellulase activity bands at 4°C in zymographic analysis, of which activity levels were corresponded to those at 37°C. Oligochaeta species in Notsuke Gulf showed 29 and 30

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

249 kDa active bands, while those in Lake Chobushi showed 36, 38, 43 and 59 kDa active
250 bands.

251 Figure 4 shows the cellulase activity of oligochaeta species in Notsuke Gulf.
252 Higher activity was detected at 30°C than at 4°C. It should be stressed that the activity
253 level at 4°C was almost corresponded with 30% of that at 30°C.

Fig.4

254
255 Relationship between the amount of organic matter and cellulase activity level

256
257 As shown in Table 1, sediment from peat fens such as Meguma Pond and
258 Notsuke Gulf contained large amounts of organic matter, 66.6% and 16.9%, respectively.
259 Sediments from lagoons such as Lake Saroma, Lake Akkeshi, and Lake Utonai
260 contained 1.5%, 6.4%, and 1.5% organic matter, respectively. Sediments from the
261 estuaries of the Teshio River, Ishikari River, and Saru River contained 1.0%, 0.1%, and
262 0.1% organic matter, respectively. There was a strong positive correlation ($r = 0.96$)
263 between the amount of organic matter and the cellulase activity level among sediments
264 collected from 17 wetlands.

266 **Discussion**

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

We measured cellulase activity in sediments collected from 17 wetlands in

Hokkaido to evaluate cellulose breakdown in cold climates. According to our

quantitative analysis (Table 1), sediments from peat fens showed the highest cellulase

activity, followed by those from lagoons and estuaries so far as measured on August

and September in the specific collecting site.

SDS-PAGE zymographic analysis revealed that the molecular size of active

cellulase bands in sediments from Notsuke Gulf (peat fen) corresponded with those

from culture medium of microorganisms. To confirm microorganism cellulases

actually act at cold temperature, we measured activity at 4°C. As shown in Fig. 3(a),

culture medium of microorganisms showed active bands of 146 and 172 kDa,

suggesting that microorganism cellulases might play any function in cellulose

breakdown in Notsuke Gulf. The molecular sizes of active cellulase bands in the

sediments of Lake Chobushi (lagoon) corresponded with those from meiobenthos.

These findings suggest that microorganisms and meiobenthos play important roles in

cellulose breakdown, especially in these wetlands in Hokkaido. However, the

possibility that the molecular sizes of cellulase active bands of sediments and

microorganisms/meiobenthos apparently coincided is not completely ruled out. Further

immunological analysis is needed to validate that the active bands of sediments were derived from microorganisms or meiobenthos.

Oligochaeta showed a strong active band that did not coincide with any bands in the sediment samples from Lake Notoro (Fig. 2c). Despite the fact, it is assumed that Oligochaeta species could play any function in cellulose breakdown in Hokkaido, together with the fact that oligochaeta played an important role in Lake Chobushi as described above. As shown in Fig.3, Oligochaeta species demonstrated the substantial cellulase activity at 4°C in qualitative analysis. Oligochaeta species in Notsuke Gulf actually demonstrated the activity at 4°C almost corresponded with 30% of that at 30°C (Fig.4), suggesting that meiobenthos might play any role to degrade plant residues at low temperature. Since same active bands were demonstrated at 4°C and 37°C, these Oligochaeta species were assumed to possess cellulases active at broad temperature range.

As shown in Table 1, a strong positive correlation was observed between the amount of organic matter and the cellulase activity level. Based on the following facts; (i) organic matters are assumed to be derived from plant residues [29], (ii) in Meguma Pond and Notsuke Gulf where high content of organic matters are detected in sediments, cellulase activity of sediments was derived from microorganisms (Figs. 2a and b), and

(iii) microorganisms secrete cellulases extracellularly[30], (iv) Liu and Toyohara reported that fungal cellulase actually bound to plant residues [31], it seems likely that cellulases secreted from microorganisms would bind to plant residues and degrade them in the wetlands of peat fen sediments. In our preliminary experiments, cellulases from *Corbicula japonica* bound to plant residues similar to fungal cellulases (data not shown), meiobenthos cellulases would function as sediment-binding form in sediment of Hokkaido wetlands.

Acknowledgements

The authors sincerely thank Dr. Chihiro Tanaka, Graduate School of Agriculture, Kyoto University for his help in culturing fungus. This study was partly supported by a grant-in-aid for scientific research from the Ministry of Education, Culture, Sports, Science, and Technology of Japan (no. 22255012).

References

1. Beukema JJ (1979) Biomass and species richness of the macrobenthic animals living on a tidal flat area in the Dutch Wadden sea- effects of a severe winter. Neth

- 321 J Sea Res 13:203-223
- 322 2. Cosgrove DJ (2005) Growth of the plant cell wall. Nat Rev Mol Cell Biol
- 323 6:850-861
- 324 3. Vries RP, Visser J (2001) *Aspergillus* enzymes involved in breakdown of plant cell
- 325 wall polysaccharides. Microbiol Rev 65:497-522
- 326 4. Watanabe H, Tokuda G (2010) Cellulolytic systems in insects. Annu Rev Entomol
- 327 55:609-632
- 328 5. Olson DG, Tripathi SA, Giannone RJ, Lo J, Caiazza NC, Hogsett DA, Hettich RL,
- 329 Guss AM, Dubrovsky G, Lynd LR (2010) Deletion of the Cel48S cellulase from
- 330 *Clostridium thermocellum*. Proc Natl Acad Sci 107:17727-17732
- 331 6. Trinci APJ, Davies DR, Gull K, Lawrence MI, Nielsen BB, Rickers A, Theodorou
- 332 MK (1994) Anaerobic fungi in herbivorous animals. Mycol Res 98:129-152
- 333 7. Chow CM, Yague E, Raguz S, Wood DA, Thurston CF (1994) The *cel3* gene of
- 334 *Agaricus-bisporus* codes for a modular cellulase and is transcriptionally regulated
- 335 by the carbon source. Appl Environ Microbiol 60:2779-2785
- 336 8. Ronsness PA (1968) Cellulolytic enzymes during morphogenesis in *Dictyostelium*
- 337 *discoideum*. J Bacterial 96:639-645
- 338 9. Bera-Maillet C, Devillard E, Cezette M, Jouany JP, Forano E (2005) Xylanases and

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

339 carboxymethylcellulases of the rumen protozoa *Polyplastron multivesiculatum*,
340 *Eudiplodinium maggii* and *Entodinium* sp. FEMS Microbiol Let 244:149-156
341 10. Watanabe H, Noda H, Tokuda G, Lo N (1998) A cellulase gene of termite origin.
342 Nature 394:330-331
343 11. Smant G, Stokkermans PWGJ, Yan Y, Boer de MJ, Baum JT, Wang XH, Hussey RS,
344 Gommers FJ, Henrissat B, Davis EL, Helder J, Schots A, Bakker J (1998)
345 Endogenous cellulases in animals: Isolation of β -1,4-endoglucanase genes from two
346 species of plant-parasitic cyst nematodes. Proc Natl Acad Sci USA 95:4905-4911
347 12. Kikuchi T, Shibuya H, Jones TJ (2005) Molecular and biochemical characterization
348 of an endo- β -1,3-glucanase from the pinewood nematode *Bursaphelenchus*
349 *xylophilus* acquired by horizontal gene transfer from bacteria. Biochem J
350 389:117-125
351 13. Xu BZ, Janson JC, Sellos D (2001) Cloning and sequencing of a molluscan
352 endo-beta-1,4-glucanase gene from the blue mussel, *Mytilus edulis*. Eur J Biochem
353 268:3718-3727
354 14. Suzuki K, Ojima T, Nishita K (2003) Purification and cDNA cloning of a cellulase
355 from abalone *Haliotis discus hannai*. Eur J Biochem 270:771-778
356 15. Nishida Y, Suzuki K, Kumagai Y, Tanaka H, Inoue A, Ojima T (2007) Isolation and

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

357 primary structure of a cellulase from the Japanese sea urchin *Strongylocentrotus*
358 *nudus*. Biochimie 89:1002-1011

359 16. Sakamoto K, Touhata K, Yamashita M, Kasai A, Toyohara H (2007) Cellulose
360 digestion by common Japanese freshwater clam *Corbicula japonica*. Fish Sci
361 73:675-683

362 17. National Institute for Environmental Studies, Japan (2003) International
363 collaborative research on the management of wetland ecosystem (in Japanese).
364 Report of special research from the National Institute for Environmental Studies,
365 Japan, pp 8-13

366 18. Toyohara H, Park Y, Tsuchiya K, Liu W (2011) Cellulase activity in meiobenthos
367 in wetlands. Fish Sci 78:133-137

368 19. Robert PH, Hjalmar T (1988) Introduction to the study of meiofauna. Smithsonian
369 Institution Press, Washington, D.C, pp 243-354

370 20. Nature Convention Bureau in Japanese Ministry of the Environment (2002) Five
371 hundred important wetlands in Japan (in Japanese). Nature Convention Bureau,
372 Tokyo

373 21. Ramsar Convention Secretariat (2011) The list of wetlands of international
374 importance. Ramsar Convention Secretariat, Ramsar, pp 21-22

- 375 22. Wit R, Stal LJ, Lomstein BA, Herbert RA, van Gernerden H, Viaroli P, Cecherelli
376 VU, Rodriguez-Valera F, Bartoli M, Giordani G, Azzoni R, Schaub B, Welsh DT,
377 Donnelly A, Cifuentes A, Anton J, Finster K, Nielsen LB, Pedersen AGU, Neubauer
378 AT, Colangelo MA, Heijs SK (2001) The role of buffering capacities in stabilising
379 coastal lagoon ecosystems. *Continent Shelf Res* 21:2021-2041
- 380 23. Artz RRE, Anderson IC, Chapman SJ, Hagn A, Schlöter M, Potts JM, Campbell CD
381 (2007) Changes in fungal community composition in response to vegetational
382 succession during the natural regeneration of cutover peatlands. *Microbiol Ecol* 54:
383 508-522
- 384 24. Soil Microbiological Society (1992) Experimental methods for soil microbiology
385 (in Japanese). Yokendo, Tokyo, pp 370-371
- 386 25. Chong K J, Peter N L (1985) Determination of reducing sugars in the nanomole
387 range with tetrazolium blue. *J Biochem Biophys Methods* 11:109-115
- 388 26. Joei WM, Gorge ED (2001) An updated classification of the recent crustacean.
389 Natural History Museum of Los Angeles Country, Los Angeles, pp 1-124
- 390 27. Niiyama T, Toyohara H (2011) Widespread distribution of cellulase and
391 hemicellulase activities among aquatic invertebrates. *Fish Sci* 77: 649-655
- 392 28. Hwang I H, Ouchi Y, Matsuto T (2007) Characteristics of leachate from pyrolysis

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

393 residue of sewage sludge. Chemosphere 68:1913-1919

394 29. Nguyen LM (2000) Organic matter composition, microbial biomass and microbial

395 activity in gravel-bed constructed wetlands treating farm dairy wastewaters. Ecol

396 Eng 16:199-221

397 30. Barnet CC, Berka RM, Fowler T (1991) Cloning and amplification of the gene

398 encoding an extracellular bold β -glucosidase from *Trichoderma reesei*: Evidence for

399 improved rates of saccharification of cellulosic substrates. Nat Biotechnol

400 9:562-567

401 31. Liu W, Toyohara H (2012) Sediment-complex-binding cellulose breakdown in

402 wetlands of rivers. Fish Sci. Doi: 10.1007/s12562-012-0471-y

403

Figure captions

Figure 1 Sampling sites of wetlands in Hokkaido. Geological types of wetlands are classified into 3 types: lagoon, peat fen, or estuary. Letters indicating sampling sites correspond to those in Table 1 and Table 2.

Figure 2 Qualitative analysis of cellulase activity by SDS-PAGE cellulose zymography at 37°C. (a) Meguma Pond: Lane 1, sediment; lane 2, Nematoda; lane 3, Oligochaeta; lane 4, Maxillopoda; lane 5, microorganisms. (b) Notsuke Gulf: Lane 1, sediment; lane 2, Oligochaeta; lane 3, microorganisms. (c) Lake Notoro: Lane 1, sediment; lane 2, Oligochaeta; lane 3, *Ostracoda*; lane 4, microorganisms. (d) Lake Chobushi: lane 1, sediment; lane 2, Oligochaeta (24h-incubation); lane 3, Oligochaeta (10 h-incubation). (e) Lake Utonai: Lane 1, sediment; lane 2, Nematoda; lane 3, Oligochaeta; lane 4, microorganisms. Note that active bands of each animal do not reflect the enzyme activity level correctly. *Asterisks* mean that the animal belongs to meiobenthos.

Figure 3 Qualitative analysis of cellulase activity of oligochaeta species from Notsuke Gulf (a) and Lake Chobushi (b). (a) Notsuke Gulf: Lane 1, sediment; lane 2,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

422 Oligochaeta; lane 3, microorganism. *Asterisks* mean that the animal belongs to
423 meiobenthos.
424
425 Figure 4 Cellulase activity of oligochaeta species in Notsuke Gulf at 4°C and 37°C as a
426 function of time. Values are mean ± standard deviation (n=3).

森岡克司先生

前略

このたびはご審査賜りありがとうございました。1名の審査員のコメントに対し下記のように対応いたしました。審査員の指示に従い追加実験を行ったため、訂正原稿の提出が遅れたことをお詫びいたします。ご審査のほど、よろしくお願いいたします。

草々

平成 24 年 2 月 19 日

京都大学農学研究科 豊原治彦

Major points

1. Fig. 2について

1-1 バンドパターンについて

前回から変更されたようですが、今一つ明瞭なバンドが見えているようには感じません。特に Fig. 2b の *sediment* のレーンの 121 と 172kDa のバンドはどう見ても（PC画面上でも印刷しても）はっきりとは見えません。この図ではとても読者を納得させることはできませんので、*sediment* のレーン添加量を増やすか、反応時間を長くすることにより明瞭なバンドを提示してください。Fig. 2eについてもゲル上部にスタックしているという意味では *sediment* と *microorganism* のセルラーゼは同じ性質をもつのかかもしれませんが、これで両者を同じものであると類推するのは無理があると感じます。上記二つのデータは line252-253 に記述されているように、本論文中で非常に重要な論拠となるデータですので、Fig. 2b は再試を奨めます。Fig. 2e は再試で良い結果が出ないようであれば、*sediment* の高分子量のバンドを *microorganism* に帰着させる記述を削除した方が良いと思います。

—ご指摘に従い Fig. 2b の野付湾底泥については4日間の反応を行うことで、明確な活性バンドを検出することができたので、そのデータと差し替えました。訂正した部分は以下の通りです。

L186-187: In case of sediment of Notsuke Gulf, the gel was incubated for 4 days because of low activity. The gels were destained using 1 M NaCl. The active bands were detected as nonstained bands.

L226-230: Sediment exhibited intensive active bands at 33 and 87 kDa and faint active bands at 49, 146, 172 and 244 kDa, while Oligochaeta species showed at 26, 29 and 30 kDa. On the other hand, culture medium of microorganisms showed active bands at 108, 146, 172 and 244 kDa.

—Fig. 2e については再試を行ってもバンドがスタックしてしまったため sediment の高分子バンドを microorganism に帰着させる記述を削除しました。訂正した部分は以下の通りです

L273-275: SDS-PAGE zymographic analysis revealed that the molecular size of active cellulase bands in sediments from Notsuke Gulf (peat fen) corresponded with those from culture medium of microorganisms.

1-2 分類に関して

・ *Nematoda*, *Oligochaeta*, *Harpacticoida*, *Ostracoda*, *Polychaeta* はそれぞれ *meiobenthos* に属する種類であり、*microorganism* に比して解析していることは図中に矢印、もしくは括弧等で示された方が分かりやすいと思います。

—ご指摘に従いそれぞれの生物が *Meiobenthos* であることが分かり易くなるように図中に * で示しました。またそれに伴い Figure caption に記述を加えました。訂正した部分は以下の通りです。

L418: Asterisks mean that the animal belongs to meiobenthos

L422-423: Asterisks mean that the animal belongs to meiobenthos

・ また *Harpacticoida* と *Ostracoda* は斜体になっています。一般に分類表記で斜体は学名の属、種に使うもので、*Harpacticoida* (ソコミジンコ目)、*Ostracoda* (カイクシ目) 等には使わないように感じます。もしなんらかの理由があるなら説明が必要かと思います。

—ご指摘の通り、表記を訂正しました。訂正した部分は以下の通りです。

L233: *Ostracoda* species demonstrated a weak active band at 27 kDa.

L414-415: (c) Lake Notoro: Lane 1, sediment; lane 2, *Oligochaeta*; lane 3, *Ostracoda*; lane 4, microorganisms.

L180:Maxillopoda species from Meguma Pond is 1 mm long.

L223:but Nematoda species and Maxillopoda species showed no activity.

L411-412:Meguma Pond: Lane 1, sediment; lane 2, Nematoda; lane 3, Oligochaeta; lane 4, Maxillopoda; lane 5, microorganisms.

・さらに、これは単なる *suggestion* ですが、上記の *Nematoda* (線形動物門)、*Oligochaeta* (貧毛綱)、*Polychaeta* (多毛綱)、*Harpacticoida* (ソコミジンコ目)、*Ostracoda* (カイクシ下綱) は門、綱、目と分類の階層がまちまちです。*Meiobenthos* の分類は非常に難解なようですが、統一された方が良いでしょうと思います。これに関して *Introduction* 中で *meiobenthod* に関する良い詳細な説明があった方が良いでしょう

—ご指摘の通り、綱(class)レベルに統一しました。節足動物門の分類体系は Joei らの分類に従い(26)、*Harpacticoida* (目)は *Maxillopoda* (綱)へ、*Tanaidacea* は *Malacostraca* (綱)へと表記を訂正しました。*Ostracoda* は Joei らの分類では綱であるため、そのまま表記しました。*Nematoda* に関しましては綱レベルの分類が非常に困難であるため、例外として門レベルで書きその旨を138行目に書き加えました。訂正した部分は以下の通りです。
また62-64行目にメイオベントスの定義、説明及び引用文献を加えました。訂正した部分は以下の通りです。

L62-64:Meiobenthos are defined as animal that pass through a 1-mm mesh filter and are known to be composed of a variety of fauna corresponding to 22 phyla [19].

L138-139:Classification of meiobenthos was performed at the level of Class according to Robert et al. [19] except for nematoda due to the difficulty in classification of this species. Classification of arthropods was performed according to Joei et al. [26].

L180:Maxillopoda species from Meguma Pond is 1 mm long.

L223:but Nematoda species and Maxillopoda species showed no activity.

L411-412:Meguma Pond: Lane 1, sediment; lane 2, Nematoda; lane 3, Oligochaeta; lane 4, Maxillopoda; lane 5, microorganisms.

L218: Malacostraca species in Lake Kuccharo (data not shown);

2. Fig. 3について

決まった温度の影響を定性的に見ていることに非常に疑問を持ちます。

・4℃と30℃での酵素活性を定量化することはできませんか？

—野付湾については新たに 4°C と 30°C で定量的に活性を測定し、その経時的変化を図 4 として追加致しました。この追加に伴い、測定法について新たに下記の文章を追加致しました。長節湖については試料が残っていないため前回と同じく定性的評価のみに留めました。
追加した部分は以下の通りです。

L142-152: Cellulase activity of oligochaeta from Notsuke Gulf was measured quantitatively according to the modified method of Niiyama and Toyohara [27]. Briefly, two bodies of living oligochaeta were homogenized with cold 110 μ l phosphate-buffered saline (PBS, containing 140 mM NaCl, 2.7 mM KCl, 8 mM Na_2HPO_4 , and 1.5 mM KH_2PO_4 , pH 7.4). Then, 3 μ l of meiobenthos extract, 3 μ l of 1 M sodium acetate buffer (pH 5.9), and 24 μ l of 1% CMC solution were mixed. Reactions were carried out at 30°C and 4°C for 1, 3, 7, 12, and 24 h with shaking. After incubation, the mixtures were heated at 100°C for 3 min in the block incubator described above to terminate the enzyme reaction. The amount of reducing sugar produced was measured by the tetrazolium blue method [25]. The absorbance at 660 nm was measured with a UV-mini 1240 spectrophotometer.

L251-252: Figure 4 shows the cellulase activity of oligochaeta species in Notsuke Gulf. Higher activity was detected at 30°C than at 4°C. It should be stressed that the activity level at 4°C was almost corresponded with 30% of that at 30°C.

L291-295: As shown in Fig.3, Oligochaeta species demonstrated the substantial cellulase activity at 4°C in qualitative analysis. Oligochaeta species in Notsuke Gulf actually demonstrated the activity at 4°C almost corresponded with 30% of that at 30°C (Fig.4), suggesting that meiobenthos might play any role to degrade plant residues at low temperature.

L425-426: Figure 4 Cellulase activity of oligochaeta species in Notsuke Gulf at 4°C and 37°C as a function of time. Values are mean $\pm$ standard deviation (n=3).

・また line266 にあるように *Oligochaeta* が 4°C のセルロース分解において重要であるという論調にするならば、Fig. 2b を 4°C で行い、*sediment* と *microorganism* のバンドも同時に考察すべきです。

—ご指摘に従い *sediment*、*Oligochaeta*、*microorganism* について 4°C でザイモグラフィーを行い、その結果を Fig. 3a として元のものとし差し替えました。

Fig 3 aに示すように、Oligochaeta は 4°C で活性バンド (29, 30 kDa) を示したことから、低温度においてもセルロース分解を有しており、底泥中のセルロース分解になんらかの役割を果たしていることが推測されます。しかし、底泥自体のセルロース分解バンド (172, 146 kDa) と Oligochaeta のバンドのサイズは一致しないことから、野付湾底泥において Oligochaeta のセルラーゼは主役ではないと考えられます。したがって旧原稿で 267 行目に記述しました「重要な」という表現は正確ではないので、新たな原稿では下記のように訂正いたしました。

L294-295:, suggesting that meiobenthos might play any role to degrade plant residues at low temperature.

L273-279:SDS-PAGE zymographic analysis revealed that the molecular size of active cellulase bands in sediments from Notsuke Gulf (peat fen) corresponded with those from culture medium of microorganisms. To confirm microorganism cellulases actually act at cold temperature, we measured activity at 4°C. As shown in Fig. 3(a), culture medium of microorganisms showed active bands of 146 and 172 kDa, suggesting that microorganism cellulases might play any function in cellulose breakdown in Notsuke Gulf.

3. *Sediment* とセルラーゼの関係について

・ 審査員 1 も指摘しているように *sediment* 中のセルラーゼと *meiobenthos*、*microorganism* のセルラーゼの関係が今一つ不鮮明です。*meiobenthos*、*microorganism* から分泌されたセルラーゼが *sediment* 中の因子に結合しているという記述 (出来れば引用文献) があった方がよいと思います。

—最近私どもの研究室から菌のセルラーゼが底泥成分、特に植物残渣に強く吸着するというを示す論文を発表いたしました (参考文献 31)。また予備実験ではありますが、ヤマトシジミのセルラーゼが同様に植物残渣等の底泥成分に吸着性を示す結果も得ております。これらの内容を踏まえ新たに、行目に「北海道の泥炭湿地において、微生物由来のセルラーゼが底泥成分に吸着して活性を発現している可能性があること、及びメイオベントス由来のセルラーゼも同様に北海道湿地帯において底泥成分に吸着した形で活性を発現している可能性があること」を示す文章を追加致しました。新たに挿入した部分は以下の通りです。

L303-309:(iv) Liu and Toyohara reported that fungal cellulase actually bound to plant residues [31], it seems likely that cellulases secreted from microorganisms would bind to plant residues and degrade them in the wetlands of peat fen sediments. . In our preliminary experiments, cellulases from *Corbicula japonica* bound to plant residues similar to fungal cellulases (data not shown), meiobenthos cellulases would function as sediment-binding form in sediment of Hokkaido wetlands.

Minor points

Line 25

--*Lake Utonai (lagoon) was potentially due to fungal cellulose*

→ *Lake Utonai (lagoon) was potentially due to microorganism cellulose*

—ご指摘に従い訂正いたしました。訂正した箇所は以下の通りです。

L25-26: Lake Utonai (lagoon) was potentially due to microorganism cellulose

Line 59

Recently, we showed

→ *Recently, it was shown* (文献 18 に本論文著者の名前なし)

—ご指摘に従い訂正いたしました。訂正した箇所は以下の通りです

L59-62: Recently, it was shown that the cellulase activities in these northern areas of Japan can be ascribed to meiobenthos, but not to microorganisms, and suggested that meiobenthos play an important role in the breakdown of cellulose, especially in cold climates [18].

和文要旨

成因が異なる北海道の湿地帯底泥におけるセルロース分解に果たすメイオベントスと微生物の役割

山田京平，豊原治彦（京大院農）

寒冷地湿地帯のセルロース分解機構を明らかにする目的で，北海道の湿地帯 17 か所の底泥のセルロース分解活性を測定した。その結果，泥炭湿地が特に活性が高く，海跡湖，河口域の順に活性は低下した。活性の定性分析の結果，メグマ沼（泥炭湿地），野付湾（泥炭湿地）及びウトナイ湖（海跡湖）では微生物が，長節湖（海跡湖）ではメイオベントスが分解に関わっていることが示された。以上の結果から，寒冷地湿地帯底泥のセルロース分解には微生物がやメイオベントス由来のセルラーゼが重要な働きを果たしていることが示唆された。

キーワード：寒冷地，菌類，湿地帯，セルロース，セルラーゼ，底泥，北海道，メイオベントス

Fig.1




Fig.2




Fig.3




Fig.4




Table 1 Comparison of cellulose activities among wetlands in Hokkaido. Cellulase activity was determined by the quantitative assay as described in the text

Site	Wetland	Location	Geological type	Cellulase activity (nmol/gh) ^a	Organic component ratio (%) ^a	Salinity (‰)
A	Meguma Pond	45°24' N 141°49' E	peat fen	737.88 ± 35.69	66.62	0
B	Notsuke Gulf	43°61' N 145°27' E	peat fen	92.39 ± 0.79	16.85	26
C	Onuma Pond	45°23' N 141°46' E	lagoon	6.74 ± 1.28	0.96	9
D	Lake Kuccharo	45°13' N 142°25' E	lagoon	6.31 ± 0.29	1.07	14
E	Lake Saroma	44°08' N 143°57' E	lagoon	28.48 ± 0.66	1.48	25
F	Lake Notoro	44°06' N 144°10' E	lagoon	13.86 ± 0.81	1.84	23
G	Lake Abashiri	43°59' N 144°13' E	lagoon	2.80 ± 0.26	0.78	0
H	Lake Furen	43°18' N 145°19' E	lagoon	4.22 ± 0.69	16.68	17
I	Mochirippu Pond	43°01' N 145°01' E	lagoon	4.31 ± 0.35	6.65	26
J	Lake Akkeshi	43°03' N 144°51' E	lagoon	21.42 ± 1.11	6.45	20
K	Pashikuru Pond	42°92' N 144°00' E	lagoon	6.65 ± 1.32	0.65	0
L	Lake Chobushi	42°65' N 143°61' E	lagoon	1.58 ± 0.23	1.69	3
M	Lake Utonai	42°70' N 141°70' E	lagoon	44.45 ± 2.00	1.49	0
N	Teshio River	44°54' N 141°43' E	estuary	5.88 ± 0.50	1.04	0

O	Ishikari River	43°15' N 141°22' E	estuary	2.58 ± 0.58	1.23	2
P	Mukawa River	42°33' N 141°55' E	estuary	0	1.41	0
Q	Saru River	42°30' N 142°00' E	estuary	0	1.48	0

^a Cellulase activity and organic component ratio showed a strong positive correlation ($r = 0.96$). p value was calculated as 8.78×10^{-10} , which was statistically significant ($P < 0.01$). Thus, null hypothesis that the coefficient is zero is completely excluded.

Table 2 Composition of grain size of 17 wetlands in Hokkaido

Site	Wetland	Composition by weight of grain size (%)				
		>1 mm	1 mm- 500 µm	500 µm- 250 µm	250 µm - 63 µm	63 µm>
A	Meguma Pond	ND ^a	ND ^a	ND ^a	ND ^a	ND ^a
B	Notsuke Gulf	14.34	11.12	40.38	9.58	24.58
C	Onuma Pond	46.36	19.40	25.54	7.16	1.54
D	Lake Kuccharo	40.30	10.34	25.54	21.98	1.84
E	Lake Saroma	26.90	57.60	12.58	1.50	1.42
F	Lake Notoro	6.96	39.60	37.18	15.42	0.84
G	Lake Abashiri	18.15	47.76	26.71	7.38	0
H	Lake Furen	40.94	23.96	24.98	8.66	1.46
I	Mochirippu Pond	15.58	36.64	36.96	10.24	0.58
J	Lake Akkeshi	18.80	28.26	22.24	25.40	5.30
K	Pashikuru Pond	11.38	10.10	39.28	38.76	0.48
L	Lake Chobushi	76.50	13.60	7.15	2.75	0
M	Lake Utonai	58.40	28.64	8.39	4.56	0
N	Teshio River	9.44	35.61	48.17	6.78	0
O	Ishikari River	0	2.35	88.06	9.59	0
P	Mukawa River	25.40	48.74	12.06	13.38	0.42
Q	Saru River	2.08	10.90	64.92	21.46	0.64

^a ND: not determined